



This is an accepted version of the following published document:

Nava, V., Chandra, S., Aherne, J. *et al.* Plastic debris in lakes and reservoirs. *Nature* **619**, 317–322 (2023). <https://doi.org/10.1038/s41586-023-06168-4>

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's [AM terms of use](#), but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <https://doi.org/10.1038/s41586-023-06168-4>



UNIVERSIDADE DA CORUÑA

Plastic debris in lakes and reservoirs

Author list:

1. Dr. Veronica Nava* - University of Milano-Bicocca, Milano, Italy.
2. Dr. Sudeep Chandra – University of Nevada, Reno, Global Water Center and Biology Department, Reno, NV, USA.
3. Dr. Julian Aherne - Trent University, School of Environment, Ontario, Canada.
4. Dr. María B. Alfonso - Research Institute for Applied Mechanics, Kyushu University, Japan.
5. Prof. Ana M. Antão-Geraldes - Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Bragança, Portugal; Laboratório Associado para a Sustentabilidade e Tecnologia em Regiões de Montanha, Portugal.
6. Dr. Katrin Attermeyer - WasserCluster Lunz - Biologische Station, Lunz am See, Lower Austria, Austria; University of Vienna, Department of Functional and Evolutionary Ecology, Vienna, Austria.
7. Prof. Roberto Bao - Centro Interdisciplinar de Química e Biología (CICA), GRICA group, University of A Coruña, Spain.
8. Dr. Mireia Bartrons - University of Vic - Central University of Catalonia, Biosciences Department, Vic, Spain.
9. Dr. Stella A. Berger - Leibniz Institute of Freshwater Ecology and Inland Fisheries, Department of Plankton and Microbial Ecology, Stechlin, Germany.
10. Dr. Marcin Biernaczyk - West Pomeranian University of Technology in Szczecin, Faculty of Food Science and Fisheries, Szczecin, Poland.
11. Dr. Raphael Bissen - Chulalongkorn University, Department of Mining and Petroleum Engineering, Bangkok, Thailand.
12. Prof. Justin D. Brookes - The University of Adelaide, School of Biological Science, North terrace, Adelaide, Australia.
13. David Brown - Horizons Regional Council, Department of Environmental Data, Palmerston North, New Zealand.
14. Dr. Miguel Cañedo-Argüelles - FEHM Lab, Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Jordi Girona, Barcelona, Spain.
15. Prof. Moisés Canle - Cátedra EMALCSA-UDC, React! Group, Faculty of Sciences & CICA, University of A Coruña, Spain.
16. Dr. Camilla Capelli – SUPSI, Institute of Earth Sciences, Mendrisio, Switzerland.
17. Dr. Rafael Carballeira – Centro Interdisciplinar de Química e Biología (CICA), GRICA group, University of A Coruña, Spain; Cavanilles Institute for Biodiversity and Evolutionary Biology, University of Valencia, Spain.
18. Mr. José Luis Cereijo – Water & Environment Research Group, University of A Coruña, Spain.
19. Dr. Sakonvan Chawchai – Chulalongkorn University, Department of Geology, Bangkok, Thailand.
20. Mr. Søren T. Christensen – CPH Business Laboratory and Environment, Hillerød, Denmark.
21. Professor Kirsten S. Christoffersen – University of Copenhagen, Department of Biology, Copenhagen, Denmark.
22. Dr. Elvira de Eyto – Marine Institute, Furnace, Newport, Co. Mayo, Ireland.
23. Prof. Jorge Delgado – Water & Environment Research Group, University of A Coruña, Spain.
24. Mr. Tyler N. Dornan – The University of Adelaide, School of Biological Science, North terrace, Adelaide, Australia.
25. Dr. Jonathan P. Doubek – Lake Superior State University, School of Natural Resources & Environment, Sault Sainte Marie, Michigan, USA; Lake Superior State University, Center for Freshwater Research and Education, Sault Sainte Marie, Michigan, USA.
26. Ms. Julia Dusaucy – Savoie Mont Blanc University, CNRS, Univ. Grenoble Alpes, EDYTEM, Chambéry, France.

27. Dr. Oxana Erina - Lomonosov Moscow State University, Department of Hydrology, Moscow, Russian Federation; Moscow State University of Technologies and Management (FCU), Faculty of Biotechnology and Fisheries, Moscow, Russian Federation.
28. Dr. Zeynep Ersoy - Universitat de Barcelona, Facultat de Biologia, Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals, FEHM-Lab (Freshwater Ecology, Hydrology and Management), Barcelona, Spain; Universitat de Barcelona, Institut de Recerca de la Biodiversitat (IRBio), Barcelona, Spain; Universidade de Évora, Mediterranean Institute for Agriculture, Environment and Development (MED), Rui Nabeiro Biodiversity Chair, Évora, Portugal.
29. Dr. Heidrun Feuchtmayr – UK Centre for Ecology & Hydrology, Lake Ecosystems Group, Lancaster, United Kingdom.
30. Prof. Maria Luce Frezzotti - University of Milano-Bicocca, Department of Earth and Environmental Sciences, Italy.
31. Dr. Silvia Galafassi – Water Research Institute, National Research Council, Verbania, Italy.
32. Dr. David Gateuille - Savoie Mont Blanc University, CNRS, Univ. Grenoble Alpes, EDYTEM, Chambéry, France.
33. Prof. Vitor Gonçalves - CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Pólo dos Açores - Faculdade de Ciências e Tecnologias, Universidade dos Açores, Ponta Delgada, São Miguel, Açores; BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, Vairão, Portugal.
34. Prof. Hans-Peter Grossart - Leibniz Institute of Freshwater Ecology and Inland Fisheries, Department of Plankton and Microbial Ecology, Stechlin, Germany; Potsdam University, Institute for Biochemistry and Biology, Potsdam, Germany.
35. Prof. David P. Hamilton – Australian Rivers Institute, Griffith University, Nathan, Queensland, Australia.
36. Dr. Ted D. Harris – Kansas Biological Survey and Center for Ecological Research, University of Kansas, Lawrence, Kansas, USA.
37. Dr. Külli Kangur – Estonian University of Life Sciences, Tartu, Estonia.
38. Dr. Gökben Başaran Kankılıç - Kırıkkale University, Department of Biology, Türkiye.
39. Rebecca Kessler – Kansas Biological Survey and Center for Ecological Research, University of Kansas, Lawrence, Kansas, USA.
40. Dr. Christine Kiel – Leibniz Institute of Freshwater Ecology and Inland Fisheries, Department of Plankton and Microbial Ecology, Stechlin, Germany.
41. Dr. Edward M. Krynak - University of Nevada, Reno, Global Water Center and Biology Department, Reno, NV, USA.
42. Dr. Àngels Leiva-Presa – University of Vic - Central University of Catalonia, Biosciences Department, Vic, Spain.
43. Prof. Fabio Lepori – SUPSI, Institute of Earth Sciences, Mendrisio, Switzerland.
44. Dr. Miguel G. Matias – Museo Nacional de Ciencias Naturales, CSIC, Spain; Universidade de Évora, Mediterranean Institute for Agriculture, Environment and Development (MED), Rui Nabeiro Biodiversity Chair, Évora, Portugal.
45. Dr. Shin-ichiro S. Matsuzaki – National Institute for Environmental Studies, Biodiversity Division, Tsukuba, Ibaraki, Japan.
46. Dr. Yvonne McElarney – Agri-Food & Biosciences Institute, Fisheries & Aquatic Ecosystems, Belfast, Northern Ireland.
47. Prof. Beata Messyaszyk – Adam Mickiewicz University, Institute of Environmental Biology, Department of Hydrobiology, Poznań, Poland.
48. Mr. Mark Mitchell – Horizons Regional Council, Department of Science and Innovation, Palmerston North, New Zealand.
49. Mr. Musa C. Mlambo – Albany Museum, Department of Freshwater Invertebrates, Grahamstown, Eastern Cape, South Africa.
50. Dr. Samuel N. Motitsoe – Rhodes University, Department of Zoology and Entomology, Makhanda (Grahamstown) South Africa.
51. Dr. Sarma Nandini – Universidad Nacional Autónoma de México, Mexico.

52. Ms. Valentina Orlandi - University of Milano-Bicocca, Milano, Italy.
53. Ms. Caroline Owens – University of California, Santa Barbara, Department of Ecology, Evolution, and Marine Biology, USA.
54. Dr. Deniz Özkundakci – Environmental Research Institute - Te Pūtahi Rangahau Taiao, The University of Waikato, Hamilton, New Zealand.
55. Ms. Solvig Pinnow – Leibniz Institute of Freshwater Ecology and Inland Fisheries, Department of Plankton and Microbial Ecology, Stechlin, Germany.
56. Prof. Agnieszka Pocięcha – Institute of Nature Conservation, Polish Academy of Sciences, Department of Freshwater Biology, Kraków, Poland.
57. Dr. Pedro Raposeiro – CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Pólo dos Açores - Faculdade de Ciências e Tecnologias, Universidade dos Açores, Ponta Delgada, São Miguel, Açores. BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, Vairão, Portugal.
58. Dr. Eva-Ingrid Rõõm – Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Chair of Hydrobiology and Fishery Kreutzwaldi, Tartumaa, Estonia.
59. Ms. Federica Rotta – SUPSI, Institute of Earth Sciences, Mendrisio, Switzerland.
60. Dr. Nico Salmaso – Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all'Adige, Trento, Italy.
61. Dr. SSS Sarma – Universidad Nacional Autónoma de México, Mexico.
62. Dr. Davide Sartirana – University of Milano-Bicocca, Milano, Italy.
63. Dr. Facundo Scordo – Instituto Argentino de Oceanografía, Universidad Nacional del Sur (UNS)-CONICET, Bahía Blanca, Argentina. 61.bis Departamento de Geografía y Turismo, Universidad Nacional del Sur, Bahía Blanca, Buenos Aires, Argentina.
64. Prof. Claver Sibomana – Center for Research in Natural Sciences and Environment, Faculty of Sciences, University of Burundi, Bujumbura, Burundi.
65. Mr. Daniel Siewert – Wdecki Park Krajobrazowy, Osie, Poland.
66. Prof. Katarzyna Stepanowska – West Pomeranian University of Technology in Szczecin, Faculty of Food Science and Fisheries, Szczecin, Poland.
67. Prof. Ülkü Nihan Tavşanoğlu – Çankırı Karatekin University, Department of Biology, Çankırı, Türkiye.
68. Ms. Maria Tereshina – Lomonosov Moscow State University, Department of Hydrology, Moscow, Russian Federation.
69. Mr. James Thompson – Agri-Food & Biosciences Institute, Fisheries & Aquatic Ecosystems, Belfast, Northern Ireland; Ulster University, School of Geography and Environmental Sciences, Coleraine, Northern Ireland.
70. Dr. Monica Tolotti – Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all'Adige, Trento, Italy.
71. Dr. Amanda Valois – National Institute of Water and Atmospheric Research, Freshwater Ecology, Hamilton and Wellington, New Zealand.
72. Dr. Piet Verburg – National Institute of Water and Atmospheric Research, Freshwater Ecology, Hamilton, New Zealand.
73. Ms. Brittany Welsh – Trent University, School of Environment, Ontario, Canada.
74. Mr. Brian Wesolek – Bay Mills Indian Community, Biological Services Department, Brimley, Michigan, USA.
75. Prof. Gesa A. Weyhenmeyer – Uppsala University, Dept. Ecology and Genetics/Limnology, Uppsala, Sweden.
76. Prof. Naicheng Wu – Department of Geography and Spatial Information Techniques, Ningbo University, Ningbo, China.
77. Prof. Edyta Zawisza – Institute of Geological Sciences, Polish Academy of Sciences, Warsaw, Poland.
78. Ms. Lauren Zink – University of Lethbridge, Department of Biological Sciences, Lethbridge, Alberta, Canada.
79. Prof. Barbara Leoni - University of Milano-Bicocca, Milano, Italy.

* email corresponding author: veronica.nava@unimib.it

Abstract

Plastic debris is thought to be widespread in freshwater ecosystems globally. We sample surface waters of 38 lakes and reservoirs, distributed across gradients of geographical position and limnological attributes, with the aim to identify factors associated with an increased observation of plastics. We find plastic debris in all studied lakes and reservoirs, suggesting that these ecosystems play a key role in the plastic pollution cycle. Our results indicate that two types of lakes are particularly vulnerable to plastic contamination; lakes and reservoirs in densely populated and urbanized areas; and large lakes and reservoirs with elevated deposition areas, long water retention times, and high levels of anthropogenic influence. Plastic concentrations vary widely among lakes; in the most polluted concentrations reach or even exceed those reported in the subtropical oceanic gyres, marine areas collecting large amounts of debris¹. Our findings highlight the importance of including lakes and reservoirs when addressing plastic pollution, in the context of pollution management and for the continued provision of lake ecosystem services.

Main

Plastic pollution is among the major challenges of our time^{2,3}. Marine environments are considered to be the final receptors and sinks of plastic debris⁴, with most research focusing on the impacts of plastics in these systems. However, the spotlight on marine ecosystems has overshadowed the role of freshwaters, particularly lakes and reservoirs, as key components in global plastic pathways. Freshwaters may accumulate plastics at rates similar to, or higher than those in marine systems⁵⁻⁷. Nonetheless, global research on plastic debris in freshwaters has been hindered by two main challenges. First, studies have focused on a limited number of freshwater systems in restricted geographical regions⁸. Second, a lack of standardised sampling methods has prevented direct quantitative comparisons across studies⁹. So far, results from plastic research in lakes and reservoirs have been compared and synthesised through meta-analyses and reviews, which have acknowledged that comparability is limited due to the different methods used across studies^{10,11}. No single study has yet been extended to a global scale, nor has there been any attempt to identify and quantify the drivers of plastic pollution in lentic systems at this scale.

We addressed these fundamental gaps by conducting a globally coordinated study aimed at (1) assessing the occurrence and abundance of plastic debris in surface waters of freshwater lakes and reservoirs, and (2) identifying natural and anthropogenic landscape factors associated with the occurrence of plastic debris. Specifically, we evaluated the ‘signature’ (i.e., abundance and types) of plastics in lakes and reservoirs as a function of potential sources of contamination, and of the hydro-morphological and watershed features that can affect plastic distribution. We performed standardised sampling and analysis of plastics (>250 µm) in surface waters from 38 lakes and reservoirs (hereafter lakes) in 23 countries, covering a wide range of hydro-morphological and anthropogenic impact features.

The study sites, albeit concentrated in the northern hemisphere, are spread out geographically and encompass a wide gradient of lake features and catchment characteristics (Extended Data Fig. 1). As a result, the sample of study sites is representative of global lake variability in several key characteristics¹² (Extended Data Fig. 1). In the absence of any concerted effort or feasible method to obtain a globally representative sample of >100 million lakes, this gradient approach was considered to be the most suitable for this coordinated international effort to study microplastics in lentic systems. We included lakes spanning 0.04 to 32,600 km² in size and 1 to 1,470 m in depth, with population densities of 0 to 3,411 inhabitants km⁻² and urban land cover of 0 to 98% in their watersheds. As we performed a snapshot sampling, the temporal and spatial variation of plastic abundance was not included. However, snapshot sampling events such as this provide valuable information, covering environmental gradients across space¹³. All samples were collected by horizontal trawls of a plankton net perpendicular to the lake outflow, following the same protocol. The concentrated sample was subsequently treated with hydrogen peroxide (concentration 15% for 24 h at 60°C) to reduce adhered substances and organisms on the plastic particles. A total of 9,425 plastic particles were identified and classified based on shape, colour, and size. Polymer composition was identified on a subset of 2,295 (~25%) particles using micro-Raman spectroscopy following Kedzierski *et al.*¹⁴ (see Methods). We related the occurrence, abundance, and features of plastics to variables describing hydro-morphology (e.g., area, depth, shoreline length, and residence time) and anthropogenic impact (e.g., land cover, presence of wastewater treatment facilities, and population density) affecting each lake. We used a geographical information system to delineate the watershed of each lake and derive information about human impact. Regression tree and redundancy analyses (RDA) were used to identify the predictors of concentration and features of plastics.

Plastic debris occurred in all 38 lakes; however, the plastic signature differed greatly among systems¹⁵. Concentration of plastics spanned four orders of magnitude, from 10^{-3} to 10^1 particles m^{-3} (mean \pm standard error, S.E. = 1.82 ± 0.37 particles m^{-3} ; median = 0.85 particles m^{-3} ; Fig. 1a-b). Most of the study sites (55%, 21 lakes) had concentrations below 1 particle m^{-3} , whereas 14 (37%) had concentrations between 1 and 5 particles m^{-3} , and three (8%) had concentrations higher than 5 particles m^{-3} .

The results for the latter three lakes are remarkable because they show that plastic concentrations in lentic systems can even exceed those detected in some of the ocean's most impacted locations. The greatest concentrations in our study (i.e., Lake Lugano with 11.5 particles m^{-3} , Lake Maggiore with 8.2 particles m^{-3} , and Lake Tahoe with 5.4 particles m^{-3}) are higher than those observed in the subtropical oceanic gyres, which are currently considered some of the greatest plastic accumulation zones in the world¹. Maximum concentrations detected in ocean gyres using a similar sampling protocol to this study (trawl nets with a 333 μm mesh) were 1.62 particles m^{-3} in the North Atlantic subtropical gyre¹⁶. It should be noted that we focused on particles larger than 250 μm ; if we included plastics with lower size limits, the concentrations would have been greater than the maximum observed here. Previous studies have identified even higher plastic concentration in some lakes (e.g., Lake Poyang, China: 5,000–34,000 particles m^{-3} ¹⁷). However, samples with such high concentrations are usually collected using a grab method. While this grab method has the advantage of capturing micro- and nano-scale plastics, the small sample volume may result in higher variability in plastic concentrations, and therefore, unrepresentative characterisation of the diversity of plastics compared to methods similar to ours, where nets were used to filter an average of 140 m^3 of lake water per site¹⁸.

The variability among the three replicates (i.e., trawls) collected within each lake was generally low (mean S.E. value = 0.47 particles m^{-3}), especially for lakes with a low average plastic

concentration (mean S.E. value = 0.14 particles m⁻³, see Extended Data Fig. 2). Small scale (among-replicate) variation in plastic distribution may arise from hydrodynamic processes¹⁹, lake morphology, and topography²⁰. Within-lake variability may cause uncertainty when estimating the overall plastic concentration, but the substantial lakewater volume filtered at each site enabled us to obtain reliable average concentrations and capture spatial differences of interest.

Plastic signatures differed depending on the morphometric characteristics of lakes. As shown by cluster analysis (Extended Data Fig. 3), we observed a higher percentage of fibres (mean ± S.E. = 77 ± 0.6%), mainly black or blue and composed of polyester (mean ± S.E. = 39 ± 1.6%), in lakes with comparatively small surface area, volume, maximum depth, and shoreline length. In large, deep lakes with more extensive shorelines, the plastic signature was dominated by fragments (mean ± S.E. = 53 ± 0.9%), mainly transparent and white, with a polymer composition of polypropylene (mean ± S.E. = 35 ± 1.5%) and polyethylene (mean ± S.E. = 31 ± 1.8%). The difference between the two clusters of lakes was statistically significant for surface area, volume, maximum depth, and shoreline length (Wilcoxon test p-values all <0.01). Most plastics from all sites were classified as microplastics (<5 mm, 93.8%). Only 4.7% were mesoplastics (5-10 mm), and we observed very few macroplastics (>10 mm, 1.5%). More than 90% of the plastic particles belonged to two shape categories (Fig. 1c), fibres (49.4%) and fragments (41.0%). Fibres, which were widespread, dominated mainly where the total plastic concentration was low (<1 particle m⁻³). We found textile fibres even in lakes and reservoirs located in remote areas with limited human presence, such as Avery Lake in Michigan, USA. While atmospheric deposition may be a relatively important source of fibres for pristine systems²¹, inputs from tributaries are likely to be more important for lakes and reservoirs with a greater human presence in their catchments. It is well established that fibres from textile materials are a major source of plastic contamination²²; more than 700,000 fibres can be

released into the water system from the washing of 6 kg of laundry²³. Moreover, recreational activities such as swimming can increase direct input of fibres into aquatic systems²⁴.

The fact that fibres and fragments were the most abundant types of microplastics is consistent with field evidence suggesting that secondary microplastics (i.e., particles that result from the fragmentation of larger plastic items) are common in the aquatic environment²⁵. Pellets and spheres, whose shape suggests a primary origin, accounted for less than 1% of plastics (Fig. 1c). Indeed, primary microplastics, which are produced either for indirect use as precursors (nurdles or virgin resin pellets) or for direct use, such as in cosmetics, scrubs and abrasives, are generally less abundant in aquatic systems and are expected to decrease in concentration, at least in some countries, as a result of regulatory measures on single-use plastics²⁶. The relevance of secondary plastic pollution emphasises the need to focus mitigation on preventing plastics from entering waterways or removing them before degradation occurs²⁷.

Recording the shape and other characteristics of plastics helps not only to identify the possible sources of pollution, but also to characterise the impact of plastic pollution. Different shape-dependent impacts are reported in the literature^{28,29}. For instance, up to 10-fold greater adverse effects of fibres compared to beads have been observed on the freshwater zooplankton *Ceriodaphnia dubia* Richard, 1894, with reduced reproductive output at fibre concentrations higher than environmental levels³⁰. Particle size is even more critical in influencing both the toxicokinetics and toxicodynamics of plastics, highlighting the importance of considering size when evaluating the potential risks associated with microplastic exposure³¹. Colour can also affect the toxicity of plastics, and selective feeding on different colours of microplastics has been observed in fish and other organisms because plastics can be mistaken for food of similar coloration³². Manual colour assignment during sample analysis can be difficult due to weathering of particles, and different colour perception among researchers³³. Nonetheless, it is still recommended to record particle colour during visual assessments. Whilst source derivation

is not likely possible based on colour alone, recording colour may help to identify broad trends, such as ingestion preference. In our study, the most common colour was black (30%), followed by transparent (24%), blue (18%), and white (13%). The remaining colours were present in lower abundances (<5%). We found a considerable number of bluish plastics, contrasting with the very low number of reddish ones (i.e., red, orange). Similarly, Martí et al.³⁴, analysing marine plastic items collected in the five subtropical gyres and semi-enclosed regions, reported that white, transparent/translucent, black/grey, and blue particles were particularly common (31%, 16%, 12%, and 11% of the total, respectively), with a very low number of red particles. A proposed explanation for the difference in colour prevalence, apart from different proportions in the waste stream cycle, is that blue is a camouflage colour in aquatic systems (i.e., crypsis mechanisms). Non-blue items (e.g., red items) would have a higher probability of detection and ingestion by visual predators resulting in a progressive enrichment in blue plastic debris³⁴.

Polyester (PES), polypropylene (PP), and polyethylene (PE) constituted the majority of polymers identified, with a mean percentage (\pm S.E.) of $30.4 \pm 3.3\%$, $20.3 \pm 2.9\%$ and $15.7 \pm 2.9\%$, respectively (Fig. 1e). This is not surprising as PE and PP account for more than half of global plastic production (36% and 21%, respectively), while PES, most of which is polyethylene terephthalate (PET), accounts for 70% of all polyester, polyamide, and acrylic fibres production³⁵. The dominance of these polymers is in agreement with previous observations in marine³⁶ and freshwater ecosystems^{37,38} and likely reflects their use in short life-cycle and mass produced products.

Population density and surface area of lakes were the most important predictors of plastic signature, as highlighted by our regression tree analysis (Fig. 2a). The tree with the lowest cross-validated relative error had two splits and three terminal nodes. The first split

differentiated lentic systems with surface area greater than 213 km² (terminal node 1). Lakes with lower surface area were not split further, and the mean plastic concentration for these systems was 4.1 particles m⁻³. Sites with higher surface area were then split based on watershed population density, giving terminal node 2 (lakes > 25 inhabitants km⁻², mean plastic concentration = 1.6 particles m⁻³) and terminal node 3 (lakes < 25 inhabitants km⁻², mean plastic concentration = 0.4 particles m⁻³).

The positive association between urban-related watershed attributes and abundance of plastic debris has been widely observed^{39,40}. In studies where this correlation has not been identified, it has been hypothesised that the study design encompassed a limited number of sites or included sites representing only population density extremes^{5,41}. Lakes were divided based on surface area, highlighting that high concentrations of plastic were found in sites with larger surface area. There may be several explanations for this observation. Larger lakes and reservoirs are usually associated with larger watersheds, greater water inflows, greater shoreline length, and more shoreline development⁴², which implies a larger deposition area and a greater number of point and nonpoint (including atmospheric deposition) sources of contamination. Large lakes are particularly exposed to anthropogenic stressors, as cities and other urban developments are usually widespread on the shores of large lakes, and they may receive inputs from larger and more polluted rivers⁴². In addition, larger lakes have longer residence times than smaller, shallower systems⁴³. For instance, in Lake Tahoe, which has a hydraulic residence time of approximately 650 years⁴⁴, we recorded one of the highest concentrations of plastics (i.e., 5.4 particles m⁻³). Lakes have the potential to act as “traps” and accumulate substantial amounts of plastic debris over time⁴⁵. Additional research is warranted that addresses how the landscape position of lakes and lake characteristics affect microplastic abundance, their fate within lakes due to sedimentation, or transport within from nearshore to offshore habitats, and out of the ecosystem through outflows.

Concentration and features of plastics varied based on environmental factors representing the impact of human activity and morphometric lake characteristics. Based on redundancy analysis, these environmental variables explained 55.9% of the variation in plastic concentration across sampling sites. Lakes with lower plastic concentration were clustered, while lakes with high plastic concentration were more spread out in ordination space (scaling 1, Extended Data Fig. 4), suggesting plastic features were more similar in less polluted lakes. There were two categories of lakes with high plastic concentration (scaling 2, Fig. 2b). The first group was characterised by a high number of wastewater treatment plants and high surface area, depth, volume and shoreline length, and a high abundance of fragments. The second group was characterised by high human impact (human footprint, population density, and urban land cover), and high abundance of fibres. The remaining sites with lower plastic concentrations (<1 particle m^{-3}) had negative relationships with measured anthropogenic variables and seemed weakly related to presence of cropland, which in turn was negatively related to urban land cover and population density.

Our analysis indicated that two types of lakes are particularly vulnerable to plastic contamination: (1) those located within highly urbanised and populated watersheds, and (2) those with high surface area, where we found high surface concentrations of plastic fragments, due to their elevated drainage area and long retention time. This result is particularly relevant because the elevated concentration of plastics in lakes with large water volumes implies high plastic loading, which could impact their ecosystem services locally, regionally and globally⁴².

Given the relatively high concentration of plastic debris, particularly in large lakes and reservoirs, lakes may be 'sentinels of plastic pollution' because they act as collectors and integrators of different sources of plastics from the watersheds and atmosphere, and 'active pipes' as they may retain, process and transport plastics through watersheds to the oceans. We find that plastic concentrations in freshwater ecosystems can be higher than those in marine

ecosystems, which are generally considered final receptors of plastic debris. This underpins the relevance of lakes as key components in the global ‘plastic cycle’. Optimising management policies to mitigate plastic pollution in upstream freshwater lakes is therefore essential to prevent plastics from entering waterways²⁷, and ending up in marine systems.

Some of the lakes most contaminated with plastic debris, including lakes Maggiore (IT), Lugano (CH-IT), Tahoe (USA), and Neagh (UK), are important sources of drinking water for local populations and support important recreational based economies. The proportion of plastic debris that ends up in the water supply is unknown, but we suggest that the potential contamination of microplastics in drinking water should become a global management and research priority.

In addition to contaminating the water supply for human needs, plastic pollution has detrimental effects on aquatic organisms and ecosystem functioning. Detecting the concentration of plastics is possible through the methods employed in this study but understanding their fate and ecological impacts remains an important and novel area of research. For instance, plastics at the surface of aquatic systems can aid the release of methane and other greenhouse gases, demonstrating that the effects of plastics can span across ecosystem boundaries^{46,47}. Plastics can go beyond the hydrosphere and interact with the atmosphere, biosphere and lithosphere, potentially affecting biogeochemical cycles through mechanisms that still need to be understood. These multiple and potentially synergistic effects call for a holistic assessment of plastic pollution in lentic systems.

Furthermore, our findings indicate that microplastics occur even in lakes that are not subjected to direct anthropogenic impacts. Microplastics, therefore, add a new stressor to these lakes and the organisms that live within them, which already face a range of pressures, including climate change⁴⁸, salinization⁴⁹, increased nutrient deposition, and nearshore filamentous algal blooms⁵⁰, to name a few. Therefore, even in remote areas away from direct human pressure,

no lake can be considered to be truly 'pristine' with respect to plastic pollution. These results demonstrate the global reach of plastic pollution and serve as yet another reminder of the unfortunate and indelible signature of humanity on lakes.

References

1. Eriksen, M., Thiel, M. & Lebreton, L. Nature of Plastic Marine Pollution in the Subtropical Gyres. *Hazardous Chemicals Associated with Plastics in the Marine Environment* 135–162 (2016).
2. Rochman, C. M. & Hoellein, T. The global odyssey of plastic pollution. *Science (1979)* **368**, 1184–1185 (2020).
3. MacLeod, M., Arp, H. P. H., Tekman, M. B. & Jahnke, A. The global threat from plastic pollution. *Science (1979)* **373**, 61–65 (2021).
4. Waldschläger, K., Lechthaler, S., Stauch, G. & Schüttrumpf, H. The way of microplastic through the environment – Application of the source-pathway-receptor model. *Science of The Total Environment* **713**, 136584 (2020).
5. Talbot, R. & Chang, H. Microplastics in freshwater: A global review of factors affecting spatial and temporal variations. *Environmental Pollution* **292**, 118393 (2022).
6. Horton, A. A., Walton, A., Spurgeon, D. J., Lahive, E. & Svendsen, C. Microplastics in freshwater and terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research priorities. *Science of the Total Environment* **586**, 127–141 (2017).
7. Ockenden, A., Tremblay, L. A., Dikareva, N. & Simon, K. S. Towards more ecologically relevant investigations of the impacts of microplastic pollution in freshwater ecosystems. *Science of The Total Environment* **792**, 148507 (2021).
8. Blettler, M. C. M., Abrial, E., Khan, F. R., Sivri, N. & Espinola, L. A. Freshwater plastic pollution: Recognizing research biases and identifying knowledge gaps. *Water Res* **143**, 416–424 (2018).
9. Hartmann, N. B. *et al.* Are We Speaking the Same Language? Recommendations for a Definition and Categorization Framework for Plastic Debris. *Environ Sci Technol* **53**, 1039–1047 (2019).
10. Guo, Z. *et al.* Global meta-analysis of microplastic contamination in reservoirs with a novel framework. *Water Res* **207**, 117828 (2021).
11. Li, C., Busquets, R. & Campos, L. C. Assessment of microplastics in freshwater systems: A review. *Science of The Total Environment* **707**, 135578 (2020).
12. Messenger, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nat Commun* **7**, 13603 (2016).
13. Mantzouki, E. *et al.* Snapshot Surveys for Lake Monitoring, More Than a Shot in the Dark. *Front Ecol Evol* **6**, (2018).
14. Kedzierski, M. *et al.* Microplastics in Mediterranean Sea: A protocol to robustly assess contamination characteristics. *PLoS One* **14**, 1–16 (2019).
15. Nava, V. Data for ‘Plastic debris in lakes and reservoirs’. (2023) doi: 10.5281/zenodo.7824882.
16. Courtene-Jones, W., van Gennip, S., Penicaud, J., Penn, E. & Thompson, R. C. Synthetic microplastic abundance and composition along a longitudinal gradient traversing the subtropical gyre in the North Atlantic Ocean. *Mar Pollut Bull* **185**, 114371 (2022).
17. Yuan, W., Liu, X., Wang, W., Di, M. & Wang, J. Microplastic abundance, distribution and composition in water, sediments, and wild fish from Poyang Lake, China. *Ecotoxicol Environ Saf* **170**, 180–187 (2019).
18. Barrows, A. P. W., Neumann, C. A., Berger, M. L. & Shaw, S. D. Grab vs. neuston tow net: a microplastic sampling performance comparison and possible advances in the field. *Anal. Methods* **9**, 1446–1453 (2017).
19. Free, C. M. *et al.* High-levels of microplastic pollution in a large, remote, mountain lake. *Mar Pollut Bull* **85**, 156–163 (2014).
20. Deng, C. *et al.* Impacts of underwater topography on the distribution of microplastics in lakes: A case from Dianchi Lake, China. *Science of The Total Environment* **837**, 155708 (2022).

21. Brahney, J., Hallerud, M., Heim, E., Hahnenberger, M. & Sukumaran, S. Plastic rain in protected areas of the United States. *Science* (1979) **368**, 1257–1260 (2020).
22. Carney Almroth, B. M. *et al.* Quantifying shedding of synthetic fibers from textiles; a source of microplastics released into the environment. *Environmental Science and Pollution Research* **25**, 1191–1199 (2018).
23. Napper, I. E. & Thompson, R. C. Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and washing conditions. *Mar Pollut Bull* **112**, 39–45 (2016).
24. Gao, F. *et al.* The seasonal distribution characteristics of microplastics on bathing beaches along the coast of Qingdao, China. *Science of The Total Environment* **783**, 146969 (2021).
25. Shim, W. J., Hong, S. H. & Eo, S. Marine Microplastics: Abundance, Distribution, and Composition. in *Microplastic Contamination in Aquatic Environments: An Emerging Matter of Environmental Urgency* (ed. Zeng, E. Y.) 1–26 (Elsevier, 2018).
26. Waldman, W. R. & Rillig, M. C. Microplastic Research Should Embrace the Complexity of Secondary Particles. *Environ Sci Technol* **54**, 7751–7753 (2020).
27. Schmaltz, E. *et al.* Plastic pollution solutions: emerging technologies to prevent and collect marine plastic pollution. *Environ Int* **144**, 106067 (2020).
28. Frydkjær, C. K., Iversen, N. & Roslev, P. Ingestion and Egestion of Microplastics by the Cladoceran *Daphnia magna*: Effects of Regular and Irregular Shaped Plastic and Sorbed Phenanthrene. *Bull Environ Contam Toxicol* **99**, 655–661 (2017).
29. Botterell, Z. L. R. *et al.* Bioavailability of Microplastics to Marine Zooplankton: Effect of Shape and Infochemicals. *Environ Sci Technol* **54**, 12024–12033 (2020).
30. Ziajahromi, S., Kumar, A., Neale, P. A. & Leusch, F. D. L. Impact of Microplastic Beads and Fibers on Waterflea (*Ceriodaphnia dubia*) Survival, Growth, and Reproduction: Implications of Single and Mixture Exposures. *Environ Sci Technol* **51**, 13397–13406 (2017).
31. Mehinto, A. C. *et al.* Risk-based management framework for microplastics in aquatic ecosystems. *Microplastics and Nanoplastics* **2**, 17 (2022).
32. Santos, R. G., Andrades, R., Fardim, L. M. & Martins, A. S. Marine debris ingestion and Thayer’s law – The importance of plastic color. *Environmental Pollution* **214**, 585–588 (2016).
33. Lusher, A. L., Bråte, I. L. N., Munno, K., Hurley, R. R. & Welden, N. A. Is It or Isn’t It: The Importance of Visual Classification in Microplastic Characterization. *Appl Spectrosc* **74**, 1139–1153 (2020).
34. Martí, E. *et al.* The Colors of the Ocean Plastics. *Environ Sci Technol* **54**, 6594–6601 (2020).
35. Geyer, R., Jambeck, J. R. & Law, K. L. Production, use, and fate of all plastics ever made. *Sci Adv* **3**, e1700782 (2017).
36. Erni-Cassola, G., Zadjelovic, V., Gibson, M. I. & Christie-Oleza, J. A. Distribution of plastic polymer types in the marine environment; A meta-analysis. *J Hazard Mater* **369**, 691–698 (2019).
37. Dusaucy, J., Gateuille, D., Perrette, Y. & Naffrechoux, E. Microplastic pollution of worldwide lakes. *Environmental Pollution* **284**, 117075 (2021).
38. Li, C., Busquets, R. & Campos, L. C. Assessment of microplastics in freshwater systems: A review. *Science of the Total Environment* **707**, 135578 (2020).
39. Baldwin, A. K., Corsi, S. R. & Mason, S. A. Plastic Debris in 29 Great Lakes Tributaries: Relations to Watershed Attributes and Hydrology. *Environ Sci Technol* **50**, 10377–10385 (2016).
40. Tanentzap, A. J. *et al.* Microplastics and anthropogenic fibre concentrations in lakes reflect surrounding land use. *PLoS Biol* **19**, e3001389 (2021).
41. Dikareva, N. & Simon, K. S. Microplastic pollution in streams spanning an urbanisation gradient. *Environmental Pollution* **250**, 292–299 (2019).
42. Jenny, J.-P. *et al.* Scientists’ Warning to Humanity: Rapid degradation of the world’s large lakes. *J Great Lakes Res* **46**, 686–702 (2020).

43. Messenger, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nat Commun* **7**, 13603 (2016).
44. Jassby, A., Goldman, C., Reuter, J., Richards, R. & Heyvaert, A. Lake Tahoe: Diagnosis and rehabilitation of a large mountain lake. in *The Great Lakes of the World (GLOW)* 431–454 (2001).
45. Windsor, F. M. *et al.* A catchment-scale perspective of plastic pollution. *Glob Chang Biol* **25**, 1207–1221 (2019).
46. Royer, S.-J., Ferrón, S., Wilson, S. T. & Karl, D. M. Production of methane and ethylene from plastic in the environment. *PLoS One* **13**, e0200574- (2018).
47. Sheridan, E. A. *et al.* Plastic pollution fosters more microbial growth in lakes than natural organic matter. *Nat Commun* **13**, 4175 (2022).
48. Woolway, R. I. *et al.* Global lake responses to climate change. *Nat Rev Earth Environ* **1**, 388–403 (2020).
49. Dugan, H. A. *et al.* Salting our freshwater lakes. *Proceedings of the National Academy of Sciences* **114**, 4453–4458 (2017).
50. Vadeboncoeur, Y. *et al.* Blue Waters, Green Bottoms: Benthic Filamentous Algal Blooms Are an Emerging Threat to Clear Lakes Worldwide. *Bioscience* **71**, 1011–1027 (2021).

Figure




Fig. 1: Concentration and features of plastics identified in the 38 lakes and reservoirs.

a, Map showing distribution and concentration of plastics in the 38 lakes and reservoirs included in the study. The dots are coloured based on the concentration of plastics (particles m^{-3}) detected. **b**, Boxplot showing the concentration of plastic (particles m^{-3}) in the 38 lakes and reservoirs. Boxplot statistics: the lower and upper hinges correspond to the first and third quartiles. The upper (lower) whisker extends from the hinge to the largest (smallest) value no further than 1.5 x IQR (Inter Quartile Range) from the hinge. Data beyond the end of the whiskers are outlying points and are plotted with triangular shape. **c**, Donut pie chart with percentage abundances of the different shapes and relative colours for all the plastic particles analysed. **d**, Width distribution (from 0 to 10 mm) of the different particle shapes. **e**, Average percentage \pm standard error (S.E.) of polymer composition for all the plastic particles analysed.




Fig. 2: Relationship of plastic concentration and features with environmental and anthropogenic drivers.

a, Results of regression tree analysis based on the total concentration of plastics in the studied lakes, with three terminal nodes. **b**, Redundancy analysis between plastic concentration in lakes, features of plastics, and environmental and anthropogenic drivers potentially able to explain the distribution and characteristics of plastic contamination (scaling 2). The dots are coloured based on the concentration of plastics (particles m^{-3}) detected.

Methods

Study sites and sample collection

The samples were collected from 38 lakes and reservoirs located in 23 countries distributed in both hemispheres, but with a higher density in the northern hemisphere. The sampled sites represent a wide assortment of limnological conditions (Supplementary Table 1). Surface area ranged from 0.04 to 32,600 km² (median = 19.50 km²), mean depth from 0.5 to 580 m (median = 9.7 m), and volume from 1.8 x 10⁻⁵ to 18,980 km³ (median = 0.18 km³). The lakes spanned different mixing regimes (i.e., polymictic, 11; monomictic, 12; dimictic, 8; and meromictic, 5) and trophic states (i.e., ultra-oligotrophic, 3; oligotrophic, 10; mesotrophic, 12; eutrophic, 11; and hyper-eutrophic, 2).

During 2020–2021, samples of plastic debris were collected following a standardised protocol. The samples were collected from a boat using horizontal net trawls (mesh sizes ranging from 50 to 300 µm) and three replicates were obtained from each lake. Sampling occurred in the pelagic zone, near the main lake outlet, on a calm day to minimise the risk of missing an unknown portion of the sample area, because rough water may cause nets to rise above or below the water's surface. The three trawl transects were oriented perpendicular to the outflow. The net was placed at the port side and the boat speed was maintained at around 1–1.5 m s⁻¹, following GESAMP⁵¹. At least 50 m³ of water was filtered for each trawl and GPS tracks were recorded to estimate the exact volume filtered. In case of net clogging, the trawls were divided into different sub-trawls to allow net cleaning.

Sample analysis

All the samples were analysed at the Laboratory of Freshwater Ecology and Management of the University of Milano-Bicocca (Italy) following a common standardised procedure. The samples were wet-sieved on a 250 µm mesh to align the lower limit size across samples,

because nets with different mesh sizes were used for the sampling work. Then, the samples were treated with 15% H₂O₂ for 24 h at 60°C to eliminate organic matter and organisms adhering to the plastic particles. This procedure was selected to reduce potential damage to plastic particles^{52–54}. The samples were then filtered onto 0.45 µm glass microfibre filters (GF/F, 47 mm Ø, Whatman), which were stored in clean glass Petri dishes. The filters were examined under a dissecting microscope (40x, Heerbrugg WILD M3Z) and particles recognized as plastics were transferred to glass slides for the subsequent spectroscopic analysis. Particles were either accepted or rejected as microplastics based on a catalogue of morphological criteria. Visual classification was considered reliable as a first step since we focused on the larger size fraction of microplastics (>250 µm). If the morphological classification was uncertain, Raman spectra were acquired to confirm or reject the hypothesised classification (for more details see ‘Micro-spectroscopy analyses’ section). Pictures of all plastic particles were taken using a high-resolution camera (Leica ICC50). All particles were counted, and their Feret’s diameter was measured using the software ImageJ (version 1.52q). The largest plastic that we found had a Feret’s diameter equal to 8.6 cm. Based on their dimensions, plastics were assigned to three different size categories: microplastics (250 µm to 5 mm)^{55,56}, mesoplastics (5 to 10 mm), and macroplastics (>10 mm; Extended Data Fig. 5). For shape categorization, we adopted a modified version of the classification proposed by Hartmann *et al.*⁹. In particular, plastics were classified as fibres, fragments, films, spheres/pellets, or lines. A lines class was added to identify those plastics that have a shape similar to fibres (longer in one dimension), but have a larger diameter, to differentiate them from fibres derived from textiles (Extended Data Fig. 6). Moreover, plastic particles were classified based on colour (i.e., red, orange, yellow, green, blue, violet, black, white, transparent, or multi-coloured), following a RAL standard colour scale, according to Lusher *et al.*³³.

Micro-spectroscopy analyses

Raman micro-spectroscopy was used to provide reliable data on the total number of plastics identified in each sample, because visual classification alone is insufficient to determine microplastic abundance³³. The particles for which the result of the visual classification was uncertain were subjected to Raman analysis to decide whether to include or exclude them. Then, to estimate the percent occurrence of the different polymers of plastics collected, Raman micro-spectroscopy was performed on a random subsample of the visually identified microplastic particles, as widely suggested in the literature⁵⁷⁻⁵⁹. In particular, a robust procedure was adopted to first determine the minimum number of particles to be studied to reach a certain confidence level in the estimated proportion of different polymers in the sample, following Kedzierski *et al.*¹⁴. The size of the subsample, n , for each lake was derived as follows:

$$n = \frac{\frac{1}{4} + \frac{\varepsilon^2}{\left(u_{1-\frac{\alpha}{2}}\right)^2}}{\frac{\varepsilon^2}{\left(u_{1-\frac{\alpha}{2}}\right)^2} + \frac{1}{4N}}$$

with ε being the desired accuracy; $u_{1-\alpha/2}$ the fractal of order α of the standardised normal law; and N the total number of plastics counted in each sample. We chose a degree of confidence of 95% (i.e., $\alpha = 0.05$; $u_{1-\alpha/2} = 1.96$) and $\varepsilon = 0.1$.

Misidentification at this stage (after visual analysis and Raman pre-checking) was infrequent ($< 3\% \pm 2\%$ on average among samples). Any particles found to be non-plastic at this stage were removed from the total count, and, when this happened, additional particles equal to the amount removed have been analysed spectroscopically to increase the confidence of the estimate. For textile fibres, anthropogenic fibres (i.e., fibres containing a mixture of dyes and cellulose) were included in the total count in agreement with previously published papers that

highlights their relevance for aquatic toxicity⁶⁰. The category ‘anthropogenic’ was also assigned to fibres when the dye masked the polymer and no information other than the colourants could be obtained. Also in these cases, to improve the accuracy of the estimate, an equal number of extra particles was examined using Raman micro-spectroscopy.

Raman spectra were acquired using a Horiba Jobin Yvon LabRAM HR Evolution Raman System at the Department of Earth and Environmental Sciences, University of Milano – Bicocca (Italy), characterised with 800 mm of focal distance coupled with an air-cooled 1024×256 px CCD detector. The spectra were obtained by using an attenuated green Nd 532.06 nm laser source (300 mW) with a 50× magnification (Olympus BXFM). The grating was 600 g mm⁻¹ and the spectral per pixel resolution was about 1.6 cm⁻¹. Two spectra were acquired for each particle with a spectral interval from 222.86 to 1,899.01 cm⁻¹ and from 1,762.24 to 3,177.02 cm⁻¹. Depending on the particles analysed, the acquisition parameters were changed: accumulation ranged between 1–3, integration time between 20–60 s, and power between 0.3–300 mW. Instrument calibration was performed daily based on the auto-calibration process performed by the Raman System Service with respect to the zero line and the silicon standard (520.7 cm⁻¹), according to the ASTM 1840–96 prescription^{61,62}. Raman spectra were baseline corrected and processed using the Fityk software^{63,64}. Further analyses on polymer spectra were performed in R (4.0.3), using the package ‘RamanMP⁶⁵. The final identification of microplastics was based on individual assessment of each spectrum, by identifying the characteristic bands of the suspected polymer in the sample spectrum.

Quality controls (QA/QC) of plastic abundance data

Prior to sampling, the nets were thoroughly cleaned using ultrapure water and then 5 L of ultrapure water was filtered through the net. This volume was collected to account for possible contamination of the sampling equipment. Laboratory-based quality assurance and quality

control (QA/QC) included procedural blanks. To assess potential contamination from laboratory materials or air, laboratory blanks were also collected. Moreover, all laboratory equipment was rinsed three times with ultrapure water. Glassware equipment was used where possible, and all the surfaces were cleaned before use. In addition, the samples were covered with aluminium foil, and cotton laboratory coats were worn. Plastic particles observed on the blanks were subtracted from the total values in environmental samples by randomly removing particles that matched the colour, shape, and polymeric composition of the blank particles (Extended Data Table 1).

Watershed and lake attributes

Sixteen variables were extracted to characterise the level of anthropogenic impact across the different lakes. We delineated the watershed boundary of all lakes (i.e., the land and water areas that drain toward the lake) using ArcMap 10.7 (Spatial Analyst tool) Geographic Information Systems (GIS) software, and the ASTER Global Digital Elevation Model (GDEM) Version 3 (ASTGTM), which provides a spatial resolution of 1 arc second (approximately 30 m horizontal spacing at the equator)⁶⁶.

Land cover in each watershed was obtained by clipping the 100 m resolution land cover map provided by the Copernicus Global Land Service⁶⁷. Data on watershed population were obtained from the UN WPP-adjusted population estimates for the year 2020⁶⁸. Information about wastewater treatment plants (WWTPs) was retrieved using the HydroWASTE dataset, a spatially explicit global database of 58,502 WWTPs and their characteristics²⁰. Additionally, the mean Global Human Footprint Index was derived for each system. This index is the Human Influence Index (HII) normalised by biome and realm. The HII is a global dataset of 1 km grid cells, created from nine global data layers of human population pressure (population density), human land use and infrastructure (built-up areas, night time lights, and land use/land cover),

and human access (coastlines, roads, railroads, and navigable rivers). A HII value of zero represents the least influenced part of the biome with a value of 100 representing the most influenced⁶⁹.

Data analyses

To identify coherent groups of lakes based on the percentage of occurrence of the different plastic features (shape, colour, and polymer composition), we performed K-means clustering. The optimal number of clusters (2) was determined using the Elbow Method⁷⁰. Statistical differences among lake attributes within the two clusters were evaluated using Wilcoxon rank-sum tests.

We evaluated the association among the 16 explanatory variables through Pearson product-moment correlation tests and variables that were highly correlated ($r > 0.85$, $p < 0.001$) were removed from subsequent analyses to avoid collinearity. Moreover, we used the Variance Inflation Factor technique to remove further variables that were highly correlated⁷¹.

To determine which candidate explanatory variables were associated with variation in the concentration of plastics, we used univariate regression trees⁷². Regression trees use recursive partitioning algorithms to reveal the structure in the data, by successive binary partitions based on the different predictors, in a way that minimises the sum of squares in the concentration of plastic within each group (node). The algorithm functioned by maximising the between-node sum of squares (minimising the within-node sum of squares) and then repeating the procedure until an overly large regression tree was constructed. The dataset was divided into training data (80%) and testing data (20%), and the mean squared error was calculated. The tree was then pruned to avoid overfitting based on the complexity parameter, which is the amount by which splitting a node improved the relative error⁷³.

Redundancy Analysis was then performed to explicitly model response variables (i.e., concentration of the shapes of plastics: fibres, fragments, film and others) as a function of explanatory variables. We reported values based on both scaling 1 and 2: scaling 1 shows similarities between objects in the response matrix, while scaling 2 shows the effects of explanatory variables⁷⁴.

All statistical analyses were completed in R (version 4.2.2), using the packages ‘corrplot’⁷⁵, ‘factoextra’⁷⁶, ‘ggplot2’⁷⁷, ‘rpart’⁷⁸, and ‘vegan’⁷⁹.

Data availability statement

The datasets generated during and/or analysed during the current study are available in the Zenodo repository, 10.5281/zenodo.7824882¹⁵

Methods references

51. GESAMP. *Guidelines or the monitoring and assessment of plastic litter and microplastics in the ocean*. ((IMO/FAO/UNESCO-IOC/UNIDO/WMO/IAEA/UN/UNEP/UNDP/ISA Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection), 2019).
52. Wiggan, K. J. & Holland, E. B. Validation and application of cost and time effective methods for the detection of 3–500 μm sized microplastics in the urban marine and estuarine environments surrounding Long Beach, California. *Mar Pollut Bull* **143**, 152–162 (2019).
53. Zhao, S., Danley, M., Ward, J. E., Li, D. & Mincer, T. J. An approach for extraction, characterization and quantitation of microplastic in natural marine snow using Raman microscopy. *Analytical Methods* **9**, 1470–1478 (2017).
54. Hurley, R. R., Lusher, A. L., Olsen, M. & Nizzetto, L. Validation of a Method for Extracting Microplastics from Complex, Organic-Rich, Environmental Matrices. *Environ Sci Technol* **52**, 7409–7417 (2018).
55. European Chemicals Agency (ECHA). *Annex XV restriction report - proposal for a restriction*. (2019).
56. Coffin, S. *Proposed definition of 'microplastics in drinking water'*. California Water Boards https://www.waterboards.ca.gov/drinking_water/certlic/drinkingwater/docs/stffrprt_jun3.pdf (2020).
57. K  ppler, A. *et al.* Analysis of environmental microplastics by vibrational microspectroscopy: FTIR, Raman or both? *Anal Bioanal Chem* **408**, 8377–8391 (2016).
58. Kazour, M., Jemaa, S., Issa, C., Khalaf, G. & Amara, R. Microplastics pollution along the Lebanese coast (Eastern Mediterranean Basin): Occurrence in surface water, sediments and biota samples. *Science of the Total Environment* **696**, 133933 (2019).
59. Alfonso, M. B. *et al.* First evidence of microplastics in nine lakes across Patagonia (South America). *Science of The Total Environment* **733**, 139385 (2020).
60. Athey, S. N. & Erdle, L. M. Are We Underestimating Anthropogenic Microfiber Pollution? A Critical Review of Occurrence, Methods, and Reporting. *Environ Toxicol Chem* **41**, 822–837 (2022).
61. Hutsebaut, D., Vandenabeele, P. & Moens, L. Evaluation of an accurate calibration and spectral standardization procedure for Raman spectroscopy. *Analyst* **130**, 1204–1214 (2005).
62. Remigi, S., Mancini, T., Ferrando, S. & Frezzotti, M. L. Interlaboratory Application of Raman CO₂ Densimeter Equations: Experimental Procedure and Statistical Analysis Using Bootstrapped Confidence Intervals. *Appl Spectrosc* **75**, 867–881 (2021).
63. Frezzotti, M. L. Diamond growth from organic compounds in hydrous fluids deep within the Earth. *Nat Commun* **10**, 4952 (2019).
64. Wojdyr, M. Fityk: A general-purpose peak fitting program. *J Appl Crystallogr* **43**, 1126–1128 (2010).
65. Nava, V., Frezzotti, M. L. & Leoni, B. Raman Spectroscopy for the Analysis of Microplastics in Aquatic Systems. *Appl Spectrosc* **75**, 1341–1357 (2021).
66. NASA/METI/AIST/Japan Spacesystems and U.S./Japan ASTER Science Team. ASTER Global Digital Elevation Model V003. *NASA EOSDIS Land Processes DAAC*. <https://lpdaac.usgs.gov/products/astgtmv003/> (2019) doi:10.5067/ASTER/ASTGTM.003.
67. Buchhorn, M. *et al.* Copernicus Global Land Service: Land Cover 100m: collection 3: epoch 2019: Globe. (2020) doi:10.5281/ZENODO.3939050.
68. Center for International Earth Science Information Network - CIESIN - Columbia University. Gridded Population of the World, Version 4 (GPWv4): Administrative Unit Center Points with Population Estimates, Revision 11. <https://doi.org/10.7927/H4BC3WMT> (2020).

69. Wildlife Conservation Society, Center for International Earth Science Information Network & Columbia University. Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human Footprint Dataset (Geographic). <https://doi.org/10.7927/H4M61H5F> (2005).
70. Han, J., Kamber, M. & Pei, J. Cluster Analysis: Basic Concepts and Methods. in *Data Mining Concepts and Techniques* (Elsevier, 2011).
71. Zuur, A. F., Ieno, E. N. & Elphick, C. S. A protocol for data exploration to avoid common statistical problems. *Methods Ecol Evol* **1**, 3–14 (2010).
72. James, G., Witten, D., Hastie, T. & Tibshirani, R. *An introduction to statistical learning with applications in R*. (Springer, 2021).
73. Breiman, L., Friedman, J. H., Olshen, R. A. & Stone, C. J. *Classification And Regression Trees*. (Routledge, 1984). doi:10.1201/9781315139470.
74. Zuur, A. F., Ieno, E. N. & Smith, G. M. *Analysing Ecological Data*. Springer Science (2007).
75. Wei, T. & Simko, V. R package ‘corrplot’: Visualization of a Correlation Matrix (Version 0.84). *R package version* (2017).
76. Kassambara, A. & Mundt, F. Factoextra: extract and visualize the results of multivariate data analyses. *R package version* Preprint at (2020).
77. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, 2016). doi:10.1007/978-0-387-98141-3.
78. Therneau, T. & Atkinson, B. rpart: Recursive Partitioning and Regression Trees. Preprint at (2022).
79. Oksanen, J. *et al.* Vegan: Community Ecology Package. R package version 2.0-2. Preprint at (2020).

Acknowledgments

This manuscript benefited from conversations at meetings of the Global Lake Ecological Observatory Network (GLEON; supported by funding from US NSF grants 1137327 and 1702991). This work was supported by the University of Milano-Bicocca (UNIMIB). Raman facilities were provided by the Department of Earth and Environmental Sciences (DISAT – UNIMIB), and the ‘Interdepartmental Network of Spectroscopy’ (UNIMIB). We gratefully acknowledge Giulia Candian and Emma Caprini for their assistance in the laboratory activities and data analysis.

A.M.A.G. acknowledges the Foundation for Science and Technology (FCT, Portugal) for financial support through national funds FCT/MCTES (PIDDAC) to CIMO (UIDB/00690/2020 and UIDP/00690/2020) and SusTEC (LA/P/0007/2020). R.B. acknowledges support from Project IMPACOM (PID2019-107424RB-I00) of the Spanish Ministry of Science and Innovation. M.C.A. was supported by a Ramon y Cajal contract funded by the Spanish Ministry of Science and Innovation (RYC2020-029829-I). M.C. acknowledges support from Cátedra EMALCSA-UDC (industrial chair). R.C. is grateful for a PhD contract from the Xunta de Galicia (Plan IC2) co-financed by European Social Fund (PRE/2013/044). Z.E. and M.G.M. acknowledge support from the Portuguese Science and Technology Foundation (FCT) project no: PTDC/CTA- AMB/30793/2017 (AdaptAlentejo—Predicting ecosystem-level responses to climate change). H.P.G. and S.P. were supported by the European Union Horizon 2020 Research and Innovation 772 programme under Grant Agreement number 965367 (PlasticsFatE). D.P.H. acknowledges support of the Australian Research Council (DP190101848). S.M. acknowledges support from Rhodes University and the University Capacity Development Programme. P.M.R. acknowledges support from Portuguese Science Foundation (FCT) (DL57/2016/ICETA/EEC2018/25). K.K. acknowledges support from grant PRG 1266 of Estonian Research Council. S.N. AND S.S.S.S. acknowledge support PAPIIT UNAM IG200820. E.R. acknowledges support from grant PUT1598 of Estonian Research Council. C.S. acknowledges support from Flemish Interuniversity Council through VLIR-UOS/UB Programme. G.A.W. acknowledges support from the Swedish Research Council (Grant No. 2020-03222) and Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS; Grant No. 2020-01091). N.W.

from the National Natural Science Foundation of China (No. 52279068). F.S. acknowledges support from an IAI-CONICET special grant.

Author contributions

V.N. and B.L. designed the study, and drafted the manuscript; V.N. compiled the data, conducted analyses; V.N., J.A., M.B.A, A.M.A.G., S.A.B., R.B., M.C.A, M.C., C.C., S.C., E.D.E., J.P.D., O.E., Z.E., H.F., S.G., H.P.G., D.P.H., T.D.H., K.K., C.K., A.L.P., M.G.M., M.C.M., S.N., C.O., D.Ö., S.P., F.R., F.S., C.S., U.N.T., P.V., G.A.W., L.Z., B.L. contributed to the discussion and conceptualization of the paper; V.N., J.A., M.B.A, A.M.A.G, K.A., R.B., S.A.B, R.B., M.C, C.C., R.C., J.C., S.C., E.D.E, T.N.D, J.P.D., J.D., O.E., Z.E., H.F., S.G., H.P.G., D.P.H., T.D.H., K.K., G.B.K., B.E.K., C.K., E.M.K., A.L.P., Y.M., M.C.M., S.N., V.O., D.Ö., S.P., A.P., P.M.R., E.I.R., F.R., F.S., C.S., D.S., K.S., K.S.C., U.N.T., M.T., J.T., P.V., G.A.W., N.W., E.Z. collected the samples; M.L.F. provided guidance and support for Raman analyses; J.A., M.C.A, T.D.H., M.G.M., C.O., D.S., B.W. contributed to data analysis; J.A., S.A.B, M.C.A, E.D.E., Z.E., H.F., S.G., D.P.H., E.M.K., A.L.P., Y.M., S.N., C.O., P.V., G.A.W., L.Z. performed language editing; J.A., M.B.A., A.M.A.G, M.C.A, M.C, C.C., R.C., S.C., T.N.D., J.P.D., Z.E., H.F., M.L.F., H.P.G, D.P.H., K.K., K.S.C., E.M.K., F.L., M.G.M., M.C.M., C.O., S.P., S.N., S.S.S.S., P.V., G.A.W., L.Z., B.L. provided constructive reviews to the paper.

Competing interest declaration

The authors declare no competing interests.

Additional information

Supplementary Information is available for this paper.

Correspondence and requests for materials should be addressed to Veronica Nava (veronica.nava@unimib.it) and Barbara Leoni (barbara.leoni@unimib.it).

Extended data figure/table legends

Extended Table 1. Blank levels for laboratory-based QA/QC, reporting the absolute number of fibres detected in the blank filters used as control for each replicated sample (i.e., trawl).

Extended Data Fig. 1. Comparison of the density distribution of the features of our study lakes and reservoirs (in yellow) to the boxplot of freshwater systems included in the ‘HydroLAKES’ global dataset. The features being compared are: a) lake area in km²; b) mean depth in m; c) lake volume in km³; d) residence time of lakes in years.

Extended Data Fig. 2. Means and standard errors of plastic concentration (particles m⁻³) resulting from the three trawls collected in each lake. The lakes are ranked in descending order based on their particle concentration, from highest to lowest.

Extended Data Fig. 3. Clusters of lakes based on the percentage occurrence of the different plastic shapes, colours, and polymeric compositions.

Extended Data Fig. 4. Scaling 1 of redundancy analysis between plastic concentration in lakes, features of plastics, and environmental and anthropogenic drivers. The dots are coloured based on the concentration of plastics (particles m⁻³) detected.

Extended Data Fig. 5. Density plots and histogram of the Feret’s diameter (width, mm) of the 9,425 particles identified in the 38 lakes analysed. The median trend is indicated by the vertical blue and dashed line.

Extended Data Fig. 6. Images of different shapes of plastic fragments collected in water samples: a, b, c) fragment; d, e, f) fibre; g, h, i) filament; j, k) film; l) sphere/pellet.