

# Sphaerodoridae (Annelida: Polychaeta) from the Bellingshausen Sea (Antarctica) with the description of two new species

J. Moreira<sup>1</sup>; J. Parapar<sup>2</sup>

<sup>1</sup> Estación de Biología Mariña da Graña, Universidade de Santiago de Compostela, Rúa da Ribeira 1, 15590 A Graña, Ferrol, Spain e-mail: [juan.moreira@usc.es](mailto:juan.moreira@usc.es)

<sup>2</sup> Departamento de Biología Animal, Facultade de Ciencias, Universidade da Coruña, 15008 A Coruña, Spain

**Polar Biology**, Volume 34, Issue 2, pp 193–204

Received: 21 April 2010 / Revised: 30 June 2010 / Accepted: 14 July 2010 / Published online: 29 July 2010

## Cite this article as:

Moreira, J. & Parapar, J. *Polar Biol* (2011) 34: 193. doi:10.1007/s00300-010-0869-x

The final publication is available at Springer via <http://dx.doi.org/10.1007/s00300-010-0869-x>

## Abstract

The examination of polychaete collections obtained during the Spanish Bentart 2006 expedition to the Bellingshausen Sea (Antarctica) revealed the presence of several sphaerodorid species. In this work, species belonging to the genera *Sphaerodorum* Örsted, 1843, *Ephesiella* Chamberlin, 1919, *Clavodorum* Hartman and Fauchald, 1971 and *Sphaerephesia* Fauchald, 1972 are reported including two new species belonging to *Sphaerodorum* and *Sphaerephesia*, respectively. A specimen identified as *Ephesiella* sp. might also represent a new species but, due to its poor state of preservation, a formal description is not possible yet. Furthermore, *Sphaerodoropsis polypapillata* Hartmann-Schröder and Rosenfeldt, 1988 is transferred to the genus *Clavodorum* Hartman and Fauchald, 1971 after examination of the type series and specimens obtained from the Bellingshausen Sea.

## Keywords

Polychaeta; Sphaerodoridae; Bellingshausen Sea; Antarctica; Benthos; New species

## Introduction

Sphaerodorids (Polychaeta, Sphaerodoridae) are small polychaetes characterized by having a variable number of tubercles and papillae on the dorsum, which can be sessile or stalked, or arranged in a definite, non-random pattern (Fauchald 1974). In general, sphaerodorids are presumed to be overlooked in many sampling programmes due to their small size and inappropriate processing of samples (Borowski 1994). Although taxonomic and ecological knowledge of this family is still scarce in many geographic areas, sphaerodorids of Antarctic seas are comparatively better known (Fauchald 1974; Hartman 1978; Hartmann-Schröder and Rosenfeldt 1992; Schüller and Ebbe 2007). To date, 13 species have been described as having their type locality in Subantarctic and/or Antarctic

waters; these belong to the genera *Sphaerodorum* Örsted, 1843 (one species), *Ephesiella* Chamberlin, 1919 (3), *Clavodorum* Hartman and Fauchald, 1971 (2) and *Sphaerodoropsis* Hartman and Fauchald, 1971 (7) (see Ehlers 1913; Benham 1921; Hartman 1967; Averincev 1972; Fauchald 1974; Hartmann-Schröder and Rosenfeldt 1988, 1990, 1992). In addition, Hartmann-Schröder and Rosenfeldt (1990) reported the presence of the genus *Ephesiopsis* Hartman and Fauchald, 1971 in Elephant Island after one fragmented specimen. Other species, which were supposed to have a cosmopolitan distribution, such as *Sphaerodorumgracilis* (Rathke, 1843) and *Sphaerodoropsis minuta* Webster and Benedict, 1887, have also been recorded in Antarctic latitudes; in many cases, these records refer actually to other sphaerodorid taxa which turned out to be new species (Fauchald 1974).

The Bellingshausen Sea (Antarctica) constitutes a natural connection between the Ross Sea and the Antarctic Peninsula and is of great zoogeographical importance because of its role in the dispersion of species around the waters of the Antarctic continent and with South America through the Scotia Arc (Sáiz et al. 2008). Nevertheless, the Bellingshausen Sea has been less studied than the Weddell and Ross Seas and the Antarctic Peninsula, where many research programmes have been carried out in the last years (e.g., Arntz et al. 1994; Hartmann-Schröder and Rosenfeldt 1988; Brandt et al. 2004; Hilbig et al. 2006). This is due, in part, to its remoteness and the prevalence of ice during most of the year there. In consequence, little information is available on the polychaete fauna of the Bellingshausen Sea in general, including that referred to sphaerodorids. In this context, the Spanish expeditions *Bentart 2003* and *Bentart 2006* were carried out in order to study the composition and distribution of the benthic fauna in the shelf and slope of the Bellingshausen Sea and adjacent areas.

The examination of the polychaete collections obtained during the *Bentart 2006* expedition revealed the presence of several sphaerodorid species. All specimens belonging to the genera *Sphaerodorum*, *Ephesiella*, *Clavodorum* and *Sphaerephesia* Fauchald, 1972 are reported here, including two new species belonging to *Sphaerodorum* and *Sphaerephesia*; *Sphaerodoropsis* material will be described elsewhere. In addition, *Sphaerodoropsis polypapillata* Hartmann-Schröder and Rosenfeldt, 1988 is transferred to the genus *Clavodorum* after examination of the type series and specimens obtained from the Bellingshausen Sea.

## Materials and methods

The material examined comes from samples obtained during the Spanish *Bentart 2006* expedition at the Bellingshausen Sea (from the Antarctic Peninsula to Thurston Island) and off the western Antarctic Peninsula (from Gerlache Strait to Marguerite Bay). Most of the sphaerodorid specimens were collected with a modified Macer-GIRO-Q Epibenthic Sledge (Cartes et al. 1994); an additional specimen was obtained by means of an Agassiz trawl (Station BS 29). Sampling methodology, location and physical features of the stations where sphaerodorids were found are described in detail in San Vicente et al. (2009).

Observations, drawings and measurements of specimens were made using an Olympus BX40 compound microscope connected to a drawing tube. All specimens were deposited at the Museo Nacional de Ciencias Naturales (MNCN), Madrid, Spain. Specimens used for scanning electron microscopy (SEM) were dehydrated in a graded ethanol series, prepared by critical-point drying using CO<sub>2</sub>, coated with gold in a BAL-TEC SCD 004 evaporator and examined and photographed under a JEOL JSM-6400 scanning electron microscope at the Servicios de Apoyo á Investigación, Universidade da Coruña (SAIN), Spain. The nomenclature of prostomial appendages follows that of Aguirrezabalaga and Ceberio (2005). The macrotubercle located dorsally to the parapodium has been

considered by several authors as the dorsal notopodial cirrus (e.g., Ruderman 1911; Pleijel 2001). In this paper, dorsal cirri are not specifically described because, if present, they would not differ from the other dorsal macrotubercles. The code of stations refers to the sampling area: BS, Bellingshausen Sea; WAP, western Antarctic Peninsula. The abbreviations for the structures referred in the figures are the following: ap, dorsal antenniform papilla; cl, chaetal lobe; dp, distal papilla; ia, intermediate lateral antenna; ldp, latero-dorsal papilla; lp, lateral papilla; ma, median antenna; mt, microtubercle; MT, macrotubercle; pa, palps; pc, peristomial cirrus; vc, ventral cirrus; vp, ventral papilla.

## Results and discussion

### Taxonomic account

Family *Sphaerodoridae* Malmgren, 1867

*Sphaerodorum olgae* sp. nov. (Figs. 1, 2, 3)

### Material examined

**St. BS 38**, one complete specimen, 10.7 mm long, 0.3 mm wide, with 77 chaetigers (Holotype, MNCN 16.01/13158). **St. BS 30**, one spec. (Paratype, MNCN 16.01/13159). **St. BS 31**, one spec. (Paratype, MNCN 16.01/13160). **St. BS 33**, 3 spec. (Paratypes, MNCN 16.01/13161). **St. BS 34**, one spec. (Paratype, MNCN 16.01/13162).


### Description

Specimens measuring 6.1–11.2 mm long, 0.17–0.37 mm wide, with 60–90 chaetigers. Body long, tapering distally; yellowish-whitish in ethanol. Tegument wrinkled, with a granulated appearance (Fig. 2b). Prostomium bluntly rounded, fused to peristomium (Figs. 1a, b, 2a). Median antenna and three pairs of lateral prostomial appendages (palps, intermediate antennae, and dorsal antenniform papillae). Median antenna clavate. Intermediate antennae and palps digitiform, longer than median antenna. Dorsal antenniform papillae digitiform, shorter than intermediate antennae and palps; contracted and difficult to detect in some specimens. One pair of peristomial cirri, digitiform, slightly shorter than palps and intermediate antennae. Prostomium and peristomium with papillae; about 8–10 papillae among prostomial paired appendages. Eyes and pharynx not seen. Two dorso-lateral macrotubercles per chaetiger (Fig. 1e); spherical to pear-shaped, provided with terminal papilla. From chaetiger 2 posteriorly two microtubercles per chaetiger, dorsal to macrotubercles, with a basal collar and a long terminal papilla (Fig. 1d). Macrotubercles and microtubercles arranged in four longitudinal rows. Papillae on all body surfaces. About four dorsal transversal rows of papillae per chaetiger, numbering approximately 20 papillae. About 5–6 ventral transversal rows of papillae per chaetiger, numbering more than 30 papillae. Some papillae on lateral surfaces among parapodia. Total number and arrangement of body papillae difficult to ascertain due to contraction of specimens and wrinkled appearance of tegument.

Parapodia uniramous, longer than wide (Figs. 1f, g, 2c–e). Parapodia of first chaetiger in ventral position; following parapodia lateral. Digitiform chaetal lobe, arising dorsally among chaetae, projecting beyond acicular lobe. Ventral cirri cylindrical, usually with a distal protuberance, longer and thicker than chaetal lobe, surpassing acicular lobe tip. Parapodia with about 9–11 digitiform papillae, shorter than chaetal lobe: one distal papilla, one on latero-dorsal parapodial face, 2–3 on each lateral face and 2–3 on ventral surface medial to ventral cirrus; papillae on ventral surfaces more spherical than those on lateral and dorsal surfaces. Parapodia of last chaetigers reduced. Recurved

hooks in first chaetiger (one per parapodium), visible in ventral view (Fig. 1b, 2f). Simple chaetae usually numbering 4 per fascicle, with small subdistal boss (Fig. 3a, b); distal end falciform, pointed. Spinulation along internal border, from distal end to subdistal boss; some spinulation along external border. Most chaetae with a suture-like depression somewhat separating distal end and subdistal boss (Fig. 3c–d); when observed in the compound microscope appearing as if blade is fused to shaft.


Pygidium with two dorsal macrotubercles and midventral digitiform anal cirrus (Fig. 1c).



**Fig. 1** *Sphaerodorum olgae* sp. nov. **a** anterior end, dorsal view, **b** anterior end, ventral view, **c** pygidium, ventral view, **d** dorsal microtubercle, **e** dorsal macrotubercle, **f** parapodium, chaetiger 7, dorsal view, **g** parapodium, chaetiger 16, ventral view. **a–c, f–g**, same scale

## Remarks

The new species differs mostly from the other six species of the genus in the features of the parapodium including number and arrangement of papillae. Thus, *S. olgae* sp. nov. differs from *Sphaerodorum indutum* Fauchald, 1974 and *S. papillifer* Moore, 1909 in having less parapodial papillae; those of *S. indutum* are larger and number at least 15 while those of *S. papillifer* are numerous on each parapodial face, small and conical in shape. In *S. recurvatum* Fauchald, 1974, the parapodia lack papillae and body papillae are absent from the dorsum, whereas in *S. olgae* sp. nov. body papillae are present both on the dorsum and venter. *Sphaerodorum gracilis* (Rathke, 1843) also has a parapodium with a similar number of papillae to that of *S. olgae* sp. nov. but these species differ in arrangement and size of their parapodial papillae; thus, in *S. gracilis* there are 5–6 papillae




**Fig. 2** *Sphaerodorum olgae* sp. nov. SEM micrographs. **a** anterior end, *frontal view*, **b** tegument, **c–e** mid-body parapodia, **f** first chaetiger, parapodia and hooks

regularly distributed along each lateral parapodial surface, which are proportionally smaller than those of *S. olgae* sp. nov. when compared to the length of the parapodium. In addition, the prechaetal lobe in *S. gracilis* is usually conical rather than digitiform and the body papillae on dorsal surfaces are slightly more numerous than in *S. olgae* sp. nov. On the other hand, *Sphaerodorum olgae* sp. nov. differs from *S. vietnamense* Fauchald, 1974 in the shape of the ventral cirri, which are long and slender in the latter; in *S. olgae* sp. nov., ventral cirri are thick and cylindrical in shape. In addition, the simple chaetae of *S. vietnamense* are of two types, including some strongly recurved ones which are not present in *S. olgae* sp. nov. In fact, the simple chaetae of *S. olgae* sp. nov. are strikingly similar to those of *S. ophiurophoretos* Martín and Alva, 1988; the latter also presents chaetae showing a protuberance which seems to mark the limit between the blade and the shaft (cfr. Fig. 2D in Martín and Alva 1988). Some simple chaetae of *S. olgae* sp. nov. (cfr. Fig. 2) are similar to the aforementioned chaetae, showing in this case a suture-like depression that makes the simple chaetae looking like a composite one in which the blade is fused to the shaft. Nevertheless, *S. ophiurophoretos* differs from *S. olgae* sp. nov. in having a shorter body, fewer chaetigers (8–9 vs. 60–90) and fewer parapodial papillae; the latter are spherical and similar to those found on the rest of the body instead of being digitiform as in *S. olgae* sp. nov.

### Etymology

The new species is dedicated to the memory of Dr. Olga Hartman, for her many contributions to our knowledge of the Antarctic polychaete fauna.



**Fig. 3** *Sphaerodorum olgae* sp. nov. SEM micrographs. **a–b** simple chaetae, **c–d** simple chaetae with suture-like depression between blade and shaft indicated by *white arrows*

### *Ecology and distribution*

To date, only known from the Bellingshausen Sea in muddy gravel, muddy sand, sandy mud and mud, at depths of 431–1,799 m.

### ***Ephesiella muehlenhardtae* Hartmann-Schröder and Rosenfeldt, 1988**

*Ephesiella muehlenhardtae* Hartmann-Schröder and Rosenfeldt, 1988: 47–48, figs 32–38.

### *Material examined*

**St. BS 29**, one complete specimen, 9.5 mm long, 0.37 mm wide, with 63 chaetigers (MNCN 16.01/13164).

### *Ecology and distribution*

This species is known from Elephant Island and Bransfield Strait (Hartmann-Schröder and Rosenfeldt 1990), in bottoms ranging from gravel and stones to fine sand and silt at depths of 93–342 m (Hartmann-Schröder and Rosenfeldt 1988, 1990, 1992). The only specimen found at the Bellingshausen Sea (Peter I Island; this work) was collected at a muddy bottom at a depth of 3,219–3,280 m, thus extending the known bathymetric range of this species.

### ***Ephesiella* sp.**

### *Material examined*

**St. BS 34**, one complete specimen, 6.75 mm long, 0.75 mm wide, with 48 chaetigers (MNCN 16.01/13165).

### *Remarks*

The only specimen available is in poor condition; the prostomium is contracted but one median antenna and two lateral prostomial appendages, all of similar size, are present. The parapodium has a ventral cirrus not projecting beyond the acicular lobe and also bears an oval dorsal papilla in distal position which resembles a prechaetal lobe; each parapodium also bears at least 10–12 papillae scattered along all parapodial surfaces and around the chaetal lobe. The parapodia of the first chaetiger are provided each with one protruding hook; all remaining chaetae are compound falcigers. The other known Antarctic species of the genus *Ephesiella*, namely *E. antarctica* (McIntosh, 1885), *E. pallida* Fauchald, 1974 and *E. muehlenhardtae* Hartmann-Schröder and Rosenfeldt, 1988, differ from our specimen in the number and arrangement of parapodial papillae. Thus, *E. antarctica* has a nearly smooth parapodium which bears a postchaetal lobe, *E. pallida* has one prechaetal lobe, one postchaetal lobe and one pair of papillae on each face of the parapodium, and *E. muehlenhardtae* has about 5–6 parapodial papillae and a chaetal lobe (cfr. Hartmann-Schröder and Rosenfeldt 1988, figs 35–37). Although our specimen might represent a new species to science, a formal description is not possible due to its state of preservation.

### *Ecology and distribution*

The only specimen available was collected in the Bellingshausen Sea in muddy fine sand at depths of 612–620 m.

*Clavodorum polypapillata* (Hartmann-Schröder and Rosenfeldt, 1988) **comb. nov.** (Figs. 4, 5)

*Material examined*

**St. BS 37**, one specimen (MNCN 16.01/13166). **St. WAP 41**, 3 spec. (MNCN 16.01/13167). **St. WAP 43**, 5 spec. (MNCN 16.01/13168).

*Additional material (type series)*

*Sphaerodoropsis polypapillata* Hartmann-Schröder and Rosenfeldt, 1988, Zoological Museum, Hamburg. Holotype (HZM: P-19161), paratype (HZM: P-19162).

*Description*

*Bentart 2006* specimens measuring 2.0–3.3 mm in length and 0.6–1.3 mm in width, with 22–25 chaetigers. Body short, grub-like, inflated, lacking pigmentation, transparent-whitish in ethanol (Fig. 5a). Tegument with a granulated appearance (Fig. 5b).

Prostomium bluntly rounded, fused to peristomium; in some specimens a constriction between the peristomium and the rest of the body. Median antenna and two pairs of lateral prostomial appendages (Fig. 4a). Median antenna long, digitiform (Fig. 4b). Intermediate antennae as long as median antenna, with 3 basal digitiform papillae. Palps digitiform, shorter than intermediate antennae, with 3 basal digitiform papillae. Peristomial cirri digitiform, shorter than antennae and palps, with a basal digitiform papilla. Peristomium with dorsal transversal row of about 10 digitiform papillae. Two brown eyes between palps and prostomial paired appendages. Several ventral digitiform papillae between peristomial cirri and mouth (Fig. 4c). Pharynx extending over 3–4 segments. Dorsal macrotubercles spherical, stalked (Fig. 4d). Macrotubercles numbering 12–17 per chaetiger arranged in a transverse row along dorsal and lateral surfaces (Fig. 4e). Spherical microtubercles irregularly distributed among rows of macrotubercles. Ventral surfaces densely covered with microtubercles provided with short stalk (Fig. 4f); microtubercles similar to those on dorsal and lateral body surfaces, the largest slightly smaller than dorsal macrotubercles.


Parapodia uniramous, with wrinkled surface, longer than wide (Figs. 4g, h, 5c). Digitiform prechaetal lobe from chaetiger 7 backwards, projecting beyond acicular lobe; postchaetal lobe absent. Ventral cirri digitiform, slightly shorter than or as long as prechaetal lobe, surpassing acicular lobe tip. Parapodia with 3–4 stalked papillae: one on anterior lateral parapodial face, one on posterior lateral face and 1–2 on ventral surface behind ventral cirrus. Composite chaetae numbering about 10–15 per fascicle; blades unidentate with recurved tip and thin spinules along cutting margin (Fig. 5d). Blades showing gradation in length in the same parapodium; blades about 50–30 µm long in anterior chaetigers and 35–25 µm long in posterior ones.

Pygidium with about 8 small dorsal papillae, two larger spherical lateral papillae and midventral digitiform anal cirrus, the latter slightly longer than parapodial ventral cirri.



Remarks

Hartmann-Schröder and Rosenfeldt (1988) described *Sphaerodoropsis polypapillata* from King George Island (Antarctica) and then reported the same species from Elephant Island (Hartmann-Schröder and Rosenfeldt 1992). The re-examination of the type series and the additional specimens collected during the *Bentart 2006* expedition confirmed the presence of stalked macrotubercles instead of the sessile ones reported in the original description. Macro- and microtubercles are close to each other in the examined specimens which makes difficult to distinguish any stalk. Nevertheless, some specimens including those of the type series had already lost some macrotubercles but not their stalks thus revealing their true morphology. Therefore, *S. polypapillata* is here transferred to the genus *Clavodorum* because of the possession of stalked tubercles and a median antenna which is at




**Fig. 4** *Clavodorum polypapillata* (Hartmann-Schröder and Rosenfeldt, 1988) comb. nov. **a** anterior end, *lateral view* (arrow indicates position of macrotubercle stalk), **b** anterior end, *lateral view*, detail, **c** anterior end, *ventral view*, **d** stalked macrotubercle, **e** mid-body segments, *dorsal view*, **f** mid-body segments, *ventral view*, **g** parapodium, chaetiger 8, *dorsal view*, **h** parapodium, chaetiger 8, *ventral view* (chaetae not illustrated). **e–f**, **g–h**, same scale

*Sphaerodoropsis polypapillata* Hartmann-Schröder and Rosenfeldt, 1988: 49–50, fig. 45; 1992: 107–108.

least as long as the intermediate ones. The genus *Sphaerodoridium* Lützen, 1961 also bears stalked macrotubercles but the median antenna is shorter than other prostomial paired appendages and ventral microtubercles are, in general, not stalked (Bakken 2002). *Clavodorum polypapillata* comb. nov. differs mainly from the other known species of the genus in having more than ten longitudinal rows of macrotubercles. Thus, *C. bengalorum* Fauchald, 1974 bears eight rows of macrotubercles, whereas *C. fusum* (Hartman, 1967), *C. atlanticum* Hartman and Fauchald, 1971, *C. clavatum* Fauchald, 1972, *C. adriaticum* Katzmann, 1974, *C. longipes* Fauchald, 1974, *C. fauchaldi* Desbruyères, 1980, *C. mexicanum* Kudenov, 1987 and *C. antarcticum* Hartmann-Schröder and Rosenfeldt, 1990 bear up to six rows of macrotubercles. In addition, *Clavodorum andamanense* Bakken, 2002 differs from *C. polypapillata* comb. nov. in having dorsal macrotubercles arranged in ten longitudinal rows, the body ventral papillae are less numerous, numbering about six per chaetiger, and the parapodium bears one small postchaetal lobe and is provided with only two papillae instead of 3–4.

#### *Ecology and distribution*

The specimens from the *Bentart 2006* expedition were collected at the Bellingshausen Sea and western Antarctic Peninsula in muddy gravel at depths of 246–516 m. This species has previously been reported from soft bottoms of the Weddell Sea ranging from fine sand with stones to silt-clay,



**Fig. 5** *Clavodorum polypapillata* (Hartmann-Schröder and Rosenfeldt, 1988) comb. nov. SEM micrographs. **a** habitus, ventral view, **b** tegument, **c** mid-body parapodia, ventral view, **d** composite chaetae

at depths of 96–532 m (Hartmann-Schröder and Rosenfeldt 1988, 1990, 1992; Schüller and Ebbe 2007).

*Sphaerephesia gesae* sp. nov. (Figs. 6, 7)

*Material examined*

**St. BS 34**, one complete specimen, 2.1 mm long, 0.5 mm wide, with 11 chaetigers (Holotype MNCN 16.01/13163).

*Description*

Body short, grub-like, lacking pigmentation, tegument transparent-whitish in ethanol. Prostomium bluntly rounded, fused to peristomium (Fig. 6a). Median antenna short, probably contracted; two pairs of lateral prostomial appendages (palps and intermediate antennae), digitiform, longer than the median antenna (Fig. 6b). Six small papillae encircled by lateral paired appendages; six similar papillae around mouth opening. Peristomial cirri digitiform, shorter than lateral paired prostomial appendages. Eyes and pharynx not seen. Dorsal macrotubercles sessile, mostly spherical or rounded, arranged in 10 longitudinal rows forming a zig-zag pattern. Chaetigers 1–2 with ten dorsal macrotubercles each; 12 macrotubercles per segment from chaetiger 3 posteriorly. Macrotubercles arranged in two transversal rows: six macrotubercles on each chaetiger parapodial area and six on the interparapodial area. On each parapodial area the four lateralmost macrotubercles are provided with a digitiform papilla on the top, the other two macrotubercles smooth; the four dorsalmost macrotubercles spherical in shape and the two ventralmost more digitiform (Fig. 6c, d). Macrotubercles on interparapodial areas not provided with papillae. Chaetigers 1–2 with six macrotubercles on parapodial areas and four macrotubercles on interparapodial areas; in chaetiger 2 the two dorsalmost on interparapodial area smaller than the others. Dorsal papillae present between rows of macrotubercles, numbering up to 11 per segment in mid-body, arranged in 11 longitudinal rows following a non-random pattern (Fig. 7a).

Venter with small papillae. Chaetigers 1–4 with up to 10 ventral papillae each, six on each parapodial area (three on each side arranged in a V-shape) and four on each interparapodial area arranged in a line perpendicular to the anterior-posterior body axis (Fig. 7b). Two additional ventral papillae from chaetiger 5 backwards, one on each side, close to the posteriormost papilla near the parapodial base.

Parapodia uniramous, longer than wide, with one acicula; digitiform prechaetal lobe from chaetiger 5 backwards, projecting beyond acicular lobe (Fig. 6e); postchaetal lobes absent. Ventral cirri digitiform, larger than prechaetal lobe, reaching acicular lobe tip (Fig. 6f). One small papilla on anterior parapodial surface from chaetiger 1 backwards (Fig. 6g). Composite falcigers numbering 5–7 per fascicle; distal end of shaft inflated, with 3–4 indentations (Fig. 6h). Blades unidentate with long, recurved tip; no spinulation observed along cutting margin. Blades up to 17 µm long, the ventralmost ones slightly shorter.


Pygidium terminal, with two small dorsal papillae, a pair of lateral anal cirri, similar in size and shape to the lateralmost macrotubercles with terminal papilla, and midventral digitiform anal cirrus (Fig. 6i).

## Etymology

This species is named after Dr. Gesa Hartmann-Schröder because of her many contributions to the knowledge of Antarctic polychaetes in general, and of sphaerodorids in particular.


## Remarks

The only short-body sphaerodorid genus with more than two rows of macrotubercles bearing a terminal papilla is *Sphaerephesia* Fauchald, 1972 (Fauchald 1974; Kudenov 1987). The five known species of the genus, namely *S. longisetis* Fauchald, 1972, *S. similisetis* Fauchald, 1972, *S. chilensis* Fauchald, 1974, *S. fauchaldi* Kudenov, 1987 and *S. regularis* Böggemann, 2009 present four longitudinal rows of macrotubercles, all provided with button-shaped or stout terminal papilla, and bear composite chaetae (Fauchald 1972; Kudenov 1987; Böggemann 2009); dorsal surfaces may present a number of microtubercles that are provided with a distal papilla (*S. longisetis*) or may be



**Fig. 6** *Sphaerephesiagesae* sp. nov. **a** anterior end, dorsal view, **b** anterior end, ventral view, **c** digitiform lateral macrotubercle with terminal papilla, **d** spherical latero-dorsal macrotubercle with terminal papilla, **e** parapodium, chaetiger 5, dorsal view, **f** parapodium, chaetiger 8, ventral view, **g** parapodium, chaetiger 4, anterior view, **h** composite chaeta, **i** distal end, ventral view

more or less capitated (*S. fauchaldi*). Because of the presence of more than two longitudinal rows of macrotubercles with terminal papilla and composite chaetae, the new species described here is tentatively included in that genus. Nevertheless, *Sphaerephesia gesae* sp. nov. differs from all other *Sphaerephesia* species in bearing more than four longitudinal rows of macrotubercles and in having only some macrotubercles of each parapodial area provided with a terminal papilla, which is longer than those reported from other species of *Sphaerephesia*. Indeed, the regular presence of two macrotubercles with a terminal papilla on each side of each row on the parapodial area suggests that this condition is not an artifact related to the fixation of the specimens. These features make *S. gesae* sp. nov. unique among all described sphaerodorids. On the other hand, the presence of more than four rows of macrotubercles arranged in a zig-zag pattern is a feature found in several species of the genus *Sphaerodoropsis*. In addition, the appearance and features of the parapodia of *S. gesae* sp. nov. are similar to those described from a number of *Sphaerodoropsis* species, namely *S. bisphaeroserialis* (Hartmann-Schröder, 1974), *S. arctowskyensis* Hartmann-Schröder and Rosenfeldt, 1988, *S. translucida* Borowski, 1994 and *S. garciaalvarezi* Moreira et al. 2004. The only species of *Sphaerodoropsis* having macrotubercles arranged in more or less a zig-zag pattern and bearing a terminal papilla is *S. translucida*, according to the redescription provided by Böggemann (2009). This species was originally described as having the macrotubercles with “distal end more or less set off, but without terminal papilla” (Borowski 1994), and not arranged following a definite pattern. Böggemann (2009) states that the possession of macrotubercles provided with a terminal papilla is in contrast to the accepted diagnosis of the genus *Sphaerodoropsis*. Because of that, *S. gesae* sp. nov. was not included in *Sphaerodoropsis*. In fact, *Sphaerodoropsis* is now composed by an assemblage of species which might represent several different genera. For example, species can be grouped according to the number and arrangement of macrotubercles (Borowski 1994). In addition,



**Fig. 7** *Sphaerephesiagesae* sp. nov. Schematic arrangement of: **a** dorsal macrotubercles and papillae, chaetiger 6, **b** ventral papillae, chaetigers 4–6. Symbols: large circle, smooth macrotubercle; large circle with inner small circle, macrotubercle with terminal papilla; small circle, dorsal/ventral papilla

some species with two transversal rows of macrotubercles arranged in a zig-zag pattern seem to differ from the others in having inflated ventral parapodial cirri in the 6th chaetiger, which could be interpreted as a penis structure (Moreira et al. 2004; Böggemann 2009). At this point, a revision of the whole family and of this genus, in particular, is needed (Borowski 1994; Aguado and Rouse 2006).

As only one specimen of *S. gesae* sp. nov. is available, no new genus has been erected for this species in spite of its unique features among sphaerodorids, mostly regarding the number and shape of dorsal macrotubercles. Indeed, examination of further material is necessary to assess the true position of this and other species within the Sphaerodoridae.

#### *Ecology and distribution*

This species is only known from the Bellingshausen Sea, in muddy fine sand at depths of 612–620 m.

### **General remarks**

Three genera of sphaerodorid polychaetes, represented by five species, are reported here for the first time for the Bellingshausen Sea; two of them are new to science. This raises the total number of known Antarctic sphaerodorids to 15. Two of these species have previously been reported from other Antarctic areas, namely *Ephesiella muehlenhardtae* and *Clavodorum polypapillata*; the latter has also been suggested to be present in the southern Atlantic Ocean (Schüller and Ebbe 2007). In fact, most of the sphaerodorid species previously described from Antarctic and Subantarctic waters have not been reported yet from other latitudes, the exception being *Sphaerodoropsis parva* (Ehlers, 1913), which has been cited in Australia and southern South America (Hartmann-Schröder and Rosenfeldt 1988) and in the eastern North Atlantic (Desbruyères 1980). However, we suspect that non-Antarctic records of this species might refer to other similar taxa, whether they represent new taxa or not. Thus, the specimens named *S. parva* by Wesenberg-Lund (1962) from the coast of Chile turned out to be a new species after the world revision of this family done by Fauchald (1974) and named *Sphaerephesia chilensis* Fauchald, 1974. Similarly, other polychaete taxa thought as having a wide distribution and previously reported from the Southern Ocean might actually not be represented there. This is the case of the northern trichobranchid *Terebellides stroemii* Sars, 1835 which was supposed to be represented at southern latitudes by the subspecies *T. stroemii kerguelensis* McIntosh, 1885, which recently Parapar and Moreira (2008), after revision of the type material, proposed elevated to the species rank leaving the distribution of *T. stroemii* limited to northern latitudes.

The sphaerodorid fauna from the Southern Ocean seems therefore to be highly endemic (Fauchald 1974; Schüller and Ebbe 2007). Schüller and Ebbe (2007) suggest that this fact might be the result of radiation events related to the recolonization of Southern Ocean sediments. Nevertheless, the true degree of endemism in this area cannot be fully assessed yet due to the many gaps in our knowledge of the taxonomy, distribution and ecology of sphaerodorids that still exist. On the one hand, these polychaetes are easily overlooked in sediment samples due to their small body size, and they thus need to be sampled adequately, for example, by means of gear like the Epibenthic Sledge which has previously demonstrated its usefulness in obtaining large numbers of specimens of this family of polychaetes (Schüller and Ebbe 2007; Parapar and Moreira 2009). On the other hand, the state of preservation of the specimens makes identification to the species level difficult; many individuals appear contracted, and characters of high taxonomic relevance such as the arrangement of tubercles and papillae are difficult to ascertain. In addition, Clarke (2008) points out that cryptic species may be common in Antarctic benthic faunas as suggested by molecular studies. Indeed, many sphaerodorid species are morphologically very similar and in many cases, due to this, are

distinguished only according to subtle differences. Therefore, we believe that future morphological work should be combined with molecular analyses in order to assess the identity of some taxa and thus determine their true geographic and bathymetric distribution.

Many Antarctic polychaetes are supposed to be distributed only at shelf and slope depths (Brandt et al. 2009). Here, four out of the five sphaerodorid species found were only present on the shelf and slope; it must be taken into account that most of the samples collected in the *Bentart 06* expeditions were taken on a limited range of depths (150–2000 m). Nevertheless, it is noteworthy that the only specimen of *E. muehlenhardtae* was found deeper than 3,000 m, thus greatly extending the previously known depth range of this species, which was reported between 93 and 342 m.

According to the few published papers on a number of benthic taxa, including molluscs (Troncoso et al. 2007; Troncoso and Aldea 2008) and polychaetes (Parapar et al. in press) as well as when considering higher taxonomic categories (Sáiz et al. 2008), the Bellingshausen Sea seems to be a well-defined biogeographical area within the Southern Ocean. This fact has tentatively been related to depth and the very nature of the sediment, which is mostly composed by deposits of foraminiferans and diatoms (Hillenbrand et al. 2003). The presence of the two new sphaerodorid species described here might strengthen this conclusion. However, as explained above, more data are needed to fully assess the biogeographical characteristics of the Bellingshausen Sea, at least in regard to its polychaete fauna.

## Acknowledgments

The authors wish to thank Dr. A. Brandt (Zoological Museum, Hamburg) for the loan of type series of *Sphaerodoropsis polypapillata*. Many thanks also to A. Castro and C. Sueiro (SAIN, Universidade da Coruña) and R. Barreiro (RIAIDT, Universidade de Santiago de Compostela, Spain) who assisted with the preparation of specimens and use of the SEM, to Y. Lucas for assisting with the preparation of the drawings and to J. García-Carracedo for the translation of several bibliographic references and the revision of the English version of the manuscript. The authors are also grateful to T. Bakken, B. Ebbe and D. Eibye-Jacobsen for their constructive comments on an earlier version of the manuscript. Special thanks to the crew of R/V *Hespérides* and all the colleagues which collaborated in the *Bentart 2006* expedition and especially to Carles San Vicente (Generalitat de Catalunya) and Patrick Arnaud (Station Marine de Endoume, France) who collected the specimens from the Epibenthic Sledge samples. The *Bentart* expeditions were carried out under the auspices of the Spanish Government through the Ministry of Education and Science and were supported by the Spanish Antarctic Programme (CICYT) (REN 2001-1074/ANT and CGL2004-01856).

## References

- Aguado MT, Rouse GW (2006) First record of Sphaerodoridae (Phyllodocida: Annelida) from hydrothermal vents. *Zootaxa* 1383:1–21
- Aguirrezabalaga F, Ceberio A (2005) *Sphaerodoropsis amoureuksi* and *S. stellifer*, two new species of Sphaerodoridae (Polychaeta) from the Capbreton Canyon (Bay of Biscay, NE Atlantic). *Cah Biol Mar* 46:9–20
- Arntz WE, Brey T, Gallardo VA (1994) Antarctic Zoobenthos. *Oceanogr Mar Biol Annu Rev* 32:241–304
- Averincev VG (1972) Benthic polychaetes Errantia from the Antarctic and Subantarctic collected by the Soviet Antarctic expeditions. Exploration of the fauna of the Seas. *Biol Res Sov Antarct Exped* 5:88–293 (in Russian)

- Bakken T (2002) Sphaerodoridae (Annelida: Polychaeta) from the BIOSHELF Project, Andaman Sea, Thailand. *Phuket Mar Biol Center Res Spec Publ* 24:197–204
- Benham WB (1921) Polychaeta. Australasian Antarctic Expedition 1911–14 under the leadership of Sir Douglas Mawson, D.Sc. B.E. *Sci Rep Ser C* 6:1–128
- Böggemann M (2009) Polychaetes (Annelida) of the abyssal SE Atlantic. *Org Div Evol* 9:251–428
- Borowski C (1994) Three new deep-sea species of Sphaerodoridae (Annelida, Polychaeta) from the eastern tropical South Pacific. *Zool Scr* 23:193–203
- Brandt A, De Broyer C, Gooday AJ, Hilbig B, Thomson MRA (2004) Introduction to ANDEEP (ANtartic benthic DEEP-sea biodiversity: colonization history and recent community patterns)-a tribute to Howard L. Sanders. *Deep Sea Res II* 51(14–16):1457–1465
- Brandt A, Linse K, Schüller M (2009) Bathymetric distribution patterns of Southern Ocean macrofaunal taxa: Bivalvia, Gastropoda, Isopoda and Polychaeta. *Deep Sea Res I* 56:2013–2025
- Cartes JE, Sorbe JC, Sarda F (1994) Spatial distribution of deep-sea decapods and euphausiids near the bottom in the northwestern Mediterranean. *J Exp Mar Biol Ecol* 179:131–144
- Clarke A (2008) Antarctic marine benthic diversity: patterns and processes. *J Exp Mar Biol Ecol* 366:48–55
- Desbruyères D (1980) Sphaerodoridae (Annélides Polychètes) profonds du Nord-Est Atlantique. *Bull Mus Natn Hist Nat, Paris, 4e sér, 2, section A(1)*:109–128
- Ehlers E (1913) Die Polychaeten-Sammlungen der Deutschen Südpolar-Expedition 1901–1903. *Dtsch Südpolar Exped* 13:397–598
- Fauchald K (1972) Benthic polychaetous annelids from deep water off western Mexico and adjacent areas in the eastern Pacific Ocean. *Allan Hancock Monogr Mar Biol* 7:1–575
- Fauchald K (1974) Sphaerodoridae (Polychaeta: Errantia) from world-wide areas. *J Nat Hist* 8:257–289
- Hartman O (1967) Polychaetous annelids collected by the USNS Eltanin and Staten Island cruises, chiefly Antarctic seas. *Allan Hancock Monogr Mar Biol* 2:1–387
- Hartman O (1978) Polychaeta from the Weddell Sea quadrant, Antarctica. Paper 4. Biology of the Antarctic Seas VI. *Antarct Res Ser* 26:125–223
- Hartmann-Schröder G, Rosenfeldt P (1988) Die Polychaeten der “Polarstern”—Reise ANTIII/2 in die Antarktis 1984. Teil 1: Euprosinidae bis Chaetopteridae. *Mitt Hamb Zool Mus Inst* 85:25–72
- Hartmann-Schröder G, Rosenfeldt P (1990) Die Polychaeten der Walter Herwig-Reise 68/1 nach Elephant Island (Antarktis) 1985 Teil 1: Aphroditidae bis Cirratulidae. *Mitt Hamb Zool Mus Inst* 87:89–122
- Hartmann-Schröder G, Rosenfeldt P (1992) Die Polychaeten der “Polarstern”—Reise ANT V/1 in die Antarktis 1986. Teil 1: Euprosinidae bis Iphitimidae. *Mitt Hamb Zool Mus Inst* 89:85–124
- Hilbig B, Gerdes D, Montiel A (2006) Distribution patterns and biodiversity in polychaete communities of the Weddell Sea and Antarctic Peninsula area (Southern Ocean). *J Mar Biol Assoc UK* 86:711–725
- Hillenbrand C-D, Grobe H, Diekmann B, Kuhn G, Fütterer DK (2003) Distribution of clay minerals and proxies for productivity in surface sediments of the Bellingshausen and Amundsen seas (West Antarctica)—relation to modern environmental conditions. *Mar Geol* 193:253–271
- Kudenov JD (1987) Five new species of Sphaerodoridae (Annelida: Polychaeta) from the Gulf of Mexico. *Proc Biol Soc Wash* 100:927–935
- Martin D, Alva V (1988) Un polychète nouveau *Sphaerodorum ophiurophoretos* nov. sp. (Polychaeta: Sphaerodoridae), symbiotique de l’ophiure *Amphipholis squamata* (Delle Chiaje, 1828). *Bull Koninklijk Belg Inst Natur Biol* 58:45–49
- Moreira J, Cacabelos E, Troncoso JS (2004) A new species of *Sphaerodoropsis* (Polychaeta: Sphaerodoridae) from north-east Atlantic, with comments on other species of the genus. *J Mar Biol Assoc UK* 84(5):995–1000



- Parapar J, Moreira J (2009) Polychaeta of the 'DIVA-Artabria I' project (cruise 2002) in the continental shelf and upper slope off Galicia (NW Spain). *Cah Biol Mar* 50:57–78
- Parapar J, Moreira J (2008) Redescription of *Terebellides kerguelensis* stat. nov. McIntosh, 1885 (Polychaeta: Trichobranchidae) from Antarctic and subantarctic waters. *Helg Mar Res* 62:143–152
- Parapar J, López E, Gambi MC, Núñez J, Ramos A (in press) Quantitative analysis of soft-bottom polychaetes of the Bellingshausen Sea and Gerlache Strait (Antarctica). *Pol Biol*
- Pleijel F (2001) Sphaerodoridae Malmgren, 1867. In: Rouse GW, Pleijel F (eds) *Polychaetes*. Oxford University Press, Oxford, pp 136–138
- Ruderman L (1911) Recherches sur *Ephesia gracilis* Rathke, Annélide polychète de la famille des sphaerodorides: morphologie, anatomie, histology. *Mém Soc zool Fr* 24:1–96
- Saiz JI, García FJ, Manjón-Cabeza ME, Parapar J, Peña-Cantero A, Saucède T, Troncoso J, Ramos A (2008) Community structure and spatial distribution of benthic fauna in the Bellingshausen Sea (West Antarctica). *Polar Biol* 31:735–743
- San Vicente C, Munilla T, Corbera J, Sorbe JC, Ramos A (2009) Suprabenthic fauna from the Bellingshausen Sea and western Antarctic Peninsula: spatial distribution and community structure. *Sci Mar* 73:357–368
- Schüller M, Ebbe B (2007) Global distributional patterns of selected deep-sea Polychaeta (Annelida) from the Southern Ocean. *Deep Sea Res II* 54:1737–1751
- Troncoso JS, Aldea C (2008) Macrobenthic mollusc assemblages and diversity in the West Antarctica from the South Shetland Islands to the Bellingshausen Sea. *Polar Biol* 31:1253–1265
- Troncoso JS, Aldea C, Arnaud P, Ramos A, García F (2007) Quantitative analysis of soft-bottom molluscs in the Bellingshausen Sea and around Peter I Island. *Polar Res* 26:126–134
- Wesenberg-Lund E (1962) Polychaeta Errantia. Reports of the Lund University Chile Expedition 1948–1949 (43). *Lunds Universitets Årsskrift NF Avd 2*, 57(12):1–139