

## Comprehensive in-hospital monitoring in acute heart failure: applications for clinical practice and future directions for research. A statement from the Acute Heart Failure Committee of the Heart Failure Association (HFA) of the European Society of Cardiology (ESC)

Veli-Pekka Harjola<sup>1</sup>, John Parissis<sup>2</sup>, Hans-Peter Brunner-La Rocca<sup>3</sup>, Jelena Čelutkienė<sup>4</sup>, Ovidiu Chioncel<sup>5</sup>, Sean P. Collins<sup>6</sup>, Daniel De Backer<sup>7</sup>, Gerasimos S. Filippatos<sup>2</sup>, Etienne Gayat<sup>8</sup>, Loreena Hill<sup>9</sup>, Mitja Lainscak<sup>10,11</sup>, Johan Lassus<sup>12</sup>, Josep Masip<sup>13,14</sup>, Alexandre Mebazaa<sup>15,16,17,18</sup>, Òscar Miró<sup>19</sup>, Andrea Mortara<sup>20</sup>, Christian Mueller<sup>21</sup>, Wilfried Mullens<sup>22</sup>, Markku S. Nieminen<sup>23</sup>, Alain Rudiger<sup>24</sup>, Frank Ruschitzka<sup>25</sup>, Petar M. Seferovic<sup>26</sup>, Alessandro Sionis<sup>27</sup>, Antoine Vieillard-Baron<sup>28</sup>, Jean Marc Weinstein<sup>29</sup>, Rudolf A. de Boer<sup>30</sup>, Maria G. Crespo-Leiro<sup>31</sup>, Massimo Piepoli<sup>32</sup>, and Jillian P. Riley<sup>33</sup>

<sup>1</sup> Emergency Medicine, University of Helsinki, Department of Emergency Medicine and Services, Helsinki University Hospital, Helsinki, Finland; <sup>2</sup> Athens University Hospital Attikon, Athens, Greece; <sup>3</sup> Department of Cardiology, Maastricht University Medical Center, Maastricht, The Netherlands; <sup>4</sup> Vilnius University, Faculty of Medicine, Institute of Clinical Medicine, Clinic of Cardiac and Vascular Diseases, Vilnius, Lithuania; <sup>5</sup> University of Medicine Carol Davila/Institute of Emergency for Cardiovascular Disease, Bucharest, Romania; <sup>6</sup> Department of Emergency Medicine, Vanderbilt University Medical Center, Nashville, TN, USA; <sup>7</sup> Department of Intensive Care Medicine, CHIREC Hospitals, Université Libre de Bruxelles, Brussels, Belgium; <sup>8</sup> Département d'Anesthésie-Réanimation-SMUR, Hôpitaux Universitaires Saint Louis-Lariboisière, INSERM-UMR 942, AP-HP, Université Paris Diderot, Paris, France; <sup>9</sup> Queens University, Belfast, UK; <sup>10</sup> Department of Internal Medicine and Department of Research and Education, General Hospital Murska Sobota, Murska Sobota, Slovenia; <sup>11</sup> Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia; <sup>12</sup> Cardiology, Heart and Lung Center, University of Helsinki, Helsinki University Hospital, Helsinki, Finland; <sup>13</sup> Consorci Sanitari Integral, University of Barcelona, Barcelona, Spain; <sup>14</sup> Hospital Sanitas CIMA, Barcelona, Spain; <sup>15</sup> U942 INSERM, AP-HP, Paris, France; <sup>16</sup> Investigation Network Initiative Cardiovascular and Renal Clinical Trialists (INI-CRCT), Nancy, France; <sup>17</sup> University Paris Diderot, Sorbonne Paris Cité, Paris, France; <sup>18</sup> AP-HP, Department of Anesthesia and Critical Care, Hôpitaux Universitaires Saint Louis-Lariboisière, Paris, France; <sup>19</sup> Emergency Department, Hospital Clínic, University of Barcelona, Barcelona, Catalonia, Spain; <sup>20</sup> Department of Cardiology, Policlinico di Monza, Monza, Italy; <sup>21</sup> Department of Cardiology and Cardiovascular Research Institute Basel (CRIB), University Hospital Basel, Basel, Switzerland; <sup>22</sup> Department of Cardiology, Ziekenhuis Oost Limburg, Genk – Biomedical Research Institute, Faculty of Medicine and Life Sciences, Hasselt University, Diepenbeek, Belgium; <sup>23</sup> Department of Cardiology, University of Helsinki, Helsinki, Finland; <sup>24</sup> Cardio-surgical Intensive Care Unit, University and University Hospital Zurich, Zurich, Switzerland; <sup>25</sup> University Heart Center, University Hospital Zurich, Zurich, Switzerland; <sup>26</sup> Department of Internal Medicine, Belgrade University School of Medicine and Heart Failure Center, Belgrade University Medical Center, Belgrade, Serbia; <sup>27</sup> Cardiology Department, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain; <sup>28</sup> INSERM U-1018, CESP, Team 5 (EpReC, Renal and Cardiovascular Epidemiology), UVSQ, 94807 Villejuif, France, University Hospital Ambroise Paré, AP-HP, Boulogne-Billancourt, France; <sup>29</sup> Cardiology Division, Soroka University Medical Centre, Beer-Sheva, Israel; <sup>30</sup> Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; <sup>31</sup> Complejo Hospitalario Universitario A Coruña (CHUAC), Instituto de Investigación Biomédica de A Coruña (INIBIC), CIBERCV, UDC, La Coruña, Spain; <sup>32</sup> Heart Failure Unit, Cardiology, G. da Saliceto Hospital, Piacenza, Italy and <sup>33</sup> Imperial College, London, UK

## **Introduction**

Professionals within clinical practice frequently encounter patients hospitalized with acute heart failure (AHF).<sup>1-5</sup> These patients commonly present with signs and symptoms of congestion, but some patients also present with signs of tissue hypoperfusion contributing to impaired organ function.<sup>6</sup> Regardless of their clinical profile or type of heart failure (i.e. reduced or preserved ejection fraction<sup>7</sup>), comprehensive inpatient monitoring in conjunction with thorough physical examination is crucial to optimize management. Clinicians are under increasing pressure to keep inpatient length of stay as short as possible while also preventing post-discharge emergency visits, rehospitalization and short-term mortality. More effective monitoring might help to achieve these goals and improve the accuracy of discharge readiness assessments.

The 2016 European Society of Cardiology (ESC) heart failure guidelines provide recommendations based on the available evidence for the inpatient management of AHF. It is notable that 17 of the 18 recommendations related to monitoring, all class I recommendations, are rated as level of evidence C ('consensus of opinion of the experts and/or small studies, retrospective studies, registries') (*Table 1*).<sup>6</sup> Thus, a substantial evidence gap exists regarding the optimal inpatient monitoring strategies for these severely ill patients.


**Table 1.** ESC guideline recommendations and further considerations for in-hospital monitoring of patients with acute heart failure

Topic	Guideline recommendation <sup>6</sup>
Diagnostic evaluation: baseline and initial monitoring; usually in emergency department or at point of initial admission	
Non-invasive vital signs	Standard non-invasive monitoring of heart rate, rhythm, respiratory rate, oxygen saturation and blood pressure is recommended (class I, level of evidence C).
Biomarkers	Upon presentation, a measurement of plasma natriuretic peptide level (BNP, NT-proBNP or MR-proANP) is recommended in all patients with acute dyspnoea and suspected AHF to help in the differentiation of AHF from non-cardiac causes of acute dyspnoea (class I, level of evidence A).
Electrocardiogram	12-lead ECG is recommended at admission in all patients presenting with suspected AHF (class I, level of evidence C).
Chest X-ray	Chest X-ray is recommended at admission in all patients presenting with suspected AHF to assess signs of pulmonary congestion and detect other cardiac or non-cardiac diseases that may cause or contribute to the patient's symptoms (class I, level of evidence C).
Laboratory assessments	Cardiac troponins, blood urea nitrogen, creatinine, electrolytes (sodium, potassium), glucose, complete blood count, liver function tests, and thyroid stimulating hormone are recommended at admission in all patients presenting with suspected AHF (class I, level of evidence C).
Echocardiography	Echocardiography is recommended immediately in haemodynamically unstable AHF patients and within 48 h when cardiac structure and function are either not known or may have changed since previous studies (class I, level of evidence C).
Clinical status monitoring: usually in intensive or critical care unit; less invasive components of monitoring continued on the general ward or prior to discharge	
Vital signs	Standard non-invasive monitoring of heart rate, rhythm, respiratory rate, oxygen saturation and blood pressure is recommended (class I, level of evidence C).
Body weight and fluid balance	It is recommended that patients should be weighed daily and have an accurate fluid balance chart completed. This includes accurate measurements of urine output in critically ill patients (class I, level of evidence C).
Signs and symptoms	It is recommended to evaluate signs and symptoms relevant to heart failure (e.g. dyspnoea, pulmonary rales, peripheral oedema, weight) (class I, level of evidence C).
Laboratory assessment	Frequent, often daily measurement of renal function (blood urea, creatinine) and electrolytes (potassium, sodium) during i.v. therapy and when renin-angiotensin-aldosterone system antagonists are initiated is recommended (class I, level of evidence C).
Invasive monitoring	Invasive monitoring with an arterial line is recommended in patients with cardiogenic shock (class I, level of evidence C). Intra-arterial line should be considered in patients with hypotension and persistent symptoms despite treatment (class IIa, level of evidence C). Pulmonary artery catheter may be considered in patients who, despite pharmacological treatment, present refractory symptoms (particularly with hypotension and hypoperfusion) (class IIb, level of evidence C).
Oxygenation	Transcutaneous monitoring of arterial oxygen saturation is recommended (class I, level of evidence C).
Blood gas analysis	Measurement of blood pH and carbon dioxide tension (possibly including lactate) should be considered, especially in patients with acute pulmonary oedema or history of COPD using venous blood. In patients with cardiogenic shock, arterial blood is preferable (class IIa, level of evidence C).

AHF, acute heart failure; BNP, B-type natriuretic peptide; COPD, chronic obstructive pulmonary disease; ECG, electrocardiogram; NT-proBNP, N-terminal pro B-type natriuretic peptide; MR-proANP, mid-regional pro-atrial natriuretic peptide.

Despite this lack of evidence, physicians are expected to deploy the best monitoring tools to guide management decisions for their patients. Importantly, monitoring is a dynamic process involving a spectrum of different clinical, biochemical and physiological parameters. These variables should be interpreted in the context of the whole patient, rather than as isolated measurements. Importantly, equal attention should be paid to changes in measurements over time. Moreover, the feasibility, costs, and possible harms of monitoring, as well as the required knowledge and skills for appropriate use of monitoring tools should be considered when optimizing monitoring approaches for an individual patient. This document provides the multidisciplinary team with practical applications of guideline

recommendations for the inpatient monitoring of patients with AHF (Figure 1). Recommendations that are taken from the ESC guidelines are noted by reference and the description of the accompanying recommendation class and level of evidence. Practical suggestions for clinical application do not have an associated recommendation class or level of evidence. Identifying research priorities to address the gaps in evidence (Table 2) to improve the evidence base for future guideline recommendations is also addressed.



**Figure 1.** Suggested inpatient monitoring for patients with acute heart failure. Comprehensive inpatient monitoring is crucial to the optimal management of acute heart failure patients. Many tools are available for in-hospital monitoring of patients with acute heart failure, and each plays a role at various points throughout the patient's treatment course, including the emergency department, intensive care or coronary care unit, and the general ward. As a result of diverse precipitants, left ventricular end-diastolic pressure (LVEDP) and natriuretic peptides (NPs) increase (red curve) and their elevation precedes worsening of congestion (black curve). This pattern may be useful to identify early subclinical decompensation. In spite of symptomatic improvement as a result of in-hospital therapies for a subset of patients, high LVEDP and NP values are still persistent during hospitalization and in the early post-discharge period, and identification of haemodynamic congestion is crucial for preventing early post-discharge adverse events in these patients. Yellow arrows represent continuous monitoring; yellow boxes indicate suggested monitoring repeated at various time points. BP, blood pressure; cTn, cardiac troponin; ECG, electrocardiogram; ECHO, echocardiogram; GDMT, guideline-directed medical therapy; HR, heart rate; IV, intravenous; RR, respiratory rate; SaO<sub>2</sub>, oxygen saturation. \*Invasive BP monitoring and blood gases may be indicated in the intensive or coronary care unit and in most critical patients. †Continuous telemetry monitoring in selected patients. ‡In the *de novo* patient, oral GDMT is started stepwise when the patient is stabilized. In patients with worsening chronic heart failure, ongoing medications should be continued at the highest tolerated dose as appropriate for the clinical situation.

**Table 2.** Gaps in knowledge and priorities for research

Topic	Research priorities
Body weight, urine output, and fluid balance	<ul style="list-style-type: none"> <li>• Impact of daily weight and fluid balance monitoring</li> <li>• Quantify and validate definition for diuretic responsiveness (e.g. urine sodium loss) vs. diuretic resistance</li> </ul>
Electrolytes	<ul style="list-style-type: none"> <li>• Normalization of serum sodium during hospitalization</li> <li>• Guiding diuretics or alternative treatment (e.g. haemofiltration) based on electrolytes (particularly sodium and chloride)</li> </ul>
Markers of kidney function	<ul style="list-style-type: none"> <li>• Prospectively test decongestive strategies guided by kidney biomarker profiles</li> <li>• Evaluate the impact of various strategies (i.e. primarily vasodilator-based vs. primarily diuretic-based) on kidney marker profiles and clinical outcomes</li> <li>• Examine pathophysiology of serum creatinine increases in the context of decongestion</li> <li>• Identify and validate new kidney markers for accurate reflection of renal function in acute settings</li> </ul>
Liver function	<ul style="list-style-type: none"> <li>• Assess utility of liver function markers to evaluate decongestion and guide treatment decisions</li> <li>• Prospectively test whether markers of cholestasis (reflecting congestion) and/or markers of hypoxic hepatitis (reflecting hypoperfusion) offer additional prognostic and therapeutic information to clinical evaluation</li> </ul>
Cardiac biomarkers	<ul style="list-style-type: none"> <li>• Validate the multimarker approach for informing diagnosis and prognosis</li> <li>• Establish the incremental value of a multimarker approach, including the combination of biomarkers with imaging markers</li> <li>• Identify markers of low-risk that can facilitate safe early discharge from the emergency department or hospital</li> <li>• Identify biomarkers to guide therapy selection (i.e. precision medicine) or monitor response to therapy and improve patient outcome</li> </ul>
Rhythm monitoring	<ul style="list-style-type: none"> <li>• Delineate which patients need continuous ECG monitoring</li> <li>• Validate risk prediction tools (e.g. scores, biomarkers, clinical assessment, history, or a combination of these) used to triage patients for 'high-intensity inpatient monitoring' or 'low-intensity inpatient monitoring'</li> </ul>
Echocardiography	<ul style="list-style-type: none"> <li>• Determine the best indices of elevated LAP</li> <li>• Evaluate using LAP indices or cardiac output to guide decongestive therapy</li> <li>• Identify echocardiography findings that indicate a need for inotropic support</li> <li>• Determine criteria that indicate failure of medical therapy and need for mechanical support</li> </ul>
Lung ultrasound	<ul style="list-style-type: none"> <li>• Evaluate lung ultrasound (i.e. changes in the quantity of B-lines) to guide decongestive therapy</li> <li>• Determine utility of lung ultrasound to indicate discharge readiness</li> <li>• Directly compare B-lines with chest X-ray</li> </ul>
Non-invasive monitoring	<ul style="list-style-type: none"> <li>• Validate thoracic bioimpedance to assess volume responsiveness and central haemodynamic response to physiological and pharmacological stimulation</li> <li>• Identify target values for microvascular flow</li> <li>• Determine how therapeutic interventions could be tailored to microvascular flow data</li> <li>• Evaluate the usefulness of microcirculation assessments for guiding treatment decisions</li> <li>• Comparison of direct vs. surrogate measurements of microcirculation</li> </ul>
Risk score	<ul style="list-style-type: none"> <li>• Externally validate risk stratification and prognostic scores</li> <li>• Evaluate generalizability of scores</li> <li>• Determine whether application of risk scores improves patient outcome (i.e. through better risk stratification, application of appropriate therapies or better health care utilization)</li> </ul>

AHF, acute heart failure; ECG, electrocardiogram; LAP, left atrial pressure.

## Clinical monitoring

### Overview

Congestion is present in >90% of patients and is the leading cause of hospitalization in patients with AHF.<sup>6,8,9</sup> Most patients present as 'wet and warm' with normal or elevated blood pressure and adequate perfusion, symptoms (e.g. dyspnoea, orthopnoea, breathlessness on minimal exertion, abdominal discomfort, or loss of appetite), and signs (e.g. pulmonary rales, oedema, elevated jugular venous pressure, hepatomegaly, ascites) of congestion due to either volume overload or fluid redistribution.<sup>6</sup> Since a single clinical symptom or sign may lack sensitivity or specificity, clinical decisions should be based on comprehensive evaluation of all findings.<sup>10</sup> Congestion in the setting of AHF is also associated with organ dysfunction and poor prognosis.<sup>11-13</sup> Far fewer patients present with symptoms or clinical signs of hypoperfusion or cardiogenic shock (e.g. cool sweaty extremities, oliguria, mental confusion, dizziness, hypotension, narrow pulse pressure), but these situations require urgent interventions to be taken to prevent ensuing death.<sup>14</sup> In the ESC Heart Failure Long-Term Registry, 70% presented with congestion without hypoperfusion, 13.6% with congestion and hypoperfusion, and 0.9% with hypoperfusion without congestion.<sup>15</sup>

The clinical picture may be more dramatic in new-onset, acute *de novo* patients than in patients with chronic heart failure who present with gradually worsening dyspnoea. In the EuroHeart Failure Survey II, the prevalence of pulmonary oedema was 26% in patients with *de novo* AHF and 10.4% in patients with acutely decompensated chronic heart failure. Cardiogenic shock occurred in 6.8% of acute *de novo* heart failure patients and in 2.2% of acutely decompensated chronic heart failure patients.<sup>8</sup>

The initial primary therapeutic objectives in AHF are the control of symptoms and relief of congestion and fluid overload while preventing further organ injury.<sup>6,13</sup> The ESC guidelines recommend daily assessment of weight and accurate fluid balance charting.<sup>6,16</sup> While perhaps intuitive, data are scarce linking these practices with improved management or outcome.

### Non-invasive vital sign monitoring

In accordance with the ESC guidelines, non-invasive monitoring of blood pressure, respiratory rate, and pulse oximetry should be performed at the time of initial presentation (class I, level of evidence C).<sup>6,17</sup> Although not specified in the guidelines, these measurements should be repeated at intervals based on the changing clinical status of the patient. Rapid changes in vital signs may indicate multi-organ injury.<sup>13</sup> For those more severely ill or with a very dynamic clinical profile, this may be every 15 min. In those patients whose profile changes gradually and who are less severely ill, this could be as infrequent as three times daily. Of note, recent advances in pulse oximetry technology may provide broader information, including perfusion index and some indirect indices of cardiac output,<sup>18</sup> but more research is needed.

### Body weight, urine output, and fluid balance

Monitoring weight change and targeting a negative net fluid balance assumes that congestion is always accompanied by fluid overload and weight gain. However, as many as 50% of patients with AHF are hospitalized without significant weight gain in the days to weeks before presentation.<sup>19-21</sup> Pulmonary congestion, manifested as a rise in left- and right-sided filling pressures, often can occur without significant weight gain or even relative hypovolaemia through volume redistribution.<sup>13,22,23</sup> Thus, first establishing the patient's volume phenotype (i.e. redistribution vs. overload recognizing there is overlap) is critical to facilitate the value of body weight monitoring. Nevertheless, it is reasonable to monitor weight (class I, level of evidence C) and targeting a specific euvolaemic weight may be appropriate in many patients. Weight gain that immediately precedes a hospitalization may help identify patients who

have been non-adherent to diet or medications, and reinforcement of education during the in-hospital stay may be warranted for such patients. Weight gain in hospital is a predictor of 30-day death or rehospitalization.<sup>24</sup>

Exact urine output measurement and accurate fluid charting are basic measures in intensive and cardiac care units. However, in regular wards these are more susceptible to errors. Moreover, calculation of net fluid balance should also include evaporation, which at best is an estimate. Exact and continuous measurement of urine output requires an indwelling urinary catheter, which is associated with patient discomfort, infection risk, and long-term complications such as incontinence.<sup>6,17</sup> The ESC guidelines do not recommend the routine use of a urinary catheter.<sup>6</sup> Intensive care unit (ICU) patients are an exception in whom more intensive monitoring most often includes a more precise evaluation of urine output.

In case of incontinence, weighing an incontinence pad<sup>25</sup> or use of disposable urinary sheaths in males<sup>26</sup> may be a better approach than estimation. Emerging data suggest that urinary characteristics (i.e. urine sodium excretion/mg loop diuretic administered) can indicate either diuretic efficiency since sodium content/urine volume tends to drop quickly during decongestion, or diuretic resistance.<sup>27-29</sup> Other quantitative measures of diuretic response, such as weight or net fluid loss per unit of diuretic (e.g. per 40 mg furosemide or equivalent), have also been proposed to assess response to decongestive therapy. There are no single cut-offs for these indices; in general, the higher the values, the better the response. Haemoconcentration has also been proposed as a measure of decongestion.<sup>30</sup> Poor diuretic response in patients with AHF has been associated with less symptom relief, in-hospital worsening of heart failure, increased mortality, and higher rehospitalization rates.<sup>28,31-33</sup>

## **Biomarker monitoring**

### *Arterial blood gas analysis and acid–base balance*

Arterial blood gas analysis provides vital information on oxygenation, ventilation, metabolic balance, an indication of tissue hypoxia. The selective use of these may assist with determining prognosis. It is usually performed on point of care analyzers measuring the partial pressures of oxygen (pO<sub>2</sub>), carbon dioxide (pCO<sub>2</sub>) and pH. Haemoglobin oxygen saturation (SpO<sub>2</sub>) can be either calculated or measured directly. Bicarbonate and base deficit/excess are derived values from pH and pCO<sub>2</sub>. Arterial blood gas analysis is generally restricted to patients with pulmonary oedema, severe AHF (on the verge of needing artificial ventilation), cardiogenic shock, or a history of chronic obstructive pulmonary disease (class IIa, level of evidence C) and can be obtained hourly as dictated by the patient's changing condition.<sup>6</sup>

Practical interpretation of arterial blood gas measurements consists of assessing pH and the respiratory and metabolic (i.e. base deficit/excess) components, calculating the anion gap in the presence of metabolic acidosis, and evaluating the primary disturbance and compensatory response. The most common acid–base imbalance in AHF is respiratory alkalosis, but metabolic acidosis, most often as a sign of poor end-organ perfusion, has been associated with an increased mortality risk.<sup>34</sup> Of note, both metabolic acidosis and alkalosis are associated with increased mortality.<sup>35</sup> The prognostic value of admission pO<sub>2</sub> and pCO<sub>2</sub> remains controversial.<sup>36</sup> The ratio of pO<sub>2</sub>/fraction of inspired oxygen gives a better impression of the adequacy of pulmonary gas exchange than pO<sub>2</sub> alone. A high correlation has been demonstrated between pCO<sub>2</sub> and pH measured in venous and arterial blood;<sup>37,38</sup> thus, venous blood gas analysis is a very good alternative method, especially in those patients in whom pulse oximetry is over 90%.

The veno–arterial gradient in pCO<sub>2</sub> measured in central lines can be used as a marker of tissue perfusion, although not specifically recommended in the ESC guidelines. Veno–arterial pCO<sub>2</sub> gradient is affected by microvascular perfusion, as well as metabolism. As flow adapts to metabolism, an increased veno–arterial gradient in pCO<sub>2</sub> (>6 mmHg) suggests that tissue perfusion is inadequate.<sup>39</sup> An increased veno–arterial pCO<sub>2</sub> gradient has been associated with poor outcome.<sup>40</sup>

## ***Lactate***

Lactate, measured from either venous or arterial blood, is a very important indicator of tissue hypoperfusion. Elevated lactate levels may result from either excess production, impaired lactate clearance (e.g. liver dysfunction), or both. While lactate is often of anaerobic origin in the early hours after the onset of circulatory failure, other sources of lactate predominate at later stages.<sup>41</sup> Hence, lactate is a good marker of tissue perfusion mostly in the early stages of circulatory failure. Elevations in lactate can cause metabolic acidosis, and lactic acidosis is a common cause of raised anion gap. Lactate concentrations >2 mmol/L are generally considered as abnormal and associated with higher mortality, but levels at the higher end of the normal reference range (>0.75 mmol/L) have also been associated with slightly increased mortality in patients admitted to the ICU.<sup>42</sup> Furthermore, lactate levels that do not decrease following appropriate treatment are associated with a poor outcome.<sup>43,44</sup> The optimal measurement intervals are not specifically described in the guidelines, but practically, lactate levels should be assessed in haemodynamically unstable or hypoxaemic patients with AHF at the time of admission and repeated at shorter intervals initially (e.g. every 1–2 h) during the acute phase of hospitalization. The lactate time interval can then be increased as the patient stabilizes.<sup>44</sup>

## ***Electrolytes***

Electrolytes should be routinely measured on admission in patients with AHF (class I, level of evidence C) together with serum creatinine and urea nitrogen (*Table 1*).<sup>6</sup> During administration of vasoactive and diuretic therapy, electrolytes should be reassessed at least daily until discharge, particularly if intravenous loop diuretics are used, with more frequent testing if clinically indicated<sup>6</sup> (e.g. with deteriorating renal function or initiation of renin–angiotensin–aldosterone system antagonists). Serum hyponatraemia at admission predicts poor prognosis in patients hospitalized for AHF.<sup>45,46</sup> Additional testing is needed to distinguish between depletion and dilutional hyponatraemia as this will guide further therapy.<sup>47</sup> Hypo- or hyperkalaemia are associated with arrhythmias (e.g. QT prolongation, torsades de pointes, or other life-threatening arrhythmias), and confirmed abnormal potassium levels should be corrected. Hypokalaemia as well as hypomagnesaemia are risk factors for new-onset atrial fibrillation.<sup>48</sup> Chloride plays a role in renal salt sensing, neurohormonal activation, and regulation of diuretic targets, and hypochloraemia predicts mortality in acute and chronic heart failure.<sup>49</sup>

## ***Markers of kidney function***

The optimal measurement of renal function is uncertain, particularly in the acute setting and interpretation should consider the absolute value as well as the change from baseline. The use of serum creatinine to estimate glomerular filtration rate (eGFR), which is only one of the three major aspects of kidney physiology (filtration, absorption, secretion) has known limitations (e.g. imprecision in non-steady state situations, influenced by muscle mass, tubular secretion affected by certain drugs, influenced by diet),<sup>50,51</sup> but it remains the standard in most hospitals. Measuring serum creatinine is inexpensive, and studies of other renal markers (e.g. cystatin C, neutrophil gelatinase-associated lipocalin) have not generated sufficient evidence of superiority over repetitive serum creatinine measurements.

Frequent, most often daily, monitoring of renal function is recommended by the current guidelines when intravenous treatments are administered or when renin–angiotensin–aldosterone system antagonists are initiated (class I, level of evidence C; *Table 1*), although guidelines do not give specific recommendations for responding to these measurements (*Table 3*).<sup>6</sup> Importantly, decreases in eGFR most often occur during the first days of hospitalization, which suggests they are caused either by administered treatments or were secondary to congestion and already in the process of worsening before admission.<sup>52,53</sup> Decongestion can result in either improvement or worsening of renal function, such as an initial improvement followed by decline or vice versa. Worsening renal function is associated with poor outcome both in acute and chronic heart failure.<sup>54</sup> Importantly, small (<15–20%) changes in eGFR can be accepted if the patient is effectively being decongested, a concept based on recent data that suggest such



changes are not associated with worse outcome.<sup>55,56</sup> Additionally, most of the changes observed are related to alterations in intraglomerular haemodynamics that result from differences in the tonus of the afferent and efferent glomerular arterioles and are not linked with adverse outcomes.<sup>57</sup> However, in the case of worsening eGFR, decreasing urinary output, and persistent congestion, further actions need to be taken.<sup>57</sup> Congestion is an important driver of worsening renal function and, as outlined previously, also related to worse prognosis.

**Table 3.** Potential use of biomarkers in acute heart failure

Potential clinical use	Biomarkers	Evidence level*	Current role in clinical practice	Limitations
Diagnosis of AHF	NPs	Level 1	Well established, recommended†	NPs also influenced by age, obesity, renal function, atrial fibrillation, concomitant medications (e.g. ARNI and BNP)
Diagnosis of concomitant diseases	Infection (CRP, WBC, PCT)	Level 3	Well established, recommended†	
	Renal dysfunction (creatinine, urea, others)	Level 3	Well established, recommended†	
	Anaemia (Hb, ferritin, transferrin saturation)	Level 3	Well established, recommended†	
	Liver function (AST, ALT, alkaline phosphatase, GGT, INR), lactate	Level 3	Well established, recommended†	
Risk stratification	Many biomarkers (e.g. NPs, MR-proADM, troponins, sST2, GDF-15, renal markers, multi-markers)	Level 4	NP and troponin generally available; other markers have limited clinical availability in some locations; risk scores should be applied when available	Useful for assessing prognosis, but lack of evidence that biomarkers facilitate clinical actions that improve outcome
Select treatment or predict response to treatment	PCT for use of antibiotics in AHF	Level 3	Currently investigational	Limited evidence, prospective study ongoing
Determine treatment effects	NPs, lactate, sST2, haemoconcentration	Level 4	No general recommendation	Treatment-induced changes in biomarkers not yet directly linked to clinical outcome
Monitor safety	Renal function, liver function	Level 5	Highly recommended†, but expert opinion only. No recommendations on how to act based on biomarker(s)	Current renal function markers (i.e. SCr) do not adequately reflect changing eGFR in acute settings
Monitor post-discharge outcome	NPs	Level 4	Recommended† to measure at discharge, but clinical consequences unclear	NPs also influenced by age, obesity, renal function, atrial fibrillation, concomitant medications (e.g. ARNI and BNP)
	Other biomarkers such as sST2, Gal-3, GDF-15	Level 5	Limited clinical availability in some locations or investigational	

AHF, acute heart failure; ALT, alanine aminotransferase; ARNI, angiotensin receptor–neprilysin inhibitor; AST, aspartate aminotransferase; BNP, B-type natriuretic peptide; CRP, C-reactive protein; eGFR, estimated glomerular filtrate rate; GGT, gamma-glutamyl transferase; Gal-3, galectin-3; GDF-15, growth differentiation factor-15; Hb, haemoglobin; INR, international normalized ratio; MR-proADM, mid-regional pro-adrenomedullin; NP, natriuretic peptide; PCT, procalcitonin; SCr, serum creatinine; sST2, soluble ST2; WBC, white blood cells.

\* Level 1: different prospective studies showing that biomarker(s) change management of patients with positive effect on outcome.

Level 2: one prospective study showing that biomarkers may change management with positive effect on outcome.

Level 3: retrospective studies suggesting that biomarker may change management, but not yet prospectively tested.

Level 4: different prospective studies showing positive findings, but no consequences of changes in therapy/management tested.

Level 5: retrospective studies only/limited results with no impact on management/therapy.

† Recommended by the authors.

### ***Markers of liver function***

Liver function tests should be performed on admission in accordance with current guidelines (class I, level of evidence C).<sup>6</sup> Liver dysfunction in AHF is most commonly the result of elevated central venous pressure and manifested by increases in alkaline phosphatase and gamma-glutamyl transferase<sup>58</sup> or total bilirubin,<sup>59</sup> these abnormalities often improve with decongestion. Elevated transaminases occur with hypoxic liver injury in the setting of hypoperfusion, which is less common, but associated with worse outcome.<sup>13</sup> Alternative causes for liver dysfunction should be sought if liver function tests do not normalize after decongestion has been achieved and perfusion corrected.

### ***Cardiac biomarkers***

Cardiac biomarkers are evidence-based tools for the diagnosis of AHF and determining prognosis. The utility of biomarkers to predict response to therapy, determine treatment effects, and monitor changes in disease processes or organ-specific function over time is an area of intense interest and research (*Table 3*).<sup>3</sup> Recently, recommendations for their use was discussed by the American Heart Association,<sup>60</sup> but definitive evidence of their role in clinical practice for these purposes has not been confirmed.<sup>61</sup>

### ***Natriuretic peptides***

The ESC guidelines recommend measuring a natriuretic peptide level at admission for all patients with acute dyspnoea and suspected AHF. This recommendation is based on a strong body of evidence (class I, level of evidence A; *Table 1*).<sup>6</sup> Clinicians should be aware of factors that can confound interpretation of natriuretic peptide values (e.g. renal impairment, atrial fibrillation, other co-morbidities, age, obesity). Pre-discharge natriuretic peptide levels are also useful for prognosis,<sup>6</sup> and pre-discharge levels have demonstrated stronger prediction of post-discharge outcomes than admission levels.<sup>62-64</sup> Thus, pre-discharge assessment of natriuretic peptides may be informative in determining readiness for discharge since a reduction of natriuretic peptide levels after decongestive therapy portends a better prognosis, but it has not yet been tested if this approach causally improves outcome and therefore it is not a specific guideline recommendation. A >30% decrease in N-terminal pro B-type natriuretic peptide (NT-proBNP) from admission was associated with a lower risk of subsequent death or readmission compared to patients without this threshold NT-proBNP decrease.<sup>65</sup> In patients who, based on the clinician's judgment, fail to adequately reduce BNP, a more intensified follow-up might be warranted.

While using natriuretic peptide levels to guide treatment is of substantial interest, it has not improved clinical outcomes compared to standard care in randomized studies to date;<sup>66</sup> most guided studies have been in the setting of chronic heart failure. In some randomized trials evaluating treatments for AHF, natriuretic peptide and other biomarker profiles improved after treatment with the agent under study, but long-term outcomes were not improved. For example, despite favourable effects on biomarkers, patient outcomes were not improved in the SURVIVE (Survival of Patients with Acute Heart Failure in Need of Intravenous Inotropic Support),<sup>67</sup> RELAX-AHF-2 (Relaxin in Acute Heart Failure-2),<sup>68</sup> or TRUE-AHF (Trial of Ularitide Efficacy and Safety in Acute Heart Failure)<sup>69</sup> trials. Post-hoc analyses have shown an association between decreasing natriuretic peptide levels during an admission for AHF and all-cause mortality after multivariate adjustment that included treatment allocation.<sup>70</sup> These findings provide more evidence that clinical outcomes in AHF are influenced by multiple pathways, some of which may not be primarily related to heart failure, and natriuretic peptides alone do not appear to sufficiently reflect all relevant processes that contribute to mortality.

### *Cardiac troponin*

Cardiac troponin obtained at admission in patients with AHF (class I, level of evidence C) is useful to evaluate the possibility of acute coronary syndrome as a precipitating factor,<sup>6,71</sup> but a dynamic change pattern should be observed to confirm the acute myocardial infarction diagnosis according to current acute coronary syndrome guidelines.<sup>72</sup> The majority of patients presenting with AHF will have elevated cardiac troponin levels, and although elevated cardiac troponin is associated with adverse prognosis,<sup>6</sup> it does not necessarily indicate an acute coronary event if other criteria for acute myocardial infarction are absent. Current evidence therefore does not support the role for serial cardiac troponin monitoring in patients with AHF without an acute coronary syndrome.

### *Other innovative markers*

Many new biomarkers have emerged in the field of AHF, but several studies have suffered from inadequate or poorly described methods, optimistically interpreted results, publication bias, and mixing of opinion vs. fact.<sup>73</sup> Thus, guideline recommendations have not yet been made regarding their use, but future studies should determine their role. Even more importantly, no study has tested specific strategies for altering AHF treatment based on biomarker profiles.

Most biomarker studies in AHF have investigated the prognostic value of single biomarkers, but the MOCA (Multinational Observational Cohort on Acute heart failure) study showed that dual biomarker combinations improved risk stratification over a single biomarker for 30-day [C-reactive protein (CRP) in combination with mid-regional pro-adrenomedullin, soluble ST2 (sST2), or NT-proBNP] and 1-year survival (CRP and sST2 reclassified risk to the greatest extent).<sup>74</sup> Similar improvements in prognostication using combined biomarkers have been demonstrated in other studies.<sup>75</sup> However, use of these biomarker panels in clinical practice will depend on their routine availability.

The detection of important co-morbidities is a promising application for biomarkers in patients with AHF, not only for risk assessment but primarily to assess the need for treatment of co-morbidities (e.g. detection of bacterial infection such as pneumonia). Procalcitonin is more specific for infection than other inflammatory markers (e.g. CRP), and retrospective data suggest that both untreated bacterial infection and unnecessary antibiotic treatment in AHF are associated with worse outcome.<sup>76</sup> IMPACT-EU (Improve Management of Heart Failure with Procalcitonin, NCT02392689) is an ongoing prospective study of the effect of procalcitonin-guided antibiotic therapy vs. physician-determined antibiotic therapy on the primary endpoint of 90-day all-cause mortality. If effective, this approach would be one of the first proven biomarker-guided treatment approaches in AHF.

## **Electrocardiography and telemetry**

Rhythm disturbances and ischaemia may be a primary cause of an AHF episode.<sup>77</sup> Conversely, haemodynamic instability may give rise to arrhythmias, or these may be iatrogenic in origin (e.g. secondary to inotrope or vasopressor administration).

Obtaining an electrocardiogram (ECG) during the initial evaluation is routine practice (class I, level of evidence C).<sup>6</sup> Telemetry monitoring is recommended by the American Heart Association in all patients with AHF who are at significant risk of an immediate, life-threatening arrhythmia and should be continued until the patient is stabilized.<sup>78</sup> Telemetry monitoring may also help with detecting episodes of ischaemia. Patients hospitalized for AHF commonly have multiple co-morbidities, and telemetry monitoring may be recommended with concomitant conditions such as acute respiratory failure, shock, acute pulmonary embolus, or renal failure with electrolyte abnormalities (e.g. hyperkalaemia) where the risk of arrhythmia is increased. It is also recommended in patients on inotropic support.

Telemetry monitoring is presumed to improve patient safety, but there is no definitive evidence that it improves patient outcomes.<sup>79-81</sup> Continuous telemetry beds are increasingly available in many hospitals. Risk stratification tools to detect high-risk patients and specific protocols for them may be helpful to triage which patients need telemetry to ensure efficient resource utilization since continuously monitoring all patients with AHF may be inappropriate and not cost-effective.

## **Imaging**

### ***Chest X-ray***

Chest X-ray can be an informative test in the evaluation of patients with suspected AHF, but its main utility is in identifying alternate causes for the patients' symptoms. Pulmonary venous congestion, pleural effusion, interstitial or alveolar oedema and cardiomegaly are the most specific findings for AHF.<sup>82,83</sup> However, up to 20% of patients with AHF will have no or little congestion visible on their emergency department chest radiograph;<sup>84</sup> thus, chest X-ray alone cannot rule out AHF. Supine chest radiographs have limited value in the evaluation of patients with AHF.<sup>85,86</sup> A chest X-ray is recommended at admission in patients with suspected AHF (class I, level of evidence C). To avoid unnecessary irradiation, serial chest X-rays are not routinely recommended.<sup>87</sup> Evidence is lacking to support monitoring patients with AHF by chest X-ray, and few data are available comparing chest X-ray to other clinical and haemodynamic tools. There is old evidence from patients with chronic cardiomyopathy showing decreasing wedge pressure<sup>88</sup> as well as clinical improvement<sup>86</sup> before remission of chest X-ray findings. Repeat chest X-rays should be reserved for patients with a clearly worsening clinical condition or in whom concurrent pulmonary diseases are suspected.<sup>6</sup> Other imaging modalities (e.g. lung ultrasound) may be used as an alternative to repeat chest X-ray to assess congestion or diagnose pleural effusions.<sup>89</sup>

Distinguishing between pulmonary oedema and acute respiratory distress syndrome is one of the most challenging differential diagnoses in the emergency department and the ICU.<sup>90</sup> Cardiogenic pulmonary oedema is the main radiographic differential diagnosis, but acute respiratory distress syndrome is favoured when cardiomegaly, septal lines, or pleural fluid are absent, and peripheral alveolar opacities are present.<sup>91</sup> The Berlin definition of acute respiratory distress syndrome consists of timing, chest imaging (chest X-ray or computed tomography scan), origin of oedema, and degree of oxygenation.<sup>92</sup> In practice, acute respiratory distress syndrome and cardiogenic pulmonary oedema can coexist. Discrimination between the two is often difficult using chest radiography alone, but it may be improved by echocardiography to detect cardiac failure.<sup>91,93</sup>

### ***Echocardiography***

Specific guideline recommendations for the use of echocardiography in AHF are provided in *Table 1*. Echocardiography may also be useful for monitoring filling pressure and cardiac output; research supporting this role is further described below. The value of strain imaging in the monitoring of AHF requires further research.

#### ***Echocardiography for monitoring filling pressures***

There is little evidence to date of either clinical utility or diagnostic accuracy of Doppler echocardiography indices for monitoring left ventricular filling pressures in the setting of AHF. Only a few studies simultaneously performed invasive and non-invasive estimation of left ventricular filling pressures.<sup>94-98</sup>

The American Society of Echocardiography (ASE) and the European Association of Cardiovascular Imaging (EACVI) proposed an algorithm for estimating mean left atrial pressure based on a comprehensive assessment of diastolic indices (*Table 4*).<sup>99</sup> It was originally based on expert consensus but recently has been validated in a multicentre European study against invasive measurement of left ventricular filling pressure.<sup>100</sup> Solely assessing mitral annular velocities is poorly correlated with left ventricular filling pressures in several clinical scenarios (e.g. patients with coronary artery disease which affected basal septal or lateral segments, significant mitral annular calcification, surgical rings or prosthetic mitral valves, left bundle branch block, right ventricular pacing, cardiac resynchronization therapy, and pericardial disease).<sup>99</sup> Several studies have also challenged the utility of E/e' ratio for estimation of pulmonary capillary occluded pressure in patients with advanced or decompensated heart failure with reduced ejection fraction.<sup>98,101</sup>

**Table 4.** Echocardiography parameters to estimate cardiac pressures

Parameter and findings	Interpretation
For estimation of LAP	
E/A ratio <0.8 Peak E velocity $\leq 50$ cm/s <sup>99</sup>	Normal or low
E/A ratio $\geq 2$ Deceleration time <160 ms (in patients with reduced LVEF) <sup>99</sup>	Elevated LAP
Average E/e' <8 <sup>99</sup>	Normal or low LAP
Average E/e' >14 <sup>99</sup>	Elevated LAP
Increased systolic and diastolic PAP estimated by CW Doppler TR systolic jet velocity	Elevated in approximately 70% of HF cases <sup>157</sup> (match of left and right); can also be encountered in normal or low LAP
E/A ratio >0.8 and <2 <sup>99</sup>	Recommended assessment of peak velocity of TR jet by CW Doppler, E/e' ratio, and LA volume index for accurate evaluation
For estimation of RAP	
IVC <2.1 cm that collapses >50% with a sudden inspiratory manoeuvre (i.e. sniff) <sup>158</sup>	Suggests normal RAP of 3 mmHg (range 0–5 mmHg) <sup>158</sup>
IVC diameter >2.1 cm that collapses <50% with a sniff <sup>158</sup>	Suggests high RAP of 15 mmHg (range 10–20 mmHg) <sup>158</sup>
Scenarios in which IVC diameter and collapse do not fit the above paradigms <sup>158</sup>	An intermediate value of 8 mmHg (range 5–10 mmHg) may be used or, preferably, other indices of RAP should be integrated to downgrade or upgrade to the normal or high values of RAP <sup>158</sup>
IVC $\leq 12$ mm (in patients with positive pressure ventilation degree of IVC collapse cannot be used) <sup>159</sup>	RAP <10 mmHg <sup>159</sup>
Hepatic vein flow pattern Vs >Vd (systolic predominance in hepatic vein flow) <sup>158</sup>	Low or normal RAP <sup>160</sup>
Hepatic vein systolic filling fraction (VTIs/VTIs + VTId) <55% <sup>160</sup> or Vs/Vd <1 <sup>158</sup> (lost systolic predominance)	Elevated RAP <sup>160</sup>

CVP, central venous pressure; CW, continuous wave; HF, heart failure; IVC, inferior vena cava; LAP, left atrial pressure; LV, left ventricular; LVEF, left ventricular ejection fraction; PAP, pulmonary artery pressure; RAP, right atrial pressure; TR, tricuspid regurgitation; Vd, diastolic wave; Vs, systolic wave; VTI, velocity–time integral.

Despite these limitations, recent ASE guidelines favour the use of comprehensive echocardiography assessment to monitor therapeutic interventions and indicate that mitral inflow velocities and E/e' are reliable indices of left atrial pressure.<sup>102</sup> Besides, indices of diastolic dysfunction may reflect pathogenic mechanisms related to vascular and cardiac stiffness independent of filling pressures.<sup>103</sup> In the setting of atrial fibrillation, a recent review showed that diastolic parameters such as E/e', isovolumetric relaxation time, and E/Vp correlated with invasive filling pressure and demonstrated adequate reproducibility. For optimal acquisition in atrial fibrillation patients, cardiac cycles with controlled heart rate (<100 b.p.m.) and similar preceding and pre-preceding RR intervals are required.<sup>104</sup> A restrictive mitral inflow pattern on Doppler echocardiography had a diagnostic accuracy for heart failure of 91% in patients hospitalized with acute dyspnoea.<sup>105</sup> Repeat Doppler examination after initiation of diuretic and vasodilator therapy and clinical improvement often showed markedly different findings from baseline,<sup>105</sup> suggesting that it might be a tool to monitor response to therapy. In a single-centre study of 60 patients, the significant decrease in E/e' was observed as early as within the first 12 h of treatment with further decline during 48 h, contrary to the delayed response of inferior vena cava (IVC) and BNP values.<sup>106</sup> Change in pulmonary capillary occluded pressure was strongly correlated with change in E/e' at 48 h but not with BNP in 50 patients admitted to intensive care,<sup>94</sup> although many factors other than left atrial pressure can affect BNP levels. Non-invasive indices should be used cautiously in patients with heart failure with preserved ejection fraction because of the high variability of individual linear regression slopes relating E/e' and E/Vp with invasive measures. Nevertheless, these indices have been useful to identify a cardiac source in patients who fail to wean from mechanical ventilation.<sup>107,108</sup> Importantly, filling pressures do not always correlate with volume overload; thus, diuretic therapy cannot solely be guided by echocardiographic examination.<sup>6,109</sup>

Inferior vena cava diameter and respiratory variation can be used to estimate right atrial pressure, being similarly altered in heart failure with reduced or preserved ejection fraction<sup>7</sup> (*Table 4*). In patients with advanced heart failure, echocardiographic right atrial pressure prediction methods showed modest precision; a single IVC diameter measurement was non-inferior to complex prediction methods.<sup>110</sup> Nagueh *et al.*<sup>95</sup> also demonstrated a strong correlation between invasive and non-invasive estimates of mean right atrial pressure in patients with acute decompensated heart failure. A recent systematic review<sup>111</sup> showed that the diagnostic accuracy of IVC measurement parameters was generally high, with the C-statistic ranging from 0.76–0.91 for IVC diameter and 0.66–0.93 for IVC collapsibility index.<sup>111</sup> The novel 3D echocardiography right atrial volume index has been proposed and validated in conjunction with IVC diameter as having an incremental value for detection of elevated right atrial pressure in patients with AHF.<sup>112</sup> Interestingly, assessment of renal venous flow might be an emerging technique to assess early changes in renal response to volume overload before overt congestion is evident.<sup>113</sup> The reasonable diagnostic performance of pre-specified cut-points justifies the current guideline recommendations<sup>99</sup> for using IVC diameter to estimate right-sided filling pressure.

#### *Echocardiography for monitoring cardiac output*

Echocardiography is capable of monitoring cardiac output, although discontinuously, using pulsed-wave Doppler in the left ventricular outflow tract. Either transthoracic or transoesophageal echocardiography may be used to estimate cardiac output. However, few studies have validated the accuracy of echocardiography compared to the 'gold standard' pulmonary artery catheter (PAC).<sup>114</sup> One recent study in 38 mechanically ventilated patients found high correlation ( $r=0.95$ ) between cardiac output measured by PAC and cardiac output measured by transthoracic echocardiography.<sup>115</sup> The techniques are not interchangeable when measuring cardiac output, but they may be interchangeable to follow directional changes in cardiac output.<sup>114</sup>

Echocardiography has been strongly recommended by consensus groups in patients presenting with circulatory failure.<sup>116,117</sup> It should be noted that echocardiography allows performance of the full haemodynamic evaluation, including chamber volumes, systolic and diastolic function, and heart–lung interactions, filling pressure, and cardiac output.

## ***Lung ultrasound***

Lung ultrasound can be performed at the bedside using portable devices; thus, its application to both general medicine and emergency or intensive settings has expanded considerably. The performance of lung ultrasound is better than chest X-ray to rule out interstitial and alveolar oedema.<sup>89,118</sup>

Many aetiologies are possible in patients presenting with acute dyspnoea. Lung ultrasound can stratify patients according to the likely underlying pathology based on visualization of B-lines.<sup>89</sup> B-lines are an important and reliable marker of pulmonary congestion, but it is difficult to identify the source of pulmonary oedema (hydrostatic or non-hydrostatic) with lung ultrasound alone. Pleural effusion is also reliably detected with lung ultrasound, with good performance and a better detection threshold than chest X-ray. The combination of lung ultrasound and point-of-care natriuretic peptide testing may be helpful in the immediate evaluation of patients with dyspnoea in the emergency department.<sup>119</sup> Lung ultrasound is helpful to differentiate cardiac vs. non-cardiac causes of acute dyspnoea in the emergency department.<sup>120</sup> B-lines appear to be dynamic in nature, as they decrease in proportion to the amount of fluid removed in patients undergoing dialysis.<sup>121</sup> They have also been shown to decrease with treatment of AHF.<sup>106,122-124</sup> According to a recent systematic review, B-line number changed within as few as 3 h of heart failure treatment.<sup>123</sup> However, the number of studies and sample sizes were small and investigator blinding was inadequate.<sup>123</sup>

## **Invasive monitoring**

### ***Arterial, central venous, and pulmonary artery catheters***

The ESC guidelines recommend the insertion of an arterial line in patients with AHF and clinical evidence of cardiogenic shock (class I, level of evidence C; *Table 1*).<sup>6</sup> The arterial line allows for repetitive sampling of arterial blood gases, providing important information on oxygenation (PaO<sub>2</sub>), ventilation (PaCO<sub>2</sub>), acid–base balance, electrolytes and lactate levels.<sup>17</sup> The continuous measurement of arterial pressure allows for the appropriate titration of vasoactive medication, if needed. Respiratory variations of invasive arterial pressure might indicate, among others, right ventricular failure or pericardial tamponade.

The central venous catheter enables the monitoring of central venous pressure and allows the safe and continuous administration of vasoactive drugs and inotropes in patients with AHF who require intensive treatment. Central venous oxygen saturation (ScvO<sub>2</sub>) can also be monitored with the central venous catheter.<sup>125</sup> ScvO<sub>2</sub> declines when oxygen delivery decreases and, particularly in patients with AHF, when cardiac output deteriorates. In combination with increased lactate levels and signs of organ dysfunction, a ScvO<sub>2</sub> <60% indicates severe hypoperfusion and mandates further diagnostics and urgent treatment.

There is no agreement on the optimal method of extended haemodynamic monitoring in assessing and treating the patient in cardiogenic shock.<sup>17</sup> One option is the PAC, i.e. Swan–Ganz catheter. A PAC may be considered in patients who, despite pharmacological treatment, present with refractory symptoms, particularly with hypotension and hypoperfusion (*Table 1*).<sup>6</sup> However, data from the PAC-Man (Pulmonary Artery Catheters in Management of Patients in Intensive Care) study did not support either benefit or harm from use of a PAC in critically ill patients (decompensated heart failure 11% of the population).<sup>126</sup> Furthermore, the ESCAPE (Evaluation Study of Congestive Heart Failure and Pulmonary Artery Catheterization Effectiveness) trial in patients hospitalized for decompensated heart failure but not in cardiogenic shock showed that though the rate of effective decongestion was increased, the use of a PAC did not impact the number of days alive and out of the hospital at 6 months, but it was associated with more adverse events.<sup>127</sup> Based on these data and the imbalance between potential benefits and known risks, PAC should not routinely be used to monitor patients with AHF, but it can still be justified to use PAC in selected populations. PAC is most appropriately used by experienced physicians in clinically compromised situations, where rapid evaluation of vasoactive medications or fluid balance is needed.



## **Risk scores**

Risk scores can be useful to identify the lowest risk patients who could safely be managed in an observation unit or discharged and followed in the outpatient setting in contrast to identifying high-risk patients who require hospital admission and to determine prognosis.<sup>128,129</sup> Most risk scores were designed to estimate high risk, but low risk is not always simply the inverse of high risk. Calculating low risk to some extent is more useful than calculating high risk, as low-risk patients may be safely discharged, saving expensive hospital admission.<sup>130</sup> The same concept applies to hospitalized patients, in order to get a better selection of the best moment for hospital discharge to minimize the risk of rehospitalization.

Clearly, risk scores must be developed in patient cohorts matching the setting where they will be applied, as they are not interchangeable. A recent review found 11 risk scores (nine for hospitalized patients and two for emergency department patients) that could be applied in patients with AHF.<sup>131</sup> Two additional scores for use in the emergency department have subsequently been developed.<sup>132,133</sup> Although some of these risk scores reached promising high discriminatory capacity (C-statistic for EHMARG risk score 0.807; C-statistic for MEESSEI-AHF risk score 0.836), none have been sufficiently validated or prospectively tested to support widespread use.<sup>129</sup>

## **Role of nursing in clinical monitoring**

Nurses are involved across the continuum of care from early identification of AHF or decompensated heart failure symptoms, to monitoring the patient's response to therapeutic intervention,<sup>17</sup> early facilitated discharge,<sup>134</sup> and prompt community follow-up.<sup>135</sup> As key members of the multidisciplinary team, they are responsible for the 24 h monitoring of patients admitted with AHF ensuring effective communication among professionals as well as education and support to patients and caregivers. Over recent years, evidence has shown the value of nurse-led interventions, in terms of improved survival and quality of life as well as reduced length of hospital stay and rates of readmission.<sup>17,136-138</sup> Reviewing patients during the critical post-discharge phase (7–14 days) enables evaluation of clinical status and patient's self-management behaviours.

## **Potential future monitoring applications**

### ***Non-invasive haemodynamic monitoring***

The complications associated with invasive monitoring (e.g. mechanical complications and infectious risks) have contributed to the development of non-invasive technologies. Non-invasive techniques are undergoing considerable development, although none can be currently recommended for routine clinical use. Invasive techniques such as the PAC and transpulmonary thermodilution remain the reference standard.

### ***Contour of the pulse wave***

Several algorithms have been proposed to determine cardiac output based on determination of systolic area by analysis of the contour of the pulse wave.<sup>139</sup> In most cases, these signals are obtained from an arterial line. The system may be calibrated (by transpulmonary thermodilution or lithium dilution) or not. Some of the non-calibrated systems estimate arterial elastance and vascular tone.<sup>140</sup>

### ***Digital photoplethysmography***

Digital photoplethysmography is a technique for continuous measurement of blood volume changes.<sup>141,142</sup> Severe skin vasoconstriction, which is common in cardiogenic shock, impairs signal quality of blood pressure and is an important limitation of the technique. These monitors are not useful when arterial impedance is variable, such as with vasoconstrictor administration, unless given continuously and steady state has been reached. Digital photoplethysmography techniques do not appear to be sufficiently effective in assessing cardiac output in resuscitation patients with microcirculatory disorders, peripheral vasoconstriction, or high blood pressure lability.<sup>143,144</sup>

### ***Thoracic bioimpedance***

Transthoracic electric bioimpedance, also called impedance plethysmography or impedance cardiography, is a non-invasive method for measuring cardiac output.<sup>145</sup> Several haemodynamic parameters can be measured and calculated using the technique including flow (e.g. stroke volume/stroke index), resistance (e.g. systemic vascular resistance/index), contractility (e.g. cardiac power index, systolic time ratio, pre-ejection period, left ventricular ejection time, velocity index, acceleration index), and fluid (e.g. thoracic fluid content).<sup>146-148</sup>

Thoracic bioimpedance data may be informative in several clinical contexts, such as fluid management in patients with AHF,<sup>146,149</sup> and the differentiation of cardiogenic from pulmonary causes of acute dyspnoea, among others.<sup>150</sup> Bioimpedance might be useful for trend analysis, but the data should be interpreted cautiously, as the method is associated with limitations that may affect its accuracy (e.g. diseases of the aortic valve or aorta, arrhythmias, acute changes in tissue water such as pulmonary or chest wall oedema or pleural effusion, mechanical ventilation, body motion, and factors that affect conductivity between the electrodes and the skin like temperature and humidity).<sup>147,151-153</sup>

Bioreactance has a higher signal-to-noise ratio<sup>145,148</sup> and initial validation studies revealed promising results in different clinical settings.<sup>154</sup> Currently available data are insufficient to support guideline recommendations for use of bioimpedance and bioreactance in the routine monitoring of patients with AHF, although new systems show potential promise.

### ***Videomicroscopy***

Microvascular perfusion can be directly assessed by videomicroscopic techniques, mostly applied sublingually, although these are employed more often as research tools and are not widely applicable in routine clinical practice. Microcirculatory perfusion is altered in patients with AHF and cardiogenic shock.<sup>155</sup> Alterations in microvascular perfusion are more severe in non-survivors.<sup>155,156</sup> More importantly, changes over time also differ between survivors and non-survivors. Sublingual perfused capillary density may be an effective tool to measure tissue microvascular perfusion and estimate prognosis.<sup>156</sup> Measurements of veno-arterial pCO<sub>2</sub> differences may be an interesting surrogate.<sup>39</sup>

### **Conclusion**

Many tools are available for the in-hospital monitoring of patients with AHF, and each may play a role at various points throughout the patient's treatment course (*Table 5, Figure 1*). At present, clinical judgment guides the application of many of these tools. Physicians must set goals for monitored variables and tailor management according to their comprehensive interpretation of acquired clinical, biochemical, haemodynamic, and physiologic patient data. Future research initiatives should aim to identify the optimal in-hospital monitoring strategies for patients with AHF that reduce morbidity, prolong survival and are safe and cost-effective.

**Table 5.** Use of monitoring across points of care delivery

	Emergency department	Intensive care unit/coronary care unit	General ward
Patient population	All patients	Cardiogenic shock; pulmonary oedema	Congestive heart failure/‘wet and warm’
Non-invasive vital sign monitoring	At initial presentation and regularly during ED stay (blood pressure, heart rate, respiratory rate, pulse oximetry)	See haemodynamic monitoring (continuous blood pressure, respiratory rate monitoring)	Ideally three times daily, but at a minimum once daily (blood pressure, respiratory rate, pulse oximetry)
Body weight	On admission if possible/available	Daily when possible (requires specific bed with a scale)	Daily
Urine output/fluid balance	Check response to first dose of diuretic in the ED	Daily monitoring of net fluid balance	When indicated; urinary catheter not recommended
Blood gases and lactate	See ICU/coronary care unit recommendation	On admission to ICU and every 1–2 h during acute phase, then decrease frequency	As indicated by clinical situation and co-morbidities
Electrolytes and markers of renal function	All patients	Daily or more often if clinically indicated	Every 1–2 days until discharge indicated
Complete blood count	All patients	As indicated by clinical situation and co-morbidities	As indicated by clinical situation and co-morbidities
Liver function	All patients	As indicated after decongestion achieved	As indicated by clinical situation and co-morbidities
Cardiac biomarkers	NP and cTn ST2 may be considered	cTn only to evaluate for suspected AMI	NP may be considered pre-discharge
Procalcitonin	On admission Procalcitonin may be useful in patients with suspected coexisting infection	As indicated by clinical situation and co-morbidities	As indicated by clinical situation and co-morbidities
ECG and telemetry monitoring	ECG on admission, telemetry in selected cases	All	Continuous telemetry in selected cases
Chest X-ray	On admission	Only as needed	Only as needed
Echocardiography	Immediate echocardiography in patients with haemodynamic instability (cardiogenic shock) and in patients suspected of acute life-threatening structural or functional cardiac abnormalities	Upon ICU admission (unless direct admission from ED) and repeated as needed	Early (preferably within 48 h of admission) in <i>de novo</i> patients and those with unknown cardiac function. Repeated echocardiogram not needed unless relevant deterioration (or lack of improvement) in clinical status
Lung ultrasound	Lung ultrasound if available	As indicated by clinical situation and co-morbidities; assessment of decongestion	As indicated by clinical situation and co-morbidities; assessment of decongestion
Haemodynamic monitoring	Non-invasive measurements of blood pressure	Invasive blood pressure; CVC; cardiac output monitoring (various devices); PAC in selected patients in cardiogenic shock	N/A

AMI, acute myocardial infarction; cTn, cardiac troponin; CVC, central venous catheter; ECG, electrocardiogram; ED, emergency department; ICU, intensive care unit; N/A, not applicable; NP, natriuretic peptide; PAC, pulmonary artery catheter.

## Acknowledgements

The authors acknowledge the Heart Failure Association of the European Society of Cardiology for organizing the meeting during which this topic was discussed.

The authors acknowledge Wendy Gattis Stough, PharmD (Campbell University College of Pharmacy and Health Sciences), supported by the Heart Failure Association of the European Society of Cardiology for contributions to content development, writing, and editing the manuscript.

The work of Jelena Čelutkienė was supported by the grant from Lithuanian Research Council Nr. MIP-049/2015.

**Conflict of interest:** V.P.H.: research grants from Abbott Laboratories and Roche Diagnostics; personal fees from Roche Diagnostics. J.P.: personal fees from Servier, Novartis, Orion Pharma and Roche Diagnostics. H.P.B.L.R.: unrestricted research grants from Roche Diagnostics, Novartis Pharma, GSK Pharma. J.C.: research grant from Lithuanian Research Council; personal fees (lecture and investigator fees) from Novartis, Servier, Grindex, Orivas, and Amgen. O.C.: research grant from Servier, Vifor, Roche, Novartis; Steering Committee for Novartis. S.P.C.: research grants from Novartis, National Institutes of Health, PCORI, Cardiorentis; personal fees from Novartis, Roche, Trevena, Cardiorentis. D.d.B.: personal fees from Edwards Lifesciences and Fresenius Kabi. E.G.: research grants from Retia Medical, Deltex Medical, Sphingotec; personal fees from Edwards Lifescience, Magnisense, Sphingotec, Roche Diagnostics; patent pending WO 2015173785 A1. J.L.: personal fees from Bayer, Boehringer Ingelheim, Novartis, Pfizer, Roche Diagnostics, Orion Pharma, Servier, Vifor Pharma. J.M.: travel grants for congresses from Philips, Boehringer Ingelheim, and Menarini; personal fees from Novartis; consultant fees from Thermo Fisher. A.M.: research grant from Roche; personal fees (speaker honoraria) from Abbott, Novartis, Orion, Roche, Servier; personal fees (advisory board/steering committee) from Cardiorentis, Adrenomed, MyCartis, ZS Pharma, Critical Diagnostics. O.M.: competitive research grants from the Instituto de Salud Carlos III supported with funds from the Spanish Ministry of Health and FEDER (PI11/01021, PI15/01019), Fundació La Marató de TV3 (2015/2510), and from the Catalonia Govern for Consolidated Groups of Investigation (GRC 2009/1385 and 2014/0313); personal fees/honorarium for expert consultancy from Novartis. C.M.: research grants from the Swiss National Science Foundation, the Swiss Heart Foundation, the Swiss Commission for Technology and Innovation, the European Union, the Stiftung für kardiovaskuläre Forschung Basel, the University of Basel, Abbott, Beckman Coulter, Biomerieux, Brahms, Indorsia, Roche, Sanofi, Siemens, Singulex, Sphingotec, as well as speaker honoraria/consulting honoraria from Abbott, Alere, Amgen, Astra Zeneca, Biomerieux, Boehringer Ingelheim, BMS, Brahms, Cardiorentis, Indorsia, Novartis, Roche, Sanofi, Siemens, and Singulex. A.R.: personal fees for lectures and/or advisory board meetings from Orphan Pharma GmbH, Amomed Pharma GmbH, Orion Pharma AG, Novartis Pharma AG. F.R.: research grants from St Jude Medical, Novartis; personal fees for lectures from St Jude Medical, Servier, Zoll, Novartis, Bayer, and Abbott; personal fees for advisory board or steering committee clinical trial meetings from AstraZeneca, Sanofi, Cardiorentis, Amgen, BMS, Pfizer, Fresenius, Vifor, and Roche. J.M.W.: personal fees from Novartis. R.d.B.: research grants to employer (UMC Groningen) from AstraZeneca, Bristol-Myers Squibb, Trevena, Novartis; consulting fees paid to employer (UMC Groningen) from Novartis, Roche, Thermofisher; personal fees (consulting) from Novartis, Servier. M.G.C.L.: research grants from CIBERCV, (FEDER Funds); Novartis and Pfizer; personal fees from Novartis, Abbott, Astellas, and MSD. The other authors report no conflict of interests.

## References

1. Crespo-Leiro MG, Anker SD, Maggioni AP, Coats AJ, Filippatos G, Ruschitzka F, Ferrari R, Piepoli MF, Delgado Jimenez JF, Metra M, Fonseca C, Hradec J, Amir O, Logeart D, Dahlstrom U, Merkely B, Drozd J, Goncalvesova E, Hassanein M, Chioncel O, Lainscak M, Seferovic PM, Tousoulis D, Kavoliuniene A, Fruhwald F, Fazlibegovic E, Temizhan A, Gatzov P, Erglis A, Laroche C, Mebazaa A; Heart Failure Association (HFA) of the European Society of Cardiology (ESC). European Society of Cardiology Heart Failure Long-Term Registry (ESC-HF-LT): 1-year follow-up outcomes and differences across regions. *Eur J Heart Fail* 2016;18:613–625.
2. Christ M, Stork S, Dorr M, Heppner HJ, Muller C, Wachter R, Riemer U; Trend HF Germany Project. Heart failure epidemiology 2000–2013: insights from the German Federal Health Monitoring System. *Eur J Heart Fail* 2016;18:1009–1018.
3. Koudstaal S, Pujades-Rodriguez M, Denaxas S, Gho JM, Shah AD, Yu N, Patel RS, Gale CP, Hoes AW, Cleland JG, Asselbergs FW, Hemingway H. Prognostic burden of heart failure recorded in primary care, acute hospital admissions, or both: a population-based linked electronic health record cohort study in 2.1 million people. *Eur J Heart Fail* 2017;19:1119–1127.
4. Omersa D, Farkas J, Erzen I, Lainscak M. National trends in heart failure hospitalization rates in Slovenia 2004–2012. *Eur J Heart Fail* 2016;18:1321–1328.
5. Schmidt M, Ulrichsen SP, Pedersen L, Botker HE, Sorensen HT. Thirty-year trends in heart failure hospitalization and mortality rates and the prognostic impact of co-morbidity: a Danish nationwide cohort study. *Eur J Heart Fail* 2016;18:490–499.
6. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, Gonzalez-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;18:891–975.
7. Van Aelst LN, Arrigo M, Placido R, Akiyama E, Girerd N, Zannad F, Manivet P, Rossignol P, Badoz M, Sadoune M, Launay JM, Gayat E, Lam CS, Cohen-Solal A, Mebazaa A, Seronde MF. Acutely decompensated heart failure with preserved and reduced ejection fraction present with comparable haemodynamic congestion. *Eur J Heart Fail* 2018;20:738–747.
8. Nieminen MS, Brutsaert D, Dickstein K, Drexler H, Follath F, Harjola VP, Hochadel M, Komajda M, Lassus J, Lopez-Sendon JL, Ponikowski P, Tavazzi L; EuroHeart Survey Investigators; Heart Failure Association, European Society of Cardiology. EuroHeart Failure Survey II (EHFS II): a survey on hospitalized acute heart failure patients: description of population. *Eur Heart J* 2006;27:2725–2736.
9. Rudiger A, Harjola VP, Muller A, Mattila E, Saila P, Nieminen M, Follath F. Acute heart failure: clinical presentation, one-year mortality and prognostic factors. *Eur J Heart Fail* 2005;7:662–670.
10. Breidhardt T, Moreno-Weidmann Z, Uthoff H, Sabti Z, Appeli S, Puelacher C, Stallone F, Twerenbold R, Wildi K, Kozuharov N, Wussler D, Flores D, Shrestha S, Badertscher P, Boeddinghaus J, Nestelberger T, Gimenez MR, Staub D, Aschwanden M, Lohrmann J, Pfister O, Osswald S, Mueller C. How accurate is clinical assessment of neck veins in the estimation of central venous pressure in acute heart failure? Insights from a prospective study. *Eur J Heart Fail* 2018;20:1160–1162.
11. Damman K, Testani JM. The kidney in heart failure: an update. *Eur Heart J* 2015;36:1437–1444.
12. Moller S, Bernardi M. Interactions of the heart and the liver. *Eur Heart J* 2013;34:2804–2811.
13. Harjola VP, Mullens W, Banaszewski M, Bauersachs J, Brunner-La Rocca HP, Chioncel O, Collins SP, Doehner W, Filippatos GS, Flammer AJ, Fuhrmann V, Lainscak M, Lassus J, Legrand M, Masip J, Mueller C, Papp Z, Parissis J, Platz E, Rudiger A, Ruschitzka F, Schafer A, Seferovic PM, Skouri H, Yilmaz MB, Mebazaa A. Organ dysfunction, injury and failure in acute heart failure: from pathophysiology to diagnosis and management. A review on behalf of the Acute Heart Failure Committee of the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur J Heart Fail* 2017;19:821–836.
14. Vincent JL, De Backer D. Circulatory shock. *N Engl J Med* 2013;369:1726–1734.
15. Chioncel O, Mebazaa A, Harjola VP, Coats AJ, Piepoli MF, Crespo-Leiro MG, Laroche C, Seferovic PM, Anker SD, Ferrari R, Ruschitzka F, Lopez-Fernandez S, Miani D, Filippatos G,

- Maggioni AP; ESC Heart Failure Long-Term Registry Investigators. Clinical phenotypes and outcome of patients hospitalized for acute heart failure: the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail* 2017;19:1242–1254.
16. Gheorghiade M, Follath F, Ponikowski P, Barsuk JH, Blair JE, Cleland JG, Dickstein K, Drazner MH, Fonarow GC, Jaarsma T, Jondeau G, Sendon JL, Mebazaa A, Metra M, Nieminen M, Pang PS, Seferovic P, Stevenson LW, van Veldhuisen DJ, Zannad F, Anker SD, Rhodes A, McMurray JJ, Filippatos G. Assessing and grading congestion in acute heart failure: a scientific statement from the Acute Heart Failure Committee of the Heart Failure Association of the European Society of Cardiology and endorsed by the European Society of Intensive Care Medicine. *Eur J Heart Fail* 2010;12:423–433.
  17. Mebazaa A, Yilmaz MB, Levy P, Ponikowski P, Peacock WF, Laribi S, Ristic AD, Lambrinou E, Masip J, Riley JP, McDonagh T, Mueller C, deFilippi C, Harjola VP, Thiele H, Piepoli MF, Metra M, Maggioni A, McMurray J, Dickstein K, Damman K, Seferovic PM, Ruschitzka F, Leite-Moreira AF, Bellou A, Anker SD, Filippatos G. Recommendations on pre-hospital & early hospital management of acute heart failure: a consensus paper from the Heart Failure Association of the European Society of Cardiology, the European Society of Emergency Medicine and the Society of Academic Emergency Medicine. *Eur J Heart Fail* 2015;17:544–558.
  18. Nowak RM, Reed BP, DiSomma S, Nanayakkara P, Moyer M, Millis S, Levy P. Presenting phenotypes of acute heart failure patients in the ED: identification and implications. *Am J Emerg Med* 2017;35:536–542.
  19. Chaudhry SI, Wang Y, Concato J, Gill TM, Krumholz HM. Patterns of weight change preceding hospitalization for heart failure. *Circulation* 2007;116:1549–1554.
  20. Gottlieb SS, Stebbins A, Voors AA, Hasselblad V, Ezekowitz JA, Califf RM, O'Connor CM, Starling RC, Hernandez AF. Effects of nesiritide and predictors of urine output in acute decompensated heart failure: results from ASCEND-HF (Acute Study of Clinical Effectiveness of Nesiritide and Decompensated Heart Failure). *J Am Coll Cardiol* 2013;62:1177–1183.
  21. Schiff GD, Fung S, Speroff T, McNutt RA. Decompensated heart failure: symptoms, patterns of onset, and contributing factors. *Am J Med* 2003;114:625–630.
  22. Mentz RJ, Kjeldsen K, Rossi GP, Voors AA, Cleland JG, Anker SD, Gheorghiade M, Fiuzat M, Rossignol P, Zannad F, Pitt B, O'Connor C, Felker GM. Decongestion in acute heart failure. *Eur J Heart Fail* 2014;16:471–482.
  23. Zile MR, Bennett TD, St John SM, Cho YK, Adamson PB, Aaron MF, Aranda JM Jr, Abraham WT, Smart FW, Stevenson LW, Kueffer FJ, Bourge RC. Transition from chronic compensated to acute decompensated heart failure: pathophysiological insights obtained from continuous monitoring of intracardiac pressures. *Circulation* 2008;118:1433–1441.
  24. Ambrosy AP, Cerbin LP, Armstrong PW, Butler J, Coles A, DeVore AD, Dunlap ME, Ezekowitz JA, Felker GM, Fudim M, Greene SJ, Hernandez AF, O'Connor CM, Schulte P, Starling RC, Teerlink JR, Voors AA, Mentz RJ. Body weight change during and after hospitalization for acute heart failure: patient characteristics, markers of congestion, and outcomes: findings from the ASCEND-HF trial. *JACC Heart Fail* 2017;5:1–13.
  25. Galen BT. Underpad weight to estimate urine output in adult patients with urinary incontinence. *J Geriatr Cardiol* 2015;12:189–190.
  26. Chartier-Kastler E, Ballanger P, Petit J, Fourmarier M, Bart S, Ragni-Ghazarossian E, Ruffion A, Le Normand L, Costa P. Randomized, crossover study evaluating patient preference and the impact on quality of life of urisheaths vs absorbent products in incontinent men. *BJU Int* 2011;108:241–247.
  27. Singh D, Shrestha K, Testani JM, Verbrugge FH, Dupont M, Mullens W, Tang WH. Insufficient natriuretic response to continuous intravenous furosemide is associated with poor long-term outcomes in acute decompensated heart failure. *J Card Fail* 2014;20:392–399.
  28. Testani JM, Brisco MA, Turner JM, Spatz ES, Bellumkonda L, Parikh CR, Tang WH. Loop diuretic efficiency: a metric of diuretic responsiveness with prognostic importance in acute decompensated heart failure. *Circ Heart Fail* 2014;7:261–270.
  29. Verbrugge FH, Nijst P, Dupont M, Penders J, Tang WH, Mullens W. Urinary composition during decongestive treatment in heart failure with reduced ejection fraction. *Circ Heart Fail* 2014;7:766–772.
  30. Breidthardt T, Weidmann ZM, Twerenbold R, Gantenbein C, Stallone F, Rentsch K, Rubini Gimenez M, Kozhuharov N, Sabti Z, Breitenbacher D, Wildi K, Puelacher C, Honegger U, Wagener M, Schumacher C, Hillinger P, Osswald S, Mueller C. Impact of haemoconcentration during acute heart failure therapy on mortality and its relationship with worsening renal function. *Eur J Heart Fail* 2017;19:226–236.

31. ter Maaten JM, Dunning AM, Valente MA, Damman K, Ezekowitz JA, Califf RM, Starling RC, van der Meer P, O'Connor CM, Schulte PJ, Testani JM, Hernandez AF, Tang WH, Voors AA. Diuretic response in acute heart failure – an analysis from ASCEND-HF. *Am Heart J* 2015;170:313–321.
32. Valente MA, Voors AA, Damman K, van Veldhuisen DJ, Massie BM, O'Connor CM, Metra M, Ponikowski P, Teerlink JR, Cotter G, Davison B, Cleland JG, Givertz MM, Bloomfield DM, Fiuzat M, Dittrich HC, Hillege HL. Diuretic response in acute heart failure: clinical characteristics and prognostic significance. *Eur Heart J* 2014;35:1284–1293.
33. Voors AA, Davison BA, Teerlink JR, Felker GM, Cotter G, Filippatos G, Greenberg BH, Pang PS, Levin B, Hua TA, Severin T, Ponikowski P, Metra M; RELAX-AHF Investigators. Diuretic response in patients with acute decompensated heart failure: characteristics and clinical outcome – an analysis from RELAX-AHF. *Eur J Heart Fail* 2014;16:1230–1240.
34. Park JJ, Choi DJ, Yoon CH, Oh IY, Lee JH, Ahn S, Yoo BS, Kang SM, Kim JJ, Baek SH, Cho MC, Jeon ES, Chae SC, Ryu KH, Oh BH; KorHF Registry. The prognostic value of arterial blood gas analysis in high-risk acute heart failure patients: an analysis of the Korean Heart Failure (KorHF) registry. *Eur J Heart Fail* 2015;17:601–611.
35. Liborio AB, Noritomi DT, Leite TT, de Melo Bezerra CT, de Faria ER, Kellum JA. Increased serum bicarbonate in critically ill patients: a retrospective analysis. *Intensive Care Med* 2015;41:479–486.
36. Minana G, Nunez J, Banuls P, Sanchis J, Nunez E, Robles R, Mascarell B, Palau P, Chorro FJ, Llacer A. Prognostic implications of arterial blood gases in acute decompensated heart failure. *Eur J Intern Med* 2011;22:489–494.
37. Masip J, De Mendoza D, Planas K, Paez J, Sanchez B, Cancio B. Peripheral venous blood gases and pulse-oximetry in acute cardiogenic pulmonary oedema. *Eur Heart J Acute Cardiovasc Care* 2012;1:275–280.
38. Walkey AJ, Farber HW, O'Donnell C, Cabral H, Eagan JS, Philippides GJ. The accuracy of the central venous blood gas for acid–base monitoring. *J Intensive Care Med* 2010;25:104–110.
39. Ospina-Tascon GA, Umana M, Bermudez WF, Bautista-Rincon DF, Valencia JD, Madrinan HJ, Hernandez G, Bruhn A, Arango-Davila C, De Backer D. Can venous-to-arterial carbon dioxide differences reflect microcirculatory alterations in patients with septic shock? *Intensive Care Med* 2016;42:211–221.
40. Ospina-Tascon GA, Umana M, Bermudez W, Bautista-Rincon DF, Hernandez G, Bruhn A, Granados M, Salazar B, Arango-Davila C, De Backer D. Combination of arterial lactate levels and venous–arterial CO<sub>2</sub> to arterial–venous O<sub>2</sub> content difference ratio as markers of resuscitation in patients with septic shock. *Intensive Care Med* 2015;41:796–805.
41. Rimachi R, Bruzzi de Carvahlo F, Orellano-Jimenez C, Cotton F, Vincent JL, De Backer D. Lactate/pyruvate ratio as a marker of tissue hypoxia in circulatory and septic shock. *Anaesth Intensive Care* 2012;40:427–432.
42. Nichol AD, Egi M, Pettila V, Bellomo R, French C, Hart G, Davies A, Stachowski E, Reade MC, Bailey M, Cooper DJ. Relative hyperlactatemia and hospital mortality in critically ill patients: a retrospective multi-centre study. *Crit Care* 2010;14:R25.
43. Haas SA, Lange T, Saugel B, Petzoldt M, Fuhrmann V, Metschke M, Kluge S. Severe hyperlactatemia, lactate clearance and mortality in unselected critically ill patients. *Intensive Care Med* 2016;42:202–210.
44. Vincent JL, Quintairos ES, Couto L Jr, Taccone FS. The value of blood lactate kinetics in critically ill patients: a systematic review. *Crit Care* 2016;20:257.
45. Klein L, O'Connor CM, Leimberger JD, Gattis-Stough W, Pina IL, Felker GM, Adams KF Jr, Califf RM, Gheorghiuade M; OPTIME-CHF Investigators. Lower serum sodium is associated with increased short-term mortality in hospitalized patients with worsening heart failure: results from the Outcomes of a Prospective Trial of Intravenous Milrinone for Exacerbations of Chronic Heart Failure (OPTIME-CHF) study. *Circulation* 2005;111:2454–2460.
46. Rusinaru D, Tribouilloy C, Berry C, Richards AM, Whalley GA, Earle N, Poppe KK, Guazzi M, Macin SM, Komajda M, Doughty RN; MAGGIC Investigators. Relationship of serum sodium concentration to mortality in a wide spectrum of heart failure patients with preserved and with reduced ejection fraction: an individual patient data meta-analysis: Meta-Analysis Global Group in Chronic heart failure (MAGGIC). *Eur J Heart Fail* 2012;14:1139–1146.
47. Verbrugge FH, Steels P, Grieten L, Nijst P, Tang WH, Mullens W. Hyponatremia in acute decompensated heart failure: depletion versus dilution. *J Am Coll Cardiol* 2015;65:480–492.
48. Arrigo M, Bettex D, Rudiger A. Management of atrial fibrillation in critically ill patients. *Crit Care Res Pract* 2014;2014:840615.

49. Grodin JL, Simon J, Hachamovitch R, Wu Y, Jackson G, Halkar M, Starling RC, Testani JM, Tang WH. Prognostic role of serum chloride levels in acute decompensated heart failure. *J Am Coll Cardiol* 2015;66:659–666.
50. Delanaye P, Cavalier E, Pottel H. Serum creatinine: not so simple! *Nephron* 2017;136:302–308.
51. Levey AS, Inker LA. Assessment of glomerular filtration rate in health and disease: a state of the art review. *Clin Pharmacol Ther* 2017;102:405–419.
52. Gottlieb SS, Abraham W, Butler J, Forman DE, Loh E, Massie BM, O'Connor CM, Rich MW, Stevenson LW, Young J, Krumholz HM. The prognostic importance of different definitions of worsening renal function in congestive heart failure. *J Card Fail* 2002;8:136–141.
53. Mullens W, Abrahams Z, Francis GS, Sokos G, Taylor DO, Starling RC, Young JB, Tang WH. Importance of venous congestion for worsening of renal function in advanced decompensated heart failure. *J Am Coll Cardiol* 2009;53:589–596.
54. Damman K, Valente MA, Voors AA, O'Connor CM, van Veldhuisen DJ, Hillege HL. Renal impairment, worsening renal function, and outcome in patients with heart failure: an updated meta-analysis. *Eur Heart J* 2014;35:455–469.
55. Metra M, Davison B, Bettari L, Sun H, Edwards C, Lazzarini V, Piovanelli B, Carubelli V, Bugatti S, Lombardi C, Cotter G, Dei Cas L. Is worsening renal function an ominous prognostic sign in patients with acute heart failure? The role of congestion and its interaction with renal function. *Circ Heart Fail* 2012;5:54–62.
56. Testani JM, Chen J, McCauley BD, Kimmel SE, Shannon RP. Potential effects of aggressive decongestion during the treatment of decompensated heart failure on renal function and survival. *Circulation* 2010;122:265–272.
57. Mullens W, Verbrugge FH, Nijst P, Tang WH. Renal sodium avidity in heart failure: from pathophysiology to treatment strategies. *Eur Heart J* 2017;38:1872–1882.
58. Nikolaou M, Parissis J, Yilmaz MB, Seronde MF, Kivikko M, Laribi S, Paugam-Burtz C, Cai D, Pohjanjousi P, Laterre PF, Deye N, Poder P, Cohen-Solal A, Mebazaa A. Liver function abnormalities, clinical profile, and outcome in acute decompensated heart failure. *Eur Heart J* 2013;34:742–749.
59. Ambrosy AP, Vaduganathan M, Huffman MD, Khan S, Kwasny MJ, Fought AJ, Maggioni AP, Swedberg K, Konstam MA, Zannad F, Gheorghiadu M; EVEREST Trial Investigators. Clinical course and predictive value of liver function tests in patients hospitalized for worsening heart failure with reduced ejection fraction: an analysis of the EVEREST trial. *Eur J Heart Fail* 2012;14:302–311.
60. Chow SL, Maisel AS, Anand I, Bozkurt B, de Boer RA, Felker GM, Fonarow GC, Greenberg B, Januzzi JL Jr, Kiernan MS, Liu PP, Wang TJ, Yancy CW, Zile MR. Role of biomarkers for the prevention, assessment, and management of heart failure: a scientific statement from the American Heart Association. *Circulation* 2017;135:e1054–e1091.
61. Brunner-La Rocca HP, Sanders-van Wijk S, Knackstedt C. Biomarkers in patients with acute dyspnoea: what for? *Eur Heart J* 2012;33:2124–2126.
62. Kociol RD, Horton JR, Fonarow GC, Reyes EM, Shaw LK, O'Connor CM, Felker GM, Hernandez AF. Admission, discharge, or change in B-type natriuretic peptide and long-term outcomes: data from Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure (OPTIMIZE-HF) linked to Medicare claims. *Circ Heart Fail* 2011;4:628–636.
63. Cheng V, Kazanagra R, Garcia A, Lenert L, Krishnaswamy P, Gardetto N, Clopton P, Maisel A. A rapid bedside test for B-type peptide predicts treatment outcomes in patients admitted for decompensated heart failure: a pilot study. *J Am Coll Cardiol* 2001;37:386–391.
64. Dhaliwal AS, Deswal A, Pritchett A, Aguilar D, Kar B, Soucek J, Bozkurt B. Reduction in BNP levels with treatment of decompensated heart failure and future clinical events. *J Card Fail* 2009;15:293–299.
65. Bettencourt P, Azevedo A, Pimenta J, Frieiro F, Ferreira S, Ferreira A. N-terminal-pro-brain natriuretic peptide predicts outcome after hospital discharge in heart failure patients. *Circulation* 2004;110:2168–2174.
66. Felker G, Anstrom KJ, Adams KF, Ezekowitz J, Fiuzat M, Houston-Miller N, Januzzi J, Mark DB, Pina I, Passmore G, Whellan DJ, Yang H, Cooper LS, Leifer ES, Desvigne-Nickens P, O'Connor CM. Effect of natriuretic peptide-guided therapy on hospitalization or cardiovascular mortality in high-risk patients with heart failure and reduced ejection fraction: a randomized clinical trial. *JAMA* 2017;318:713–720.
67. Mebazaa A, Nieminen MS, Packer M, Cohen-Solal A, Kleber FX, Pocock SJ, Thakkar R, Padley RJ, Poder P, Kivikko M; SURVIVE Investigators. Levosimendan vs dobutamine for



- patients with acute decompensated heart failure: the SURVIVE randomized trial. *JAMA* 2007;297:1883–1891.
68. McCullough PA. How trialists and pharmaceutical sponsors have failed us by thinking that acute heart failure is a 48-hour illness. *Am J Cardiol* 2017;120:505–508.
  69. Packer M, O'Connor C, McMurray J, Wittes J, Abraham W, Anker S, Dickstein K, Filippatos G, Holcomb R, Krum H, Maggioni A, Mebazaa A, Peacock F, Petrie MC, Ponikowski P, Ruschitzka F, Van Veldhuisen D, Kowarski L, Schactman M, Holzmeister J; TRUE-AHF Investigators. Effect of ularitide on cardiovascular mortality in acute heart failure. *N Engl J Med* 2017;376:1956–1964.
  70. Cohen-Solal A, Logeart D, Huang B, Cai D, Nieminen MS, Mebazaa A. Lowered B-type natriuretic peptide in response to levosimendan or dobutamine treatment is associated with improved survival in patients with severe acutely decompensated heart failure. *J Am Coll Cardiol* 2009;53:2343–2348.
  71. Arrigo M, Gayat E, Parenica J, Ishihara S, Zhang J, Choi DJ, Park JJ, Alhabib KF, Sato N, Miro O, Maggioni AP, Zhang Y, Spinar J, Cohen-Solal A, Iwashyna TJ, Mebazaa A; GREAT Network. Precipitating factors and 90-day outcome of acute heart failure: a report from the intercontinental GREAT registry. *Eur J Heart Fail* 2017;19:201–208.
  72. Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S, Baumgartner H, Gaemperli O, Achenbach S, Agewall S, Badimon L, Baigent C, Bueno H, Bugiardini R, Carerj S, Casselman F, Cuisset T, Erol C, Fitzsimons D, Halle M, Hamm C, Hildick-Smith D, Huber K, Iliodromitis E, James S, Lewis BS, Lip GY, Piepoli MF, Richter D, Rosemann T, Sechtem U, Steg PG, Vrints C, Luis ZJ. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016;37:267–315.
  73. Januzzi JL Jr, Felker GM. Surfing the biomarker tsunami at JACC: heart failure. *JACC Heart Fail* 2013;1:213–215.
  74. Lassus J, Gayat E, Mueller C, Peacock WF, Spinar J, Harjola VP, van Kimmenade R, Pathak A, Mueller T, Disomma S, Metra M, Pascual-Figal D, Laribi S, Logeart D, Noura S, Sato N, Potocki M, Parenica J, Collet C, Cohen-Solal A, Januzzi JL Jr, Mebazaa A; GREAT Network. Incremental value of biomarkers to clinical variables for mortality prediction in acutely decompensated heart failure: the Multinational Observational Cohort on Acute Heart Failure (MOCA) study. *Int J Cardiol* 2013;168:2186–2194.
  75. Herrero-Puente P, Prieto-Garcia B, Garcia-Garcia M, Jacob J, Martin-Sanchez FJ, Pascual-Figal D, Bueno H, Gil V, Llorens P, Vazquez-Alvarez J, Romero-Pareja R, Sanchez-Gonzalez M, Miro O. Predictive capacity of a multimarker strategy to determine short-term mortality in patients attending a hospital emergency Department for acute heart failure. BIO-EAHFE study. *Clin Chim Acta* 2017;466:22–30.
  76. Maisel A, Neath SX, Landsberg J, Mueller C, Nowak RM, Peacock WF, Ponikowski P, Mockel M, Hogan C, Wu AH, Richards M, Clopton P, Filippatos GS, Di Somma S, Anand I, Ng LL, Daniels LB, Christenson RH, Potocki M, McCord J, Terracciano G, Hartmann O, Bergmann A, Morgenthaler NG, Anker SD. Use of procalcitonin for the diagnosis of pneumonia in patients presenting with a chief complaint of dyspnoea: results from the BACH (Biomarkers in Acute Heart Failure) trial. *Eur J Heart Fail* 2012;14:278–286.
  77. Arrigo M, Tolppanen H, Sadoune M, Feliot E, Teixeira A, Laribi S, Plaisance P, Noura S, Yilmaz MB, Gayat E, Mebazaa A. Effect of precipitating factors of acute heart failure on readmission and long-term mortality. *ESC Heart Fail* 2016;3:115–121.
  78. Drew BJ, Califf RM, Funk M, Kaufman ES, Krucoff MW, Laks MM, Macfarlane PW, Sommargren C, Swiryn S, Van Hare GF. Practice standards for electrocardiographic monitoring in hospital settings: an American Heart Association scientific statement from the Councils on Cardiovascular Nursing, Clinical Cardiology, and Cardiovascular Disease in the Young: endorsed by the International Society of Computerized Electrocardiology and the American Association of Critical-Care Nurses. *Circulation* 2004;110:2721–2746.
  79. Snider A, Papaleo M, Beldner S, Park C, Katechis D, Galinkin D, Fein A. Is telemetry monitoring necessary in low-risk suspected acute chest pain syndromes? *Chest* 2002;122:517–523.
  80. Saleem MA, McClung JA, Aronow WS, Kannam H. Inpatient telemetry does not need to be used in the management of older patients hospitalized with chest pain at low risk for in-hospital coronary events and mortality. *J Gerontol A Biol Sci Med Sci* 2005;60:605–606.

81. Hollander JE, Valentine SM, McCuskey CF, Brogan GX Jr. Are monitored telemetry beds necessary for patients with nontraumatic chest pain and normal or nonspecific electrocardiograms? *Am J Cardiol* 1997;79:1110–1111.
82. Collins S, Storrow AB, Kirk JD, Pang PS, Diercks DB, Gheorghide M. Beyond pulmonary edema: diagnostic, risk stratification, and treatment challenges of acute heart failure management in the emergency department. *Ann Emerg Med* 2008;51:45–57.
83. Wang CS, FitzGerald JM, Schulzer M, Mak E, Ayas NT. Does this dyspneic patient in the emergency department have congestive heart failure? *JAMA* 2005;294:1944–1956.
84. Collins SP, Lindsell CJ, Storrow AB, Abraham WT. Prevalence of negative chest radiography results in the emergency department patient with decompensated heart failure. *Ann Emerg Med* 2006;47:13–18.
85. Studler U, Kretzschmar M, Christ M, Breidthardt T, Noveanu M, Schoetzau A, Perruchoud AP, Steinbrich W, Mueller C. Accuracy of chest radiographs in the emergency diagnosis of heart failure. *Eur Radiol* 2008;18:1644–1652.
86. Chakko S, Woska D, Martinez H, de Marchena E, Futterman L, Kessler KM, Myerberg RJ. Clinical, radiographic, and hemodynamic correlations in chronic congestive heart failure: conflicting results may lead to inappropriate care. *Am J Med* 1991;90:353–359.
87. Collins S, Storrow AB, Albert NM, Butler J, Ezekowitz J, Felker GM, Fermann GJ, Fonarow GC, Givertz MM, Hiestand B, Hollander JE, Lanfear DE, Levy PD, Pang PS, Peacock WF, Sawyer DB, Teerlink JR, Lenihan DJ. Early management of patients with acute heart failure: state of the art and future directions. A consensus document from the Society for Academic Emergency Medicine/Heart Failure Society of America Acute Heart Failure Working Group. *J Card Fail* 2015;21:27–43.
88. Dash H, Lipton MJ, Chatterjee K, Parmley WW. Estimation of pulmonary artery wedge pressure from chest radiograph in patients with chronic congestive cardiomyopathy and ischaemic cardiomyopathy. *Br Heart J* 1980;44:322–329.
89. Volpicelli G, ElBarbary M, Blaivas M, Lichtenstein DA, Mathis G, Kirkpatrick AW, Melniker L, Gargani L, Noble VE, Via G, Dean A, Tsung JW, Soldati G, Copetti R, Bouhemad B, Reissig A, Agricola E, Rouby JJ, Arbelot C, Liteplo A, Sargsyan A, Silva F, Hoppmann R, Breikreutz R, Seibel A, Neri L, Storti E, Petrovic T. International evidence-based recommendations for point-of-care lung ultrasound. *Intensive Care Med* 2012;38:577–591.
90. Singh V. Acute respiratory distress syndrome. *J Lung Pulm Respir Res* 2016;3:00078.
91. Sheard S, Rao P, Devaraj A. Imaging of acute respiratory distress syndrome. *Respir Care* 2012;57:607–612.
92. Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, Camporota L, Slutsky AS; ARDS Definition Task Force. Acute respiratory distress syndrome: the Berlin definition. *JAMA* 2012;307:2526–2533.
93. Pesenti A, Musch G, Lichtenstein D, Mojoli F, Amato MB, Cinnella G, Gattinoni L, Quintel M. Imaging in acute respiratory distress syndrome. *Intensive Care Med* 2016;42:686–698.
94. Dokainish H, Zoghbi WA, Lakkis NM, Al-Bakshy F, Dhir M, Quinones MA, Nagueh SF. Optimal noninvasive assessment of left ventricular filling pressures: a comparison of tissue Doppler echocardiography and B-type natriuretic peptide in patients with pulmonary artery catheters. *Circulation* 2004;109:2432–2439.
95. Nagueh SF, Bhatt R, Vivo RP, Krim SR, Sarvari SI, Russell K, Edvardsen T, Smiseth OA, Estep JD. Echocardiographic evaluation of hemodynamics in patients with decompensated systolic heart failure. *Circ Cardiovasc Imaging* 2011;4:220–227.
96. Ommen SR, Nishimura RA, Appleton CP, Miller FA, Oh JK, Redfield MM, Tajik AJ. Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation* 2000;102:1788–1794.
97. Ritzema JL, Richards AM, Crozier IG, Frampton CF, Melton IC, Doughty RN, Stewart JT, Eigler N, Whiting J, Abraham WT, Troughton RW. Serial Doppler echocardiography and tissue Doppler imaging in the detection of elevated directly measured left atrial pressure in ambulant subjects with chronic heart failure. *JACC Cardiovasc Imaging* 2011;4:927–934.
98. Mullens W, Borowski AG, Curtin RJ, Thomas JD, Tang WH. Tissue Doppler imaging in the estimation of intracardiac filling pressure in decompensated patients with advanced systolic heart failure. *Circulation* 2009;119:62–70.
99. Nagueh SF, Smiseth OA, Appleton CP, Byrd BF 3rd, Dokainish H, Edvardsen T, Flachskampf FA, Gillebert TC, Klein AL, Lancellotti P, Marino P, Oh JK, Popescu BA, Waggoner AD. Recommendations for the evaluation of left ventricular diastolic function by echocardiography: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2016;29:277–314.

100. Lancellotti P, Galderisi M, Edvardsen T, Donal E, Goliash G, Cardim N, Magne J, Laginha S, Hagendorff A, Haland TF, Aaberge L, Martinez C, Rapacciuolo A, Santoro C, Ilardi F, Postolache A, Dulgheru R, Mateescu AD, Beladan CC, Deleanu D, Marchetta S, Auffret V, Schwammenthal E, Habib G, Popescu BA. Echo-Doppler estimation of left ventricular filling pressure: results of the multicentre EACVI Euro-Filling study. *Eur Heart J Cardiovasc Imaging* 2017;18:961–968.
101. Cameli M, Lisi M, Mondillo S, Padeletti M, Ballo P, Tsioulpas C, Bernazzali S, Maccherini M. Left atrial longitudinal strain by speckle tracking echocardiography correlates well with left ventricular filling pressures in patients with heart failure. *Cardiovasc Ultrasound* 2010;8:14.
102. Porter TR, Shillcutt SK, Adams MS, Desjardins G, Glas KE, Olson JJ, Troughton RW. Guidelines for the use of echocardiography as a monitor for therapeutic intervention in adults: a report from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2015;28:40–56.
103. Gandhi SK, Powers JC, Nomeir AM, Fowle K, Kitzman DW, Rankin KM, Little WC. The pathogenesis of acute pulmonary edema associated with hypertension. *N Engl J Med* 2001;344:17–22.
104. Kotecha D, Mohamed M, Shantsila E, Popescu BA, Steeds RP. Is echocardiography valid and reproducible in patients with atrial fibrillation? A systematic review. *Europace* 2017;19:1427–1438.
105. Logeart D, Saudubray C, Beyne P, Thabut G, Ennezat PV, Chavelas C, Zanker C, Bouvier E, Solal AC. Comparative value of Doppler echocardiography and B-type natriuretic peptide assay in the etiologic diagnosis of acute dyspnea. *J Am Coll Cardiol* 2002;40:1794–1800.
106. Ohman J, Harjola VP, Karjalainen P, Lassus J. Assessment of early treatment response by rapid cardiothoracic ultrasound in acute heart failure: cardiac filling pressures, pulmonary congestion and mortality. *Eur Heart J Acute Cardiovasc Care* 2017;2048872617708974.
107. Lamia B, Maizel J, Ochagavia A, Chemla D, Osman D, Richard C, Teboul JL. Echocardiographic diagnosis of pulmonary artery occlusion pressure elevation during weaning from mechanical ventilation. *Crit Care Med* 2009;37:1696–1701.
108. Caille V, Amiel JB, Charron C, Belliard G, Vieillard-Baron A, Vignon P. Echocardiography: a help in the weaning process. *Crit Care* 2010;14:R120.
109. Verbrugge FH, Grieten L, Mullens W. New insights into combinational drug therapy to manage congestion in heart failure. *Curr Heart Fail Rep* 2014;11:1–9.
110. Tsutsui RS, Borowski A, Tang WH, Thomas JD, Popovic ZB. Precision of echocardiographic estimates of right atrial pressure in patients with acute decompensated heart failure. *J Am Soc Echocardiogr* 2014;27:1072–1078.e2.
111. Ciozda W, Kedan I, Kehl DW, Zimmer R, Khandwalla R, Kimchi A. The efficacy of sonographic measurement of inferior vena cava diameter as an estimate of central venous pressure. *Cardiovasc Ultrasound* 2016;14:33.
112. Patel AR, Alsheikh-Ali AA, Mukherjee J, Evangelista A, Quraini D, Ordway LJ, Kuvin JT, Denofrio D, Pandian NG. 3D echocardiography to evaluate right atrial pressure in acutely decompensated heart failure *correlation with invasive hemodynamics*. *JACC Cardiovasc Imaging* 2011;4:938–945.
113. Nijst P, Martens P, Dupont M, Tang WH, Mullens W. Intrarenal flow alterations during transition from euvolemia to intravascular volume expansion in heart failure patients. *JACC Heart Fail* 2017;5:672–681.
114. Wetterslev M, Moller-Sorensen H, Johansen RR, Perner A. Systematic review of cardiac output measurements by echocardiography vs. thermodilution: the techniques are not interchangeable. *Intensive Care Med* 2016;42:1223–1233.
115. Mercado P, Maizel J, Beyls C, Titeca-Beauport D, Joris M, Kontar L, Riviere A, Bonef O, Soupison T, Tribouilloy C, de Cagny B, Slama M. Transthoracic echocardiography: an accurate and precise method for estimating cardiac output in the critically ill patient. *Crit Care* 2017;21:136.
116. Cecconi M, De Backer D, Antonelli M, Beale R, Bakker J, Hofer C, Jaeschke R, Mebazaa A, Pinsky MR, Teboul JL, Vincent JL, Rhodes A. Consensus on circulatory shock and hemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. *Intensive Care Med* 2014;40:1795–1815.
117. Vincent JL, Rhodes A, Perel A, Martin GS, Della Rocca G, Vallet B, Pinsky MR, Hofer CK, Teboul JL, de Boode WP, Scolletta S, Vieillard-Baron A, De Backer D, Walley KR, Maggiorini M, Singer M. Clinical review: Update on hemodynamic monitoring – a consensus of 16. *Crit Care* 2011;15:229.

118. Al Deeb M, Barbic S, Featherstone R, Dankoff J, Barbic D. Point-of-care ultrasonography for the diagnosis of acute cardiogenic pulmonary edema in patients presenting with acute dyspnea: a systematic review and meta-analysis. *Acad Emerg Med* 2014;21:843–852.
119. Liteplo AS, Marill KA, Villen T, Miller RM, Murray AF, Croft PE, Capp R, Noble VE. Emergency thoracic ultrasound in the differentiation of the etiology of shortness of breath (ETUDES): sonographic B-lines and N-terminal pro-brain-type natriuretic peptide in diagnosing congestive heart failure. *Acad Emerg Med* 2009;16:201–210.
120. Pivetta E, Goffi A, Lupia E, Tizzani M, Porrino G, Ferreri E, Volpicelli G, Balzaretto P, Banderali A, Iacobucci A, Locatelli S, Casoli G, Stone MB, Maule MM, Baldi I, Merletti F, Cibinel GA, Baron P, Battista S, Buonafede G, Busso V, Conterno A, Del Rizzo P, Ferrera P, Pecetto PF, Moiraghi C, Morello F, Steri F, Ciccone G, Calasso C, Caserta MA, Civita M, Condo C, D'Alessandro V, Del Colle S, Ferrero S, Griot G, Laurita E, Lazzerro A, Lo Curto F, Michelazzo M, Nicosia V, Palmari N, Ricchiardi A, Rolfo A, Rostagno R, Bar F, Boero E, Frascisco M, Micossi I, Mussa A, Stefanone V, Agricola R, Cordero G, Corradi F, Runzo C, Soragna A, Scullo D, Vercillo D, Allione A, Artana N, Corsini F, Dutto L, Lauria G, Morgillo T, Tartaglino B, Bergandi D, Cassetta I, Maserà C, Garrone M, Ghiselli G, Ausiello L, Barutta L, Bernardi E, Bono A, Forno D, Lamorte A, Lison D, Lorenzati B, Maggio E, Masi I, Maggiorotto M, Novelli G, Panero F, Perotto M, Ravazzoli M, Saglio E, Soardo F, Tizzani A, Tizzani P, Tullio M, Ulla M, Romagnoli E; SIMEU Group for Lung Ultrasound in the Emergency Department in Piedmont. Lung ultrasound-implemented diagnosis of acute decompensated heart failure in the ED: a SIMEU multicenter study. *Chest* 2015;148:202–210.
121. Noble VE, Murray AF, Capp R, Sylvia-Reardon MH, Steele DJR, Liteplo A. Ultrasound assessment for extravascular lung water in patients undergoing hemodialysis. Time course for resolution. *Chest* 2009;135:1433–1439.
122. Volpicelli G, Caramello V, Cardinale L, Mussa A, Bar F, Frascisco MF. Bedside ultrasound of the lung for the monitoring of acute decompensated heart failure. *Am J Emerg Med* 2008;26:585–591.
123. Platz E, Merz AA, Jhund PS, Vazir A, Campbell R, McMurray JJ. Dynamic changes and prognostic value of pulmonary congestion by lung ultrasound in acute and chronic heart failure: a systematic review. *Eur J Heart Fail* 2017;19:1154–1163.
124. Ohman J, Harjola VP, Karjalainen P, Lassus J. Focused echocardiography and lung ultrasound protocol for guiding treatment in acute heart failure. *ESC Heart Fail* 2018;5:120–128.
125. Marx G, Reinhart K. Venous oximetry. *Curr Opin Crit Care* 2006;12:263–268.
126. Harvey S, Harrison DA, Singer M, Ashcroft J, Jones CM, Elbourne D, Brampton W, Williams D, Young D, Rowan K; PAC-Man Study Collaboration. Assessment of the clinical effectiveness of pulmonary artery catheters in management of patients in intensive care (PAC-Man): a randomised controlled trial. *Lancet* 2005;366:472–477.
127. Binanay C, Califf RM, Hasselblad V, O'Connor CM, Shah MR, Sopko G, Stevenson LW, Francis GS, Leier CV, Miller LW; ESCAPE Investigators and ESCAPE Study Coordinators. Evaluation study of congestive heart failure and pulmonary artery catheterization effectiveness: the ESCAPE trial. *JAMA* 2005;294:1625–1633.
128. Peacock WF. When and for how long do patients with acute heart failure have to be admitted? *Emergencias* 2016;28:363–365.
129. Harjola P, Harjola VP. Can we do more for patients with acute heart failure before arrival at the hospital? *Emergencias* 2017;29:221–222.
130. Meijers WC, De Boer RA, Van Veldhuisen DJ, Jaarsma T, Hillege HL, Maisel AS, Di Somma S, Voors AA, Peacock WF. Biomarkers and low risk in heart failure. Data from COACH and TRIUMPH. *Eur J Heart Fail* 2015;17:1271–1282.
131. Miro O, Levy PD, Mockel M, Pang PS, Lambrinou E, Bueno H, Hollander JE, Harjola VP, Diercks DB, Gray AJ, DiSomma S, Papa AM, Collins SP. Disposition of emergency department patients diagnosed with acute heart failure: an international emergency medicine perspective. *Eur J Emerg Med* 2017;24:2–12.
132. Collins SP, Jenkins CA, Harrell FE Jr, Liu D, Miller KF, Lindsell CJ, Naftilan AJ, McPherson JA, Maron DJ, Sawyer DB, Weintraub NL, Fermann GJ, Roll SK, Sperling M, Storrow AB. Identification of emergency department patients with acute heart failure at low risk for 30-day adverse events: the STRATIFY Decision Tool. *JACC Heart Fail* 2015;3:737–747.
133. Miro O, Rossello X, Gil V, Martin-Sanchez FJ, Llorens P, Herrero-Puente P, Jacob J, Bueno H, Pocock S. Predicting 30-day mortality for patients with acute heart failure who are in the emergency department: a cohort study. *Ann Intern Med* 2017;167:698–705.

134. Lambrinou E, Kalogirou F, Lamnisis D, Sourtzi P. Effectiveness of heart failure management programmes with nurse-led discharge planning in reducing re-admissions: a systematic review and meta-analysis. *Int J Nurs Stud* 2012;49:610–624.
135. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA, McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel B, Sam F, Stevenson LW, Tang WH, Tsai EJ, Wilkoff BL. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;62:e147–e239.
136. Guder G, Stork S, Gelbrich G, Brenner S, Deubner N, Morbach C, Wallenborn J, Berliner D, Ertl G, Angermann CE. Nurse-coordinated collaborative disease management improves the quality of guideline-recommended heart failure therapy, patient-reported outcomes, and left ventricular remodelling. *Eur J Heart Fail* 2015;17:442–452.
137. Stromberg A, Martensson J, Fridlund B, Levin LA, Karlsson JE, Dahlstrom U. Nurse-led heart failure clinics improve survival and self-care behaviour in patients with heart failure: results from a prospective, randomised trial. *Eur Heart J* 2003;24:1014–1023.
138. Stamp KD, Machado MA, Allen NA. Transitional care programs improve outcomes for heart failure patients: an integrative review. *J Cardiovasc Nurs* 2014;29:140–154.
139. Caillard A, Gayat E, Tantom A, Dubreuil G, M'Bakulu E, Madadaki C, Bart F, Bresson D, Froelich S, Mebazaa A, Vallee F. Comparison of cardiac output measured by oesophageal Doppler ultrasonography or pulse pressure contour wave analysis. *Br J Anaesth* 2015;114:893–900.
140. Teboul JL, Saugel B, Cecconi M, De Backer D, Hofer CK, Monnet X, Perel A, Pinsky MR, Reuter DA, Rhodes A, Squara P, Vincent JL, Scheeren TW. Less invasive hemodynamic monitoring in critically ill patients. *Intensive Care Med* 2016;42:1350–1359.
141. Fortin J, Marte W, Grullenberger R, Hacker A, Habenbacher W, Heller A, Wagner C, Wach P, Skrabal F. Continuous non-invasive blood pressure monitoring using concentrically interlocking control loops. *Comput Biol Med* 2006;36:941–957.
142. Penaz J, Voigt A, Teichmann W. Contribution to the continuous indirect blood pressure measurement. *Z Gesamte Inn Med* 1976;31:1030–1033.
143. Monnet X, Picard F, Lidzboriski E, Mesnil M, Duranteau J, Richard C, Teboul JL. The estimation of cardiac output by the Nexfin device is of poor reliability for tracking the effects of a fluid challenge. *Crit Care* 2012;16:R212.
144. Bubenek-Turconi SI, Craciun M, Miclea I, Perel A. Noninvasive continuous cardiac output by the Nexfin before and after preload-modifying maneuvers: a comparison with intermittent thermodilution cardiac output. *Anesth Analg* 2013;117:366–372.
145. Jakovljevic DG, Trenell MI, MacGowan GA. Bioimpedance and bioreactance methods for monitoring cardiac output. *Best Pract Res Clin Anaesthesiol* 2014;28:381–394.
146. Yancy C, Abraham WT. Noninvasive hemodynamic monitoring in heart failure: utilization of impedance cardiography. *Congest Heart Fail* 2003;9:241–250.
147. Cotter G, Moshkovitz Y, Kaluski E, Cohen AJ, Miller H, Goor D, Vered Z. Accurate, noninvasive continuous monitoring of cardiac output by whole-body electrical bioimpedance. *Chest* 2004;125:1431–1440.
148. Jakovljevic DG, Moore S, Hallsworth K, Fattakhova G, Thoma C, Trenell MI. Comparison of cardiac output determined by bioimpedance and bioreactance methods at rest and during exercise. *J Clin Monit Comput* 2012;26:63–68.
149. Shochat M, Shotan A, Blondheim DS, Kazatsker M, Dahan I, Asif A, Shochat I, Rabinovich P, Rozenman Y, Meisel SR. Usefulness of lung impedance-guided pre-emptive therapy to prevent pulmonary edema during ST-elevation myocardial infarction and to improve long-term outcomes. *Am J Cardiol* 2012;110:190–196.
150. Stevanovic P, Scepanovic R, Radovanovic D, Bajec D, Perunovic R, Stojanovic D, Stevanovic D. Thoracic electrical bioimpedance theory and clinical possibilities in perioperative medicine. *Signa Vitae* 2008;3(Suppl 1):S22–S27.
151. Peyton PJ, Chong SW. Minimally invasive measurement of cardiac output during surgery and critical care: a meta-analysis of accuracy and precision. *Anesthesiology* 2010;113:1220–1235.
152. Raaijmakers E, Faes TJ, Scholten RJ, Goovaerts HG, Heethaar RM. A meta-analysis of three decades of validating thoracic impedance cardiography. *Crit Care Med* 1999;27:1203–1213.
153. Kamath SA, Drazner MH, Tasissa G, Rogers JG, Stevenson LW, Yancy CW. Correlation of impedance cardiography with invasive hemodynamic measurements in patients with advanced heart failure: the BioImpedance CardioGraphy (BIG) substudy of the Evaluation Study of

- Congestive Heart Failure and Pulmonary Artery Catheterization Effectiveness (ESCAPE) Trial. *Am Heart J* 2009;158:217–223.
154. Raval NY, Squara P, Cleman M, Yalamanchili K, Winklmaier M, Burkhoff D. Multicenter evaluation of noninvasive cardiac output measurement by bioimpedance technique. *J Clin Monit Comput* 2008;22:113–119.
  155. De Backer D, Creteur J, Dubois MJ, Sakr Y, Vincent JL. Microvascular alterations in patients with acute severe heart failure and cardiogenic shock. *Am Heart J* 2004;147:91–99.
  156. den Uil CA, Lagrand WK, van der Ent M, Jewbali LS, Cheng JM, Spronk PE, Simoons ML. Impaired microcirculation predicts poor outcome of patients with acute myocardial infarction complicated by cardiogenic shock. *Eur Heart J* 2010;31:3032–3039.
  157. Campbell P, Drazner MH, Kato M, Lakdawala N, Palardy M, Nohria A, Stevenson LW. Mismatch of right- and left-sided filling pressures in chronic heart failure. *J Card Fail* 2011;17:561–568.
  158. Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K, Solomon SD, Louie EK, Schiller NB. Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology, and the Canadian Society of Echocardiography. *J Am Soc Echocardiogr* 2010;23:685–713.
  159. Jue J, Chung W, Schiller NB. Does inferior vena cava size predict right atrial pressures in patients receiving mechanical ventilation? *J Am Soc Echocardiogr* 1992;5:613–619.
  160. Nagueh SF, Kopelen HA, Zoghbi WA. Relation of mean right atrial pressure to echocardiographic and Doppler parameters of right atrial and right ventricular function. *Circulation* 1996;93:1160–1169.