

## Extracellular vesicles as potential tools for regenerative therapy

Jose Antonio Rodriguez-Navarro , Jorge Pascual-Guerra , Silvia Sacristan ,  
Maria del Carmen Nogales , Juan Antonio Fafián-Labora & Ana O’Loughlen

To cite this article: Jose Antonio Rodriguez-Navarro , Jorge Pascual-Guerra , Silvia Sacristan ,  
Maria del Carmen Nogales , Juan Antonio Fafián-Labora & Ana O’Loughlen (2020): Extracellular  
vesicles as potential tools for regenerative therapy, Molecular & Cellular Oncology, DOI:  
[10.1080/23723556.2020.1809958](https://doi.org/10.1080/23723556.2020.1809958)

To link to this article: <https://doi.org/10.1080/23723556.2020.1809958>



© 2020 The Author(s). Published with  
license by Taylor & Francis Group, LLC.



Published online: 08 Sep 2020.



Submit your article to this journal [↗](#)



Article views: 75



View related articles [↗](#)



View Crossmark data [↗](#)

## Extracellular vesicles as potential tools for regenerative therapy

Jose Antonio Rodriguez-Navarro <sup>a</sup>, Jorge Pascual-Guerra <sup>a</sup>, Silvia Sacristan <sup>a</sup>, Maria del Carmen Nogales<sup>a</sup>, Juan Antonio Fafián-Labora<sup>b\*</sup>, and Ana O'Loghlen <sup>b</sup>

<sup>a</sup>Instituto Ramón y Cajal de Investigaciones Sanitarias, IRYCIS Neurobiología-Investigación, Madrid, Spain; <sup>b</sup>Epigenetics & Cellular Senescence Group, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK

### ABSTRACT

Small extracellular vesicles released by fibroblasts from young human donors diminish lipid peroxidation in senescent cells and in different old mice organs due to their enrichment in Glutathione-S-transferase Mu lipid antioxidant activity.

### ARTICLE HISTORY

Received 22 July 2020  
Revised 7 August 2020  
Accepted 10 August 2020

### KEYWORDS

Small extracellular vesicles; glutathione-S-transferase M; lipid peroxidation; aging; senescence

Aging is characterized by a progressive decline in the physiological function efficiency and by an increased susceptibility to disease. This decline is in part mediated by the activation of the cellular phenotype named senescence. Cellular senescence is characterized by a cell-cycle arrest, and by the senescence-associated secretory phenotype, or SASP, that includes the secretion of small extracellular vesicles (sEVs), soluble factors, and matrix remodeling enzymes.<sup>1</sup>

We recently showed that sEVs secreted by cells isolated from young human healthy individuals (sEV-Y) can ameliorate the senescence phenotype in old cells and in old mice by reducing lipo-oxidative stress.<sup>2</sup>



In vitro, different cell types (fibroblasts from old human donors, fibroblasts obtained from Hutchison-Gilford progeria patients, and young fibroblasts induced to senesce by over-expression of *H-RAS* oncogene) were used as old cells to determine the ability of sEV-Y to counteract senescence. Treatment of old cells with sEV-Y (but not with extracellular vesicles from young individuals (EV-Y) larger than 200 nm) reduced the levels of senescence markers and revert the proliferation arrest of fibroblasts. One of the most widely used biomarkers for senescent cells is the  $\beta$ -Galactosidase staining. We could observe that sEV-Y treatment in old mice reduced the levels of  $\beta$ -Galactosidase staining in tissue sections of different organs like liver, kidney, lung, and brown adipose tissue. Similarly, we found a decreased mRNA expression of SASP factors in the liver and kidney when old mice were treated with sEV-Y.<sup>2,3</sup>

Previous work from the O'Loghlen lab characterized the differential proteomic composition of proliferating versus senescent cells sEVs allowed us to identify enzymes of the glutathione antioxidant system enriched in sEV-Y.<sup>3</sup> In particular, Glutathione-S-Transferase Mu 2 (encoded by the

*GSTM2* gene) enzyme counteracts lipid oxidation and it decreases with age in mice and humans. The oxidation of lipids generates highly reactive products such as 4-hydroxynonenal (4-HNE) and malondialdehyde (MDA) reactive aldehydes that accumulate in senescence, aging, and age-related diseases. We observed a decrease in their levels in the liver and serum of aged mice after the treatment with sEV-Y enriched in antioxidant enzymes.<sup>2</sup> Accordingly, Yoshida et al. described that extracellular nicotinamide phosphoribosyltransferase enriched in EVs enhances Nicotinamide adenine dinucleotide (NAD<sup>+</sup>) biosynthesis once the vesicles are internalized by cells, improving motor activity and extending lifespan in old mice.<sup>4</sup> Therefore, additional components within sEV-Ys can also contribute to the observed reversion of certain characteristics of senescence and aging.

More experimental data are needed to affirm that sEV-Y have the ability to rejuvenate young tissues and counteract age-related pathologies as happens with cellular reprogramming.<sup>5</sup> We need longer-term experiments, more behavioral and physiological data. In spite of the mounting evidence supporting a protective and antioxidant role of sEVs in old individuals, we must be cautious before starting to use sEVs in clinical trials to counteract aging. EVs are implicated in the development of neurodegenerative and cardiovascular diseases, and participate in the progress of carcinomas. The sEV composition (ribonucleic acids, proteins, lipids, etc.) depends not only on the age of the donor cell, but also on the cell type, physiological condition, isolation methodology, etc. Cancer, and different stem and immune cells secrete sEVs that can influence physiological and pathological processes.<sup>6-8</sup>

However, in our experiments, sEV-Ys are helping aged cells and tissues to repair. Therefore, it will be interesting to perform long-term experiments, try different administration routes,

**CONTACT** Juan Antonio Fafián-Labora  [j.a.f.labora@qmul.ac.uk](mailto:j.a.f.labora@qmul.ac.uk)  Epigenetics & Cellular Senescence Group, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London E1 2AT, UK

\*Present address: Departamento de Fisioterapia, Medicina y Ciencias Biomédicas. Facultad de Ciencias de la Salud. Universidade da Coruña. INIBIC-Complejo Hospitalario Universitario A Coruña (CHUAC). Agrupación estratégica CICA-INIBIC. As Xubias, 15006-A Coruña, Spain

© 2020 Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

and test sEV-Y therapeutic capacity in pathological models to demonstrate their repairing and rejuvenating abilities.

## Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

## Funding

This work was supported by the Barts and The London School of Medicine and Dentistry [MGU0497]; Biotechnology and Biological Sciences Research Council [BB/P000223/1]; Instituto de Salud Carlos III [CP19-0010]; Ministerio de Economía, Industria y Competitividad, Gobierno de España [SAF2016-78666R]; Xunta de Galicia [ED481B 2017/117].

## ORCID

Jose Antonio Rodriguez-Navarro  <http://orcid.org/0000-0002-0741-6338>

Jorge Pascual-Guerra  <http://orcid.org/0000-0001-6667-3310>

Silvia Sacristan  <http://orcid.org/0000-0001-5294-7608>

Ana O'Loughlen  <http://orcid.org/0000-0002-8141-1965>

## References

1. Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013;153:1194–1217. doi:10.1016/j.cell.2013.05.039.
2. Fafian-Labora JA, Rodriguez-Navarro JA, O'Loughlen A. Small extracellular vesicles have GST activity and ameliorate senescence-related tissue damage. *Cell Metabol*. 2020;32(71–86):e75. doi:10.1016/j.cmet.2020.06.004.
3. Borghesan M, Fafián-Labora J, Eleftheriadou O, Carpintero-Fernández P, Paez-Ribes M, Vizcay-Barrena G, Swisa A, Kolodkin-Gal D, Ximénez-Embún P, Lowe R, et al. Small extracellular vesicles are key regulators of non-cell autonomous intercellular communication in senescence via the interferon protein IFITM3. *Cell Rep*. 2019;27(3956–3971):e3956. doi:10.1016/j.celrep.2019.05.095.
4. Yoshida M, Satoh A, Lin JB, Mills KF, Sasaki Y, Rensing N, Wong M, Apte RS, Imai S-I. Extracellular vesicle-contained eNAMPT delays aging and extends lifespan in mice. *Cell Metabol*. 2019;30(329–342):e325. doi:10.1016/j.cmet.2019.05.015.
5. Ocampo A, Reddy P, Martinez-Redondo P, Platero-Luengo A, Hatanaka F, Hishida T, Li M, Lam D, Kurita M, Beyret E, et al. In vivo amelioration of age-associated hallmarks by partial reprogramming. *Cell*. 2016;167(7):e1712. doi:10.1016/j.cell.2016.11.052.
6. O'Loughlen A, Marechal N, Pechalrieu D, Cura V, Franchini D-M, Faux C, Alby F, Troffer-Charlier N, Kudithipudi S, Jeltsch A. Role for extracellular vesicles in the tumour microenvironment. *Philos Trans R Soc Lond B Biol Sci*. 2018;373. doi:10.1098/rstb.2017.0072.
7. Fafian-Labora JA, O'Loughlen A. Classical and nonclassical intercellular communication in senescence and ageing. *Trends Cell Biol*. 2020;30(8):628–639. doi:10.1016/j.tcb.2020.05.003.
8. Fafian-Labora J, Carpintero-Fernández P, Jordan SJD, Shikh-Bahaei T, Abdullah SM, Mahenthiran M, Rodríguez-Navarro JA, Niklison-Chirou MV, O'Loughlen A. FASN activity is important for the initial stages of the induction of senescence. *Cell Death Dis*. 2019;10:318. doi:10.1038/s41419-019-1550-0.