

Supplementary S1. Levels of evidence of studies used in the prospective surveillance model evaluated with the Oxford Centre for Evidence-Based Medicine Level of Evidence Tool [15].

References	LEVEL OF EVIDENCE
Ali NA, O'Brien JM Jr, Hoffmann SP, et al. Acquired weakness, handgrip strength, and mortality in critically ill patients. <i>Am J Respir Crit Care Med.</i> 2008;178(3):261-268. doi:10.1164/rccm.200712-1829OC	3
Akyuz G, Kenis O. Physiotherapy modalities and rehabilitation techniques in the management of neuropathic pain. <i>Am J Phys Med Rehabil.</i> 2014;93(3):253-259. doi:10.1097/PHM.0000000000000037	5
Arribas-Romano A, Fernández-Carnero J, Molina-Rueda F, Angulo-Díaz-Parreño S, Navarro-Santana MJ. Efficacy of Physiotherapy on Nociceptive Pain Processing Alterations in Patients with Chronic Musculoskeletal Pain: A Systematic Review and Meta-analysis [published online ahead of print, 2020 Feb 25]. <i>Pain Med.</i> 2020;pnz366. doi:10.1093/pm/pnz366	1
Australian Department of Health. Australia's Physical Activity and Sedentary Behaviour Guidelines and the Australian 24-Hour Movement Guidelines 2019 for Adults (18-64 years). Available from: https://www1.health.gov.au/internet/main/publishing.nsf/Content/health-pubhlth-strateg-phys-act-guidelines#npa1864 . Accessed on 1st May 2020	1
Barreiro E, Bustamante V, Cejudo P, et al. Guidelines for the evaluation and treatment of muscle dysfunction in patients with chronic obstructive pulmonary disease. <i>Arch Bronconeumol.</i> 2015;51:384-395. doi:10.1016/j.arbres.2015.04.011	5
Bennett M. The LANSS Pain Scale: the Leeds assessment of neuropathic symptoms and signs. <i>Pain.</i> 2001;92:147-157. doi:10.1016/s0304-3959(00)00482-6	3
Bienvenu OJ, Williams JB, Yang A, Hopkins RO, Needham DM. Posttraumatic stress disorder in survivors of acute lung injury: evaluating the Impact of Event Scale-Revised. <i>Chest.</i> 2013;144(1):24-31. doi:10.1378/chest.12-0908	3
Boonstra AM, Stewart RE, Köke AJ, et al. Cut-Off Points for Mild, Moderate, and Severe Pain on the Numeric Rating Scale for Pain in Patients with Chronic	3

Musculoskeletal Pain: Variability and Influence of Sex and Catastrophizing. <i>Front Psychol.</i> 2016;7:1466. Published 2016 Sep 30. doi:10.3389/fpsyg.2016.01466	
Cesari M, Kritchevsky SB, Penninx BW, et al. Prognostic value of usual gait speed in well-functioning older people--results from the Health, Aging and Body Composition Study. <i>J Am Geriatr Soc.</i> 2005;53(10):1675-1680. doi:10.1111/j.1532-5415.2005.53501.x	3
Chen YC, Chen KC, Lu LH, Wu YL, Lai TJ, Wang CH. Validating the 6-minute walk test as an indicator of recovery in patients undergoing cardiac surgery: A prospective cohort study. <i>Medicine (Baltimore).</i> 2018;97(42):e12925. doi:10.1097/MD.00000000000012925	3
Dettling-Ihnenfeldt DS. The Post-Intensive Care Syndrome (PICS): Impact of ICU-stay on functioning and implications for rehabilitation care. 2017	4
Ekelund U, Steene-Johannessen J, Brown WJ, et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women [published correction appears in <i>Lancet.</i> 2016 Sep 24;388(10051):e6]. <i>Lancet.</i> 2016;388(10051):1302-1310. doi:10.1016/S0140-6736(16)30370-1	1
Elliott D, Denehy L, Berney S, Alison JA. Assessing physical function and activity for survivors of a critical illness: a review of instruments. <i>Aust Crit Care.</i> 2011;24(3):155-166. doi:10.1016/j.aucc.2011.05.002	5
Finnerup NB, Haroutounian S, Kamerman P, et al. Neuropathic pain: an updated grading system for research and clinical practice. <i>Pain.</i> 2016;157(8):1599-1606. doi:10.1097/j.pain.0000000000000492	2
Frontera WR, Slovik DM, Dawson DM. <i>Exercise in Rehabilitation Medicine. 2nd ed;</i> 2006.	5
Holland AE, Spruit MA, Troosters T, et al. An official European Respiratory Society/American Thoracic Society technical standard: field walking tests in chronic respiratory disease. <i>Eur Respir J.</i> 2014;44(6):1428-1446. doi:10.1183/09031936.00150314	5
Ingle L, Cleland JG, Clark AL. The long-term prognostic significance of 6-minute walk test distance in patients with chronic heart failure. <i>Biomed Res Int.</i> 2014;2014:505969. doi:10.1155/2014/505969	3

Jutte JE, Needham DM, Pfoh ER, Bienvenu OJ. Psychometric evaluation of the Hospital Anxiety and Depression Scale 3 months after acute lung injury. <i>J Crit Care.</i> 2015;30(4):793-798. doi:10.1016/j.jcrc.2015.04.006	3
Laisné F, Lecomte C, Corbière M. Biopsychosocial predictors of prognosis in musculoskeletal disorders: a systematic review of the literature (corrected and republished) *. <i>Disabil Rehabil.</i> 2012;34(22):1912-1941. doi:10.3109/09638288.2012.729362	1
Liu K, Zhang W, Yang Y, Zhang J, Li Y, Chen Y. Respiratory rehabilitation in elderly patients with COVID-19: A randomized controlled study. <i>Complement Ther Clin Pract.</i> 2020;39:101166. doi:10.1016/j.ctcp.2020.101166	2
Malfliet A, Kregel J, Meeus M, et al. Applying contemporary neuroscience in exercise interventions for chronic spinal pain: treatment protocol. <i>Braz J Phys Ther.</i> 2017;21(5):378-387. doi:10.1016/j.bjpt.2017.06.019	2
Mahoney FI, Barthel DW. Functional Evaluation: The Barthel Index. <i>Md State Med J.</i> 1965;14:61-65.	2
Mayer TG, Neblett R, Cohen H, et al. The development and psychometric validation of the central sensitization inventory. <i>Pain Pract.</i> 2012;12(4):276-285. doi:10.1111/j.1533-2500.2011.00493.x	2
Mo X, Jian W, Su Z, et al. Abnormal pulmonary function in COVID-19 patients at time of hospital discharge [published online ahead of print, 2020 May 12]. <i>Eur Respir J.</i> 2020;2001217. doi:10.1183/13993003.01217-2020	5
Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment [published correction appears in <i>J Am Geriatr Soc.</i> 2019 Sep;67(9):1991]. <i>J Am Geriatr Soc.</i> 2005;53(4):695-699. doi:10.1111/j.1532-5415.2005.53221.x	2
Norkin CC, White DJ. <i>Measurement of Joint Motion : A Guide to Goniometry. 5th edition;</i> 2017.	5
Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. <i>Eur Respir J.</i> 2005;26(5):948-968. doi:10.1183/09031936.05.00035205	5
Qiu S, Cai X, Sun Z, et al. Heart Rate Recovery and Risk of Cardiovascular Events and All-Cause Mortality: A Meta-Analysis of Prospective Cohort Studies. <i>J Am Heart Assoc.</i> 2017;6(5):e005505. Published 2017 May 9. doi:10.1161/JAHA.117.005505	2

Ramirez-Velez R, Perez-Sousa MA, Venegas-Sanabria LC, et al. Normative Values for the Short Physical Performance Battery (SPPB) and Their Association With Anthropometric Variables in Older Colombian Adults. The SABE Study, 2015. <i>Front Med.</i> 2020;7:52. doi:10.3389/fmed.2020.00052	2
Hunter J, Rawlings-Anderson K. Respiratory assessment. <i>Nurs Stand.</i> 2008;22(41):41-43. doi:10.7748/ns2008.06.22.41.41.c6576	5
Rubio Castañeda FJ, Tomás Aznar C, Muro Baquero C. Validity, Reliability and Associated Factors of the International Physical Activity Questionnaire Adapted to Elderly (IPAQ-E)]. <i>Rev Esp Salud Publica.</i> 2017;91:e201701004. Published 2017 Jan 18.	2
Santos GM, Souza AC, Virtuoso JF, Tavares GM, Mazo GZ. Predictive values at risk of falling in physically active and no active elderly with Berg Balance Scale. <i>Rev Bras Fisioter.</i> 2011;15(2):95-101. doi:10.1590/s1413-35552011000200003	3
Sidiras G, Patsaki I, Karatzanos E, et al. Long term follow-up of quality of life and functional ability in patients with ICU acquired Weakness - A post hoc analysis. <i>J Crit Care.</i> 2019;53:223-230. doi:10.1016/j.jcrc.2019.06.022	3
Smith JM, Lee AC, Zeleznik H, et al. Home and Community-Based Physical Therapist Management of Adults With Post-Intensive Care Syndrome [published online ahead of print, 2020 Apr 13]. <i>Phys Ther.</i> 2020;pzaa059. doi:10.1093/ptj/pzaa059	5
Vasconcelos KS, Dias JM, Bastone Ade C, et al. Handgrip Strength Cutoff Points to Identify Mobility Limitation in Community-dwelling Older People and Associated Factors. <i>J Nutr Health Aging.</i> 2016;20(3):306-315. doi:10.1007/s12603-015-0584-y	4
World Health Organization. Waist circumference and waist-hip ratio: report of a WHO expert consultation. Geneva, 8-11 December 2008. Accessed at https://apps.who.int/iris/bitstream/handle/10665/44583/9789241501491_eng.pdf?ua=1 ON May 5, 2020.	1
Yang LL, Yang T. Pulmonary Rehabilitation for Patients with Coronavirus Disease 2019 (COVID-19) [published online ahead of print, 2020 May 14]. <i>Chronic Dis Transl Med.</i> 2020;10.1016/j.cdtm.2020.05.002. doi:10.1016/j.cdtm.2020.05.002	5
Zhao HM, Xie YX, Wang C. Recommendations for respiratory rehabilitation in adults with COVID-19 [published online ahead of print, 2020 Apr 9]. <i>Chin Med J (Engl).</i> 2020;10.1097/CM9.0000000000000848. doi:10.1097/CM9.0000000000000848	5