

	<b>n (%)</b>
Adriamycin (other Anthracyclines)	39 (5)
Postpartum	9 (1.2)
Familial dilated cardiomyopathy	90 (11.6)
Myocarditis	46 (5.9)
Alcoholic dilated cardiomyopathy	31 (4)
Viral dilated cardiomyopathy	6 (0.8)
Other dilated cardiomyopathy	100 (12.9)
Idiopathic restrictive cardiomyopathy	43 (5.5)
Amyloidosis	39 (5)
Endomyocardial fibrosis	12 (1.5)
Sarcoidosis	3 (0.4)
Restrictive cardiomyopathy secondary to radio/chemotherapy	4 (0.5)
Other restrictive cardiomyopathy	12 (1.5)
Hypertrophic cardiomyopathy	114 (14.7)
Valvular heart disease	195 (25.1)
Other cardiac disease	34 (4.4)
Rheumatic disease	1 (0.1)

**Supplementary table 1. Breakdown of other heart failure etiologies.**

	<b>Excluded N=1,164</b>	<b>Included N=2,810</b>	<b>p-value</b>
Age (years), mean $\pm$ SD	53,03 $\pm$ 11,41	52,84 $\pm$ 11,56	0,637
Male sex, n (%)	928 (79,7)	2203 (78,4)	0,342
Heart failure etiology			0,222
Idiopathic DCM, n (%)	316 (27,1)	794 (28,3)	
Ischemic DCM, n (%)	523 (44,9)	1167 (41,5)	
Congenital, n (%)	24 (2,1)	71 (2,5)	
Other, n (%)	301 (25,9)	779 (27,7)	
Recent infection, n (%)	107 (10,1)	371 (13,2)	0,009
Pre-transplant mechanical ventilation, n (%)	123 (11,3)	384 (13,7)	0,049
Mechanical circulatory support, n (%)	245 (22,2)	737 (26,2)	0,009
1-year mortality, n (%)	290 (24,9)	587 (20,9)	0,005

**Supplementary table 2. Differences in some baseline characteristics and 1-year mortality between excluded patients due to missing values of bilirubin and creatinine and the final study cohort. DCM, dilated cardiomyopathy; SD, standard deviation.**