

Site-Selective C–H Amination of Phenol-Containing Biomolecules

Published as part of the *Organic Letters virtual special issue* “Chemoselective Methods for Labeling and Modification of Peptides and Proteins”.

Carlota Girón-Elola, Ibon Sasiain, Rosalía Sánchez-Fernández, Elena Pazos, and Arkaitz Correa*



Cite This: *Org. Lett.* 2023, 25, 4383–4387



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: A C–N bond-forming cross-dehydrogenative coupling of a collection of Tyr-containing peptides and estrogens with heteroarenes is described. This oxidative coupling is distinguished by its scalability, operational simplicity, and air tolerance and enables the appendage of phenothiazines and phenoxazines in phenol-like compounds. When incorporated into a Tb(III) metalloprotein, the Tyr-phenothiazine moiety acts as a sensitizer for the Tb(III) ion, providing a new tool for the design of luminescent probes.




Owing to their chemical versatility, amino acids are privileged motifs in a number of disciplines such as organic synthesis, medicinal chemistry, and chemical biology. The strategical incorporation of noncanonical amino acids into a given peptide template represents a unique platform to create molecular diversity within the peptide drug discovery space.¹ In fact, peptides housing nonproteinogenic amino acids often exhibit improved permeability and higher stability to those of their parent native analogues. As a result, the past decade has witnessed an exponential growth of the peptide therapeutics market,² and the development of general methods to modify peptides in a late-stage fashion poses a challenging task of paramount importance.

Metal-catalyzed C–H functionalization reactions have recently evolved into powerful yet innovative technologies toward the site-selective modification of amino acids and peptides.³ In this respect, a wide range of bioconjugation methods are available today to tag highly reactive residues such as Cys, Lys, or Trp. Conversely, despite its prevalence in a myriad of medically relevant compounds, the modification of tyrosine (Tyr) has been comparatively overlooked.⁴ The introduction of directing groups (DGs) into the oxygen atom of the phenol ring has enabled olefination,⁵ hydroxylation,⁶ acylation,⁷ acetoxylation,⁸ and acyloxylation⁹ reactions at the *ortho*-C(sp²)-H bond. Although chelation assistance constitutes a common practice within the field and results in the assembly of unprecedented peptides, the cleavage of the required DG is often a low-yielding step, which deeply jeopardizes the practicality of the latter methods. Accordingly, the performance of bioconjugation reactions in native Tyr-containing peptides stands out as a streamlined and preferred avenue. In this respect, a number of elegant C-modification methods have been reported for the appendage of different

functional groups including Mannich-type reactions,¹⁰ nitrations,¹¹ and trifluoromethylations,¹² among others. Based on the attractive features of phenothiazines,¹³ of particular importance are the methods developed by Lei¹⁴ and MacMillan¹⁵ to label Tyr derivatives upon electrochemistry and photocatalysis, respectively (Scheme 1).

The click-like phenol–phenothiazine coupling reaction¹⁶ can occur in a plethora of reaction conditions;¹⁷ however, its application at more complex biomolecules has been less explored. Whereas the reported amination protocols are effective to append phenothiazine¹⁴ and phenoxazines¹⁵ in

Scheme 1. Amination of Tyr-Containing Peptides



Received: May 12, 2023


Published: June 7, 2023



Tyr-containing proteins, the requirement of sophisticated reaction devices may reduce their use in a mainstream context. Inspired by these elegant methods, we wondered whether we could perform those couplings in an operationally simple fashion without special reaction equipment. In this Letter, we unlock the prowess of silver carbonate to assist a reliable and scalable oxidative C–H amination process. This method enables the practical appendance of benzoxazines and benzothiazines into phenol-containing biomolecules, such as Tyr-containing peptides and estrogens.

We started our studies by selecting the cross-dehydrogenative coupling (CDC) of simple Boc-Tyr-OMe (**1a**) with phenothiazine as the model reaction. After some experimentation,¹⁸ we found that Ag₂CO₃ (30 mol %) in *o*-xylene under air at 90 °C provided **2aa** quantitatively (Table 1,

Table 1. C–H Phenothiazination of **1a**^a




entry	Change from standard conditions	2aa (%) ^b
1	none	99
2	Without Ag ₂ CO ₃	0
3	Under Ar	45
4	H ₂ O instead of <i>o</i> -xylene	11
5	AgOAc instead of Ag ₂ CO ₃	23
6	Ag ₂ O instead of Ag ₂ CO ₃	47
7	Ag ₂ CO ₃ (25 mol %)	76
8	At 70 °C	82
9	Boc-Tyr-OMe (1.2 equiv)	99 (89) ^c
10	Ag ₂ CO ₃ (1.0 equiv) in H ₂ O under Ar	83

^aReaction conditions: **1a** (0.50 mmol), phenothiazine (0.25 mmol), Ag₂CO₃ (30 mol %), *o*-xylene (1.0 mL) at 90 °C for 16 h under air. ^bYield of isolated product after column chromatography. ^cReaction performed with 15.06 mmol of phenothiazine.

entry 1). Control experiments underpinned the crucial role of silver carbonate¹⁹ and air within the reaction outcome (entries 2–3). The use of other Ag(I) salts or a lower amount of silver carbonate resulted in lower yields of **2aa** (entries 5–7). Importantly, the reaction could be performed in neat water,¹⁸ albeit a stoichiometric amount of Ag₂CO₃ and an argon atmosphere were required (entry 10). The latter evidenced the potential utility of the method in more complex Tyr-containing biomolecules, which usually require an aqueous system. Moreover, the synthesis of **2aa** could be performed with 3.0 g (15.06 mmol) of phenothiazine with an excellent 89% isolated yield (entry 9).¹⁸ These preliminary studies revealed that the challenging phenothiazination of a Tyr residue could be performed in a reliable practical fashion without a sophisticated setup. Notably, the parent benzoxazine as well as substituted benzothiazines could be also used as the coupling partners (Scheme 2). It is important to highlight the tolerance to alkynes and chlorides, which represent versatile reaction sites for creating molecular diversity. In accordance with previous studies in simple phenol systems,¹⁷ other heteroarenes such as carbazole or

Scheme 2. Scope of the Heteroarene^{a,b}




^aReaction conditions: **1a** (0.30 mmol), heteroarene (0.25 mmol), Ag₂CO₃ (30 mol %), *o*-xylene (1.0 mL) at 90 °C for 16 h under air. ^bYield of isolated product after column chromatography, average of at least two independent runs.

indoles were found unreactive under the standard conditions, thus highlighting the unique capacity of phenothiazine to form *N*-centered radicals.

With the optimized conditions in hand, we next investigated the preparative scope of the method to assemble a new family of decorated Tyr-containing peptides in a simple fashion (Scheme 3). A wide variety of dipeptides underwent the corresponding C–N bond-forming coupling in good to excellent yields. Remarkably, phenothiazine was efficiently installed at dipeptides housing Leu (**4a**), Phe (**4b**), Ala (**4c–d**), Ile (**4e**), Pro (**4f**), Ser (**4g**), Thr (**4h**), Met (**4i**), Asp (**4j**), Glu (**4k**), Lys (**4l**), Arg (**4m**), and Trp (**4n–o**). Of particular importance are compounds incorporating oxidizable functional groups such as hydroxyl, thioether, and indole, which remained intact along the process. It is noteworthy that the performance of the process in water in the presence of 1.0 equiv of Ag₂CO₃ ushered in higher yields for certain highly polar dipeptides such as those incorporating Pro (**4f**), Thr (**4h**), and Asp (**4j**), which may be due to solubility issues. Interestingly, more challenging tri- and tetrapeptides (**4p–x**) could be efficiently tagged with phenothiazine. Importantly, tetrapeptides **4w** and **4x** bearing the amino acid sequence of biologically relevant *Endomorphin-1* and *Endomorphin-2*, respectively, were also labeled in a late-stage fashion. Other Tyr derived from naturally occurring compounds or active pharmaceuticals including those derived from palmitic acid (**6a**), oleic acid (**6b**), ibuprofen (**6d**), and artificial sweetener neotame (**6g**) smoothly underwent the corresponding amination reaction (Scheme 4). Likewise, non-natural Tyr residues incorporating adamantane (**6c**), photo-switchable diazobenzene (**6e**), and carboxamide (**6f**) site-selectively underwent our amination manifold. In order to demonstrate the synthetic utility of the method, we further explored its use toward the late-stage modification of estrogens, such as estrone and estradiol. The coupling of estrone with phenothiazines preferentially occurred at the C2 position, although variable amounts of the parent isomer substituted at the C4 were also formed (Scheme 4).

In order to gain some insight into the reaction mechanism, some experiments were conducted. The presence of a free-hydroxyl group within the Tyr was found key as Tyr housing

Scheme 3. C–H Phenothiazination of Tyr-Containing Peptides^{a,b}


^aReaction conditions: 3 (0.30 mmol), phenothiazine (0.25 mmol), Ag₂CO₃ (30 mol %), *o*-xylene (1.0 mL) at 90 °C for 16 h under air. ^bYield of isolated product after column chromatography, average of at least two independent runs with no more than 5% variation in yield between runs. ^c3 (0.30 mmol), phenothiazine (0.25 mmol), Ag₂CO₃ (1.0 equiv), H₂O (1.0 mL) at 90 °C for 16 h under Ar.

OTs, OAc, or OMe groups remained intact (Table S4).¹⁸ The addition of BHT as a radical trap ushered in the entire inhibition of the process. Conversely, the addition of TEMPO resulted in slightly lower yields of 2aa (68% vs 99%), but the process was not entirely shut down (Table S3).¹⁸ This reactivity pattern has been observed in similar processes,^{16,17} and some authors have suggested that TEMPO may prolong the lifetime of the transient *N*-centered radical species upon the formation of covalent intermediates.^{17c,d,19} Accordingly,²⁰ we assumed that the process would start with the Ag-assisted formation of an electrophilic *N*-centered radical I at the phenothiazine (Scheme S1).¹⁸ The latter could likely be trapped by the phenol ring of the Tyr in a polarity-matched fashion to yield radical intermediate II that would eventually evolve into the target product through further oxidation to III and aromatization (Scheme S1, path a). However, a radical–radical coupling between the phenothiazine radical I and the phenoxy radical IV derived from the starting phenol has been often proposed and cannot be ruled out at this stage (path b).²⁰ The key role of air

could be attributed to the reoxidation of the *in situ* formed Ag(0) species.

Considering the potential use of the Tyr(NPT) moiety as a fluorophore^{14,15} and the attractive luminescent properties of lanthanide ions, i.e., long lifetimes in the order of milliseconds and narrow emission bands in the visible and near-infrared region,²¹ we wondered whether the Tyr(NPT) unit could be used as an antenna for lanthanide ions, and therefore synthesized peptide 7[Tb] (Figure 1A).^{18,22} As expected, compared to parent Tyr analogue 8, peptide 7 exhibited red-shifted fluorescence with an emission band centered at 446 nm (Figure 1B). Interestingly, the time-gated emission spectrum of the metallopeptide 7[Tb] showed the characteristic emission bands of the Tb(III) ion at 489, 544, 586, and 620 nm, for the corresponding transitions ⁵D₄ → ⁷F_{*J*}, *J* = 6, 5, 4, 3, confirming that the Tyr(NPT) chromophore is an adequate Tb(III) antenna.

In summary, we have developed a CDC of Tyr-containing peptides with phenothiazines and phenoxazines. Salient features of this method are the scalability, operational

Scheme 4. C–H Phenothiazination of Tyr-Containing Compounds and Estrogens^{a,b}

^aReaction conditions: **5** (0.30 mmol), phenothiazine (0.25 mmol), Ag₂CO₃ (30 mol %), *o*-xylene (1.0 mL) at 90 °C for 16 h under air. ^bYield of isolated product after column chromatography, average of at least two independent runs with no more than 5% variation in yield between runs. ^cYield of the minor C4-functionalized derivative.




Figure 1. (A) Peptide sequence of **7**[Tb]. (B) Fluorescence spectra of 2 μM **7** (blue line) and **8** (black line) in 10 mM HEPES, 100 mM NaCl, pH 7.5, excitation 254 and 274 nm, respectively. (C) Luminescence spectra of 2 μM **7**[Tb] in 10 mM HEPES, 100 mM NaCl, pH 7.5, excitation 254 nm.

simplicity, tolerance to air and water, and application for the late-stage modification of estrogens. In addition, we have shown that the Tyr(NPT) moiety can be used as an antenna for Tb(III) ions, providing a new tool for the design of luminescent probes for biologically relevant targets.

ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.3c01560>.

Experimental procedures, syntheses and characterization of all new compounds, and tables with details of several optimization studies (PDF)

Accession Codes

CCDC 2257500 and 2257502 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Arkaitz Correa – University of the Basque Country (UPV/EHU), Department of Organic Chemistry I, Joxe Mari Korta R&D Center, 20018 Donostia-San Sebastián, Spain; orcid.org/0000-0002-9004-3842; Email: arkaitz.correa@ehu.eus

Authors

Carlota Girón-Elola – University of the Basque Country (UPV/EHU), Department of Organic Chemistry I, Joxe Mari Korta R&D Center, 20018 Donostia-San Sebastián, Spain

Ibon Sasiain – University of the Basque Country (UPV/EHU), Department of Organic Chemistry I, Joxe Mari Korta R&D Center, 20018 Donostia-San Sebastián, Spain

Rosalía Sánchez-Fernández – CICA – Centro Interdisciplinar de Química e Bioloxía and Departamento de Química, Facultade de Ciencias, Universidade da Coruña, 15071 A Coruña, Spain

Elena Pazos – CICA – Centro Interdisciplinar de Química e Bioloxía and Departamento de Química, Facultade de Ciencias, Universidade da Coruña, 15071 A Coruña, Spain; orcid.org/0000-0001-9397-2703

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.orglett.3c01560>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful to Ministerio de Ciencia e Innovación (PID2021-122889NB-I00, CTQ2017-89166-R, MCIN/AEI/FEDER, UE), the Consellería de Cultura, Educación e Universidade, Xunta de Galicia (ED431C 2018/39, ED431C 2022/39, and 508/2020), and Basque Government (IT1741-22) for financial support. We thank for technical and human support provided by Central Service of Analysis-SGIker of UPV/EHU and European funding (ERDF and ESF). R.S.F. thanks the Consellería de Cultura, Educación e Universidade, Xunta de Galicia for her PhD fellowship (ED481A-2020/008). E.P. thanks the MCIN/AEI/10.13039/501100011033 and ESF Investing in your future for her RyC contract (RYC2019-027199-I).

REFERENCES

- (1) Ding, Y.; Ting, J. P.; Liu, J.; Al-Azzam, S.; Pandya, P.; Afshar, S. Impact of non-proteinogenic amino acids in the discovery and development of peptide therapeutics. *Amino Acids* **2020**, *52*, 1207.
- (2) (a) Li Petri, G.; Di Martino, S.; De Rosa, M. Peptidomimetics: An Overview of Recent Medicinal Chemistry Efforts toward the Discovery of Novel Small Molecule Inhibitors. *J. Med. Chem.* **2022**, *65*, 7438. (b) Muttenthaler, M.; King, G. F.; Adams, D. J.; Alewood, P. F. Trends in peptide drug discovery. *Nat. Rev. Drug Discovery* **2021**, *20*, 309.
- (3) (a) Kjærsgaard, N. L.; Nielsen, T. B.; Gothelf, K. V. Chemical Conjugation to Less Targeted Proteinogenic Amino Acids. *ChemBioChem.* **2022**, *23*, No. e202200245. (b) deGruyter, J. N.; Malins, L. R.; Baran, P. S. Residue-Specific Peptide Modification: A Chemist's Guide. *Biochemistry* **2017**, *56*, 3863.
- (4) (a) Correa, A. Metal-Catalyzed C(sp²)-H Functionalization Processes of Phenylalanine- and Tyrosine-Containing Peptides. *Eur. J. Inorg. Chem.* **2021**, *2021*, 2928. (b) Dorta, D. A.; Deniaud, D.; Mével, M.; Gouin, S. G. Tyrosine Conjugation Methods for Protein Labelling. *Chem.—Eur. J.* **2020**, *26*, 14257.
- (5) Hu, Q.-L.; Hou, K.-Q.; Li, J.; Ge, Y.; Song, Z.-D.; Chan, A. S. C.; Xiong, X.-F. Silanol: a bifunctional group for peptide synthesis and late-stage functionalization. *Chem. Sci.* **2020**, *11*, 6070.
- (6) Andrade-Sampedro, P.; Matxain, J. M.; Correa, A. Ru-Catalyzed C-H Hydroxylation of Tyrosine-Containing Di- and Tripeptides toward the Assembly of L-DOPA Derivatives. *Adv. Synth. Catal.* **2022**, *364*, 2072.
- (7) (a) Urruzuno, I.; Andrade-Sampedro, P.; Correa, A. Late-Stage C-H Acylation of Tyrosine-Containing Oligopeptides with Alcohols. *Org. Lett.* **2021**, *23*, 7279. (b) San Segundo, M.; Correa, A. Site-selective aqueous C-H acylation of tyrosine-containing oligopeptides with aldehydes. *Chem. Sci.* **2020**, *11*, 11531.
- (8) Urruzuno, I.; Andrade-Sampedro, P.; Correa, A. Pd-Catalyzed Site-Selective C-H Acetoxylation of Tyrosine-Containing Peptides. *Eur. J. Org. Chem.* **2023**, *26*, No. e20220148.
- (9) Hou, X.; Kaplaneris, N.; Yuan, B.; Frey, J.; Ohyama, T.; Messinis, A. M.; Ackermann, L. Ruthenaelectro-catalyzed C-H acyloxylation for late-stage tyrosine and oligopeptide diversification. *Chem. Sci.* **2022**, *13*, 3461.
- (10) Joshi, N. S.; Whitaker, L. R.; Francis, M. B. A Three-Component Mannich-Type Reaction for Selective Tyrosine Bioconjugation. *J. Am. Chem. Soc.* **2004**, *126*, 15942.
- (11) Long, T.; Liu, L.; Tao, Y.; Zhang, W.; Quan, J.; Zheng, J.; Hegemann, J. D.; Uesugi, M.; Yao, W.; Tian, H.; Wang, H. Light-Controlled Tyrosine Nitration of Proteins. *Angew. Chem., Int. Ed.* **2021**, *60*, 13414.
- (12) Ichiishi, N.; Caldwell, J. P.; Lin, M.; Zhong, W.; Zhu, X.; Streckfuss, E.; Kim, H.-Y.; Parish, C. A.; Krska, S. W. Protecting group free radical C-H trifluoromethylation of peptides. *Chem. Sci.* **2018**, *9*, 4168.
- (13) (a) Lu, X.; Zhan, Y.; He, W. Recent development of small-molecule fluorescent probes based on phenothiazine and its derivatives. *J. Photochem. Photobiol. B, Biol.* **2022**, *234*, 112528. (b) Ohlow, M. J.; Moosmann, B. Phenothiazine: the seven lives of pharmacology's first lead structure. *Drug Discovery Today* **2011**, *16*, 119.
- (14) Song, C.; Liu, K.; Wang, Z.; Ding, B.; Wang, S.; Weng, Y.; Chiang, C.-W.; Lei, A. Electrochemical oxidation induced selective tyrosine bioconjugation for the modification of biomolecules. *Chem. Sci.* **2019**, *10*, 7982.
- (15) Li, B. X.; Kim, D. K.; Bloom, S.; Huang, R. Y.-C.; Qiao, J. X.; Ewing, W. R.; Oblinsky, D. G.; Scholes, G. D.; MacMillan, D. W. C. Site-selective tyrosine bioconjugation via photoredox catalysis for native-to-bioorthogonal protein transformation. *Nat. Chem.* **2021**, *13*, 902.
- (16) (a) Fatykhov, R.; Khalymbadzha, I.; Chupakhin, O. Cross-Dehydrogenative Coupling Reactions between Phenols and Heteroarenes: Modern Trends in Cross-Coupling Chemistry of Phenols. *Adv. Synth. Catal.* **2022**, *364*, 1052. (b) Patureau, F. W. The Phenol-Phenothiazine Coupling: an Oxidative Click Concept. *ChemCatChem.* **2019**, *11*, 5227.
- (17) (a) Benchouaia, R.; Nandi, S.; Maurer, C.; Patureau, F. W. O₂-Mediated Dehydrogenative Phenoxazination of Phenols. *J. Org. Chem.* **2022**, *87*, 4926. (b) Zhao, Y.; Huang, B.; Yang, C.; Xia, W. Visible-Light-Promoted Direct Amination of Phenols via Oxidative Cross-Dehydrogenative Coupling Reaction. *Org. Lett.* **2016**, *18*, 3326. (c) Jin, R.; Patureau, F. W. Mild, Periodate-Mediated, Dehydrogenative C-N Bond Formation with Phenothiazines and Phenols. *Org. Lett.* **2016**, *18*, 4491. (d) Louillat-Habermeyer, M.-L.; Jin, R.; Patureau, F. W. O₂-mediated dehydrogenative amination of phenols. *Angew. Chem., Int. Ed.* **2015**, *54*, 4102.
- (18) For further details, see the [Supporting Information](#).
- (19) For the use of Ag₂O in combination with a Te(II) catalyst in the coupling of simple phenols with phenothiazines, see: Vemuri, P. Y.; Cremer, C.; Patureau, F. W. Te(II)-Catalyzed Cross-Dehydrogenative Phenothiazination of Anilines. *Org. Lett.* **2022**, *24*, 1626.
- (20) Goswami, M.; Konkol, A.; Rahimi, M.; Louillat-Habermeyer, M.-L.; Kelm, H.; Jin, R.; de Bruin, B.; Patureau, F. W. Mechanism of the Dehydrogenative Phenothiazination of Phenols. *Chem.—Eur. J.* **2018**, *24*, 11936.
- (21) Parker, D.; Fradgley, J. D.; Wong, K. L. The Design of Responsive Luminescent Lanthanide Probes and Sensors. *Chem. Soc. Rev.* **2021**, *50*, 8193.
- (22) Sánchez-Fernández, R.; Sánchez-Temprano, A.; Esteban-Gómez, D.; Pazos, E. Probing Tyrosine Nitration with a Small Tb(III)-Metallopeptide. *ChemBioChem.* **2023**, No. e202300072.