

Vitamin D and Its Synthetic Analogs

Miguel A. Maestro,[†] Ferdinand Molnár,[‡] and Carsten Carlberg^{*,§}


[†]Departamento de Química-CICA, Universidade da Coruña, ES-15071 A Coruña, Spain

[‡]School of Science and Technology, Department of Biology, Nazarbayev University, KZ-010000 Astana, Kazakhstan

[§]School of Medicine, Institute of Biomedicine, University of Eastern Finland, FI-70211 Kuopio, Finland

Supporting Information

ABSTRACT: For many individuals, in particular during winter, supplementation with the secosteroid vitamin D₃ is essential for the prevention of bone disorders, muscle weakness, autoimmune diseases, and possibly also different types of cancer. Vitamin D₃ acts via its metabolite 1 α ,25-dihydroxyvitamin D₃ [1,25(OH)₂D₃] as potent agonist of the transcription factor vitamin D receptor (VDR). Thus, vitamin D directly affects chromatin structure and gene regulation at thousands of genomic loci, i.e., the epigenome and transcriptome of its target tissues. Modifications of 1,25(OH)₂D₃ at its side-chain, A-ring, triene system, or C-ring, alone and in combination, as well as nonsteroidal mimics provided numerous potent VDR agonists and some antagonists. The nearly 150 crystal structures of VDR's ligand-binding domain with various vitamin D compounds allow a detailed molecular understanding of their action. This review discusses the most important vitamin D analogs presented during the past 10 years and molecular insight derived from new structural information on the VDR protein.



INTRODUCTION

An UV-B (290–315 nm)-dependent, nonenzymatic reaction in human skin converts the cholesterol precursor 7-dehydrocholesterol into previtamin D₃ that further isomerizes into vitamin D₃ (calciferol, **1**)¹ (Figure 1). Similarly, UV-B-radiated plants and mushrooms are able to produce the isomer vitamin D₂ (ergocalciferol, **2**) based on their membrane sterol ergosterol.² Both secosteroids are themselves biologically inert and have to be activated by hydroxylation first at C-25, leading to the prehormones 25-hydroxyvitamin D₃ [25(OH)D₃, (calciol, **3**)] and 25(OH)D₂, and then at C-1, creating 1,25(OH)₂D₃ (calcitriol, **4**)³ and 1,25(OH)₂D₂, respectively.

25(OH)D₃ is the metabolically most stable and abundant vitamin D metabolite, and its serum levels serve as a biomarker of the vitamin D status of individuals.⁴ The biologically active form of vitamin D₃, 1,25(OH)₂D₃, acts via activation of the transcription factor VDR as a nuclear hormone that directly affects gene regulation.⁵ The physiological role of vitamin D is the regulation of calcium homeostasis for maintaining bone mineralization⁶ as well as the modulation of innate and adaptive immunity⁷ for improving the response to infections by microbes, such as *Mycobacterium tuberculosis*,⁸ and preventing autoimmune diseases, such as multiple sclerosis.⁹

Lifestyle decisions, such as staying predominantly indoors and covered by textile outdoors, combined with changes in seasons and climate cause, for many individuals, insufficient exposure to UV-B and thus low endogenous production of vitamin D₃. Human diet is often rather low in vitamin D

because only fatty fish and UV-B irradiated mushrooms have reasonable quantities of the vitamin D₃ or vitamin D₂, respectively. The fortification of milk, margarine, and juices with vitamin D₃ or vitamin D₂ is applied in some countries. Moreover, in winter months daily supplementation with at least 25 μ g (1000 IU) of vitamin D₃ is recommended in order to prevent vitamin D deficiency.¹⁰ The latter not only would result in rickets in children and in a higher risk of bone fractures due to osteoporosis or osteomalacia in adults,¹¹ but also will compromise the function of the immune system and the claimed preventive actions of vitamin D against cardiovascular diseases, diabetes, neuropsychiatric disorders, and cancer.¹² Supplementation with vitamin D₃ clearly increased in the general population, e.g., the sales of vitamin D supplementation products increased within 1 decade nearly 15-fold.¹³

There is no doubt that a sufficient vitamin D status is important for bone health,¹⁴ but overdosing with vitamin D₃, 1,25(OH)₂D₃, or its synthetic analogs may result in tissue calcification.¹⁵ Symptoms of hypercalcemia are (i) digestive distress, such as vomiting, nausea, and stomach pain, (ii) fatigue, dizziness, and confusion, (iii) excessive thirst, and (iv) frequent urination. However, hypercalcemia occurs rarely and no other severe side effects or toxicity of vitamin D overdosing is known. Nevertheless, higher doses of vitamin D₃ are not

Received: January 31, 2019

Published: March 27, 2019




Figure 1. Vitamin D metabolites and analogs available on the market.

Table 1. Vitamin D Compounds on the Market^a

- Vitamin D₃ (1, calciferol) is used worldwide in the prevention of vitamin D deficiency and associated diseases, such as rickets [Vigantol (Bayer), Delsterol (DuPont), Duphafrol-D₃ (multiple pharma companies), Lutavit D₃ (BASF), Vi-D₃, Videkhol, Vigosan (multiple pharma companies)].
- Vitamin D₂ (2, ergocalciferol) is also used in the prevention of vitamin D deficiency and associated diseases, such as rickets (marketed with different names by multiple pharma companies).
- Calcidiol (3, 25(OH)D₃) is used in the treatment of chronic hypocalcemia, renal osteodystrophy [Calderol (Upjohn), Hidroferol (Faes Farma)], rickets [Dedrogyl (Roussel), Hidroferol (Faes Farma)].
- Calcitriol [4, 1,25(OH)₂D₃] is prescribed for renal osteodystrophy [Rocatrol (Roche), Calcijex (Abbott)], osteoporosis [Rocatrol (Roche)] and psoriasis [Silkis (Galderma)].
- Calcipotriol [5, 22-ene-26,27-dehydro-1,25(OH)₂D₃] is used for psoriasis [Davionex (Leo Pharmaceuticals), Dovonex (Warner Chilcott)].
- Doxercalciferol [6, 1α(OH)D₃, Hectrol (Bone Care International)] is prescribed for secondary hyperparathyroidism.
- Alfalcidol (7, 1α(OH)D₃) is used for renal osteodystrophy [Alfarol (Chugai Pharmaceutical), One-Alpha (Leo Pharmaceuticals)], secondary hyperparathyroidism [Alfarol (Chugai Pharmaceutical)], osteoporosis [Alfarol (Chugai Pharmaceutical), Alpha D₃ (Teva Pharmaceuticals)] and rickets [Alfarol (Chugai Pharmaceutical)].
- Tacalcitol (8, 1α,24(OH)₂D₃) is prescribed for psoriasis [Bonalfa (Teijin), Curatoderm (Merck KGaA)].
- Paricalcitol [9, 19-nor-1,25(OH)₂D₃, Zemplar, (Abbott Laboratories)] is used for secondary hyperparathyroidism.
- Oxacalcitriol (10, 22-oxa-1,25(OH)₂D₃) is used for secondary hyperparathyroidism and psoriasis [Oxarol (Chugai Pharmaceuticals)] in Japan.
- Falecalcitriol [11, 1,25(OH)₂-26,27-F₆-D₃] is prescribed for secondary hyper-parathyroidism in Japan [Hornel (Taisho Pharmaceuticals and Sumitomo Pharmaceuticals), Fulstan (Kissei Pharmaceuticals)].
- Eldecalcitol [12, 2α-(3-hydroxypropoxy)-1,25(OH)₂D₃] is prescribed for osteoporosis only in Japan [Edirol (Chugai Pharmaceutical)].

^aOnly a few vitamin D compounds have reached the market.^{88,89} Their applications, commercial name, and company are listed. The structures of the compounds are shown in Figure 1.

recommended as nutritional supplement for reaching non-skeletal effects of the vitamin. Similarly, the main goal of the development of vitamin D analogs is to identify compounds with a low calcemic effect versus a potent antiproliferative, prodifferentiating, and/or immune-modulatory function.

In total, more than 3000 synthetic vitamin D analogs were developed by various pharmaceutical companies and academic research groups in order to advance the biological properties of the natural compound for a applications in the therapy of (i) hyperproliferative diseases, such as different types of cancer, (ii) psoriasis, an autoimmune disease of the skin,¹⁶ or (iii) bone disorders, such as osteoporosis.¹⁷ However, so far only a few vitamin D compounds made it to the market (Table 1). In

addition to vitamin D₃ being extensively used as a nutritional supplement, the commercially most successful vitamin D analog is calcipotriol (5), which is topical agent in clinical use for the treatment of psoriasis. Together with the compounds doxercalciferol (6), alfalcidol (7), tacalcitol (8), paricalcitol (9), oxacalcitriol (10), falecalcitriol (11), and eldecalcitol (12) it had been discussed in previous reviews^{18,19} (Figure 1). In contrast, despite promising in vitro results, analogs such as inecalcitol (13) or seocalcitol (14) were unsuccessful in phase II clinical trials of acute myeloid leukemia (www.hybrigenics.com/news/articles/list/type/2) or pancreatic cancer,²⁰ respectively. Interestingly, some immune-system-related vitamin D target genes, such as cathelicidin antimicrobial peptide




Figure 2. $1,25(\text{OH})_2\text{D}_3$ complexed to the VDR-LBD. The VDR-LBD has a conserved 3D architecture, which is made of a three-layer α -helical sandwich. In the lower part of the LBD the LBP is located. All the helices are labeled from N-terminus toward C-terminus and numbered in white color (A). Details on the LBP with bound $1,25(\text{OH})_2\text{D}_3$ and critical amino acids that provide anchoring contacts for the three OH groups (B). Details on the conformation of the bound $1,25(\text{OH})_2\text{D}_3$ molecule with the annotated OH groups and highlights to its contribution of its activity. The numbering of the carbon atoms is indicated (C). The figure is based on the PDB code 1DB1.





Figure 3. Vitamin D signaling. $25(\text{OH})_2\text{D}_3$ is converted by the enzyme CYP27B1 to its biologically most active form $1,25(\text{OH})_2\text{D}_3$, which binds to the transcription factor VDR. Upon binding of $1,25(\text{OH})_2\text{D}_3$ or synthetic agonists, a conformational change in the LBD is induced leading to cofactor exchanges shifting the balance toward recruitment of coactivator proteins. Co-repressor proteins dissociate from the VDR-RXR heterodimer. In parallel, the mediator complex and chromatin modifying enzymes (readers, writers, and erasers) are recruited in order to handle histone proteins of local nucleosomes around genomic VDR binding sites. In addition, chromatin remodeling complexes are recruited and rearrange nucleosomes at vitamin D-sensitive chromatin regions. Altogether, these changes lead to looping of the distal regulatory elements toward the basal transcriptional machinery with RNA polymerase II and other nuclear adaptor proteins initiating the start of $1,25(\text{OH})_2\text{D}_3$ -dependent transcription from hundreds to thousands of TSS regions throughout the whole human genome. The ultimate outcome is the increase or decrease of the of primary vitamin D target gene expression followed by changes of indicated cellular functions.

(CAMP) and CD14 (encoding for a Toll-like receptor 4 co-receptor), are very responsive,^{21,22} while there are no vitamin D target genes with comparable inducibility involved in the management of cellular growth and differentiation. The failure of anticancer trials and the success in the therapy of an immune disease as well as prominent gene regulatory effects in immune cells suggest that VDR ligands, in addition to bone-related functions, may rather have a therapeutic potential in immune diseases than in cancer.

The majority of synthetic VDR ligands are direct derivatives of $1,25(\text{OH})_2\text{D}_3$, but within the past years an increasing number of vitamin D mimics were published. $1,25(\text{OH})_2\text{D}_3$ had been modified at its side-chain, A-ring (often together with side-chain changes), triene system, and C-ring. These modifications follow the strategy to increase the VDR binding affinity while in parallel modulating the metabolic stability of the molecules.¹⁸ A reasonable number of new vitamin D analogs have been published within the past years and will be



Family	compound	VDR aff.	Prolif inh.	Transac act.	Calcemia
20(22)-yn-	15	+			
	16a	=			
	16b	=			
	16c	-			
	17a	++			
	17b	++			
Gemini	2S-18	+++			
	2R-18	+++			
	2S-19	++			
	2R-19	+++			
	2S-20	+			
	2R-20	++			
	21		+++		
	22		+++		
Carboranyl	23		+++		
	24	++	++	++	-

Ref: 1,25(OH)₂D₃: = = = =

Figure 4. Side-chain-modified vitamin D analogs. The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10× higher; (++) >100× higher; (+++) >1000× higher; (–) >10× lower; VDR aff, VDR affinity; Prolif inh, proliferation inhibition; Transac act, VDR transactivation activity; calcemia, [Ca²⁺] level changes in serum.

discussed in this review. Moreover, the number of solved VDR crystal structures with synthetic ligands has significantly grown.

■ CENTRAL ROLE OF VDR IN VITAMIN D SIGNALING

VDR is the only protein expressed by the human genome that is able to bind 1,25(OH)₂D₃ and its analogs at subnanomolar concentrations.²³ Thus, all physiological functions of vitamin D compounds are mediated by VDR and its target genes.²⁴ The VDR gene is expressed most prominently in intestine, kidneys,

and bone, but in most of the other 400 human tissues and cell types some VDR expression is found.²⁵ This means that not only tissues that relate to calcium homeostasis and bone formation but also immune cells respond to vitamin D.²⁶

VDR is an endocrine receptor and member of the superfamily of nuclear receptors; i.e., the mechanisms of its action are comparable to the receptors for glucocorticoids and estrogen.²⁷ VDR's ligand-binding domain (LBD) is structurally conserved and comprises 11–15 α -helices, modestly varies

between solved crystal complexes, and depends on the folding of the intrinsically disordered region between helices H1 and H3 and the presence of a helix HX between helices H11 and H12^{28,29} (Figure 2A). The lower part of the LBD contains a ligand-binding pocket (LBP), which is a cavity with a volume of $\sim 700 \text{ \AA}^3$ (with possible expansion beyond 1000 \AA^3) being formed by some 40 mostly nonpolar amino acids.³⁰ Three pairs of polar amino acids within the LBP fix via hydrogen bonds each one of the three OH groups (at C-1 α , C-3 β , and C-25) of $1,25(\text{OH})_2\text{D}_3$. The 1 α -OH group interacts with Y143 (helix H1) and S278 (helix H5), the 3 β -OH group contacts S237 (helix H3) and R274 (helix H5), and the 25-OH group interferes with H305 (loop between helices H6 and H7) and H397 (helix H11)²⁸ (Figure 2B).

VDR ligands induce a conformational shift to the LBD, which replaces co-repressor proteins by coactivator proteins; i.e., ligand binding induces a different protein–protein interaction profile of the receptor.³¹ VDR agonists cause an efficient dissociation of co-repressors from the LBD and allow the specific binding of coactivators and the mediator complex (Figure 3). Coactivators also attract chromatin modifying enzymes that write, erase, or read post-translational marks of histones, such as acetyl and methyl groups, to histone proteins of nucleosomes in the vicinity of genomic VDR binding sites.³² Moreover, also members of chromatin remodeling complexes interact in a ligand-dependent fashion with VDR and cause a rearrangement of nucleosomes at vitamin D-sensitive chromatin regions.³³ These epigenetic changes allow looping of VDR-bound enhancers toward accessible transcription start sites (TSSs) at hundreds to thousands of loci throughout the human genome.³⁴ These enhancer-TSS assemblies are triggered by ligand-activated VDR and finally result in an increase or decrease in the expression of hundreds of primary vitamin D target genes (Figure 3).

The structure of the human VDR-LBD complexed with $1,25(\text{OH})_2\text{D}_3$ was solved in the year 2000.²⁸ Since then altogether 143 human, rat, and zebrafish VDR-LBDs have been crystallized with a large number of synthetic analogs³⁵ (Table S1). In general, the analogs behave like $1,25(\text{OH})_2\text{D}_3$ by stabilizing the LBD in more or less the same conformation, since the three OH groups of each vitamin D compound take up a nearly identical position. This suggests that there is only one agonistic conformation of the LBD for which the interaction between the ligand's 25-OH group and the LBP amino acids H305 and H397 are most important (Figure 2C).

On the basis of the vitamin D analog's chemical modification, all solved VDR-ligand-complexes can be divided to the six groups: (i) A-ring modifications, (ii) side-chain modifications, (iii) triene system modifications, (iv) combined A-ring and side-chain modification, (v) modifications in the CD-ring, and (vi) nonsteroidal analogs. All modifications aim to either (i) maintain the three anchoring OH groups at the same position as in $1,25(\text{OH})_2\text{D}_3$ and/or (ii) fill the LBP most efficiently in order to form additional hydrogen network and/or hydrophobic contacts. More variant modifications of $1,25(\text{OH})_2\text{D}_3$ aim to alter the ligand conformation or to bounce the shape of the LBP by adding an additional side-chain at positions C-20 or C-22. Moreover, de novo designed nonsteroidal compounds carry modifications, such as the exchange the classical secosteroid ring structure by rings with aromatic character. The aim with these molecules is to maintain the hydrophobic interactions with amino acid residues lining the inner surface of the LBP as well as to

increase the stacking interaction with aromatic amino acid residues.

In this review we discuss different classes of vitamin D analogs and, where applicable, provide molecular understanding from VDR crystal structures.

■ SIDE-CHAIN MODIFICATIONS

The first locked side-chain vitamin D analogs nor-21-20(22),23(24)-diyn-1,25(OH)₂D₃ (**15**), nor-21,23,24,25,26,27-20(22)-yn-22-(3-hydroxyphenyl)-1,25(OH)₂D₃ (**16a**), nor-21,23,24,25,26,27-20(22)-yn-22-[3-(hydroxymethyl)phenyl]-1,25(OH)₂D₃ (**16b**), and nor-21,23,24,25,26,27-20(22)-yn-22-[4-(hydroxymethyl)phenyl]-1,25(OH)₂D₃ (**16c**) have been synthesized by convergent route through a Wittig–Horner approach starting from Inhoffen–Lythgoe diol³⁶ (Figure 4). These analogs lead to significant activation of VDR-dependent transcription in comparison to $1,25(\text{OH})_2\text{D}_3$. An unique structural modification on the C-22-diyne analog, a C-17-methyl substitution, was provided through a vinyl(pinacolato)boronate approach and resulted in the C-17-methyl-substituted vitamin D analogs nor-21-20(22),23(24)-diyn-17-methyl-1,25(OH)₂D₃ (**17a**) and nor-21-20(22),23(24)-diyn-17-methyl-26,26,26,27,27,27-hexafluoro-1,25(OH)₂D₃ (**17b**).³⁷ The C-22-aromatic-substituted analogs are less potent in activating VDR than the C-22-diyne isomers. The C-17-methyl analogs bind more efficiently to VDR than $1,25(\text{OH})_2\text{D}_3$.

The two side-chain analog Gemini comprises an unaltered side-chain of $1,25(\text{OH})_2\text{D}_3$ and a second chain at C-20.^{38,39} Although the volume of Gemini is increased by some 25%, it still fits into VDR's LBP.³⁰ One side-chain of Gemini takes the same place as that of $1,25(\text{OH})_2\text{D}_3$, whereas an extra subcavity opens within the LBP for the second side-chain.⁴⁰ The increase in transcriptional activity of Gemini⁴¹ motivated the preparation of Gemini-type analogs with side-chains containing double or triple bonds and isohexafluoro-2-propanol or isohexadeutero-2-propanol side-chain ends. Compounds **18–20** have been synthesized with both configurations at C-20 by a convergent approach through Wittig–Horner coupling starting from Inhoffen–Lythgoe diol.⁴² (*R*)-Analog showed higher antiproliferative potency in MCF10CA1 human breast cancer cells than their (*S*)-counterparts, and both were 100–1000 times more potent than $1,25(\text{OH})_2\text{D}_3$. Furthermore, both configurations of the Gemini derivatives are also more potent than $1,25(\text{OH})_2\text{D}_3$ in inducing the differentiation of NB4 human leukemia cells. Thus, Gemini compounds have enhanced potency in inhibiting proliferation and inducing differentiation with reduced induction of hypercalcemia when compared to $1,25(\text{OH})_2\text{D}_3$. Moreover, C-20 methyl-substituted Gemini analogs (**21–23**) are also potent in the inhibition of HL-60 human leukemia cell proliferation and the induction of *CAMP* gene expression.⁴³

1 α -Hydroxy-25,26,27-trinor-24-*o*-carboranyl-vitamin D₃ ($1,25\text{cD}_3$, **24**, Figure 4) is a rather new vitamin D analog, in which an *o*-carborane moiety replaces the 25-OH group.⁴⁴ Despite the lack of this critical group, $1,25\text{cD}_3$ is as effective as $1,25(\text{OH})_2\text{D}_3$ in inhibiting the growth of MCF-7 human breast cancer cells and in inducing the differentiation of HaCaT human keratinocytes. VDR binds $1,25\text{cD}_3$ 2 times tighter than $1,25(\text{OH})_2\text{D}_3$ and is equally potent as the natural hormone in inducing reporter gene activity while not showing adverse calcemic effects. Moreover, like most other vitamin D analogs, the conformation of the complex of $1,25\text{cD}_3$ with




Figure 5. Structure–function relationship of VDR ligands (1). The carborane group of **1,25cD₃** creates additional hydrophobic interactions that compensate for the loss of the 25-OH group. All residues that have conserved interactions are shown in gray (top). Detailed interaction with residues based on PDB code structures 2HC4 (**1,25(OH)₂D₃**) and 5E7V (**1,25cD₃**). The displayed interactions are identified under cutoff 3.5 Å (A). Destabilization of the VDR-LBD upon binding of 23,36-lactone analogs. Representation of crystal structure *b*-factors using structures PDB codes 1RK3 (**1,25(OH)₂D₃**, left) and 3A2H (TEI-9647, right). Regions with the highest *b*-factors are highlighted in red and they are helices H9–10 that may affect heterodimerization with RXR, helix H11 affecting the position of helix H12, and the coactivator peptide showing very high *b*-factors overall. The most stable part of the VDR is shown in blue through green, yellow and red monitoring the highest *b*-factor values (B). Structural implication of 22S-alkyl-2-methylene-19-nor-**1,25(OH)₂D₃** binding. The position of the helix H12 takes the same conformation in both 22S-alkyl-2-methylene-19-nor-**1,25(OH)₂D₃** and **1,25(OH)₂D₃**. Many residues around the two aliphatic chains move or rotate, such as H301 (loop helices H6/7), H393 (helix H11), or F418 (helix H12). Structural elements from 22S-alkyl-2-methylene-19-nor-**1,25(OH)₂D₃** (PDB code 2ZXM) and **1,25(OH)₂D₃** (PDB code 1RK3) complexes are highlighted in green and white color, respectively (C).


VDR's LBD is highly similar to that of **1,25(OH)₂D₃**; i.e. the protein shows the same topology. Nevertheless, the loop between helices H6 and H7, which is a critical region for the activation of the receptor, and the last part of helix H11 show shifts by 0.6 Å. The carborane side-chain is 2.4 Å longer than that of **1,25(OH)₂D₃**, but it is hydrophobic and therefore favors the interaction with hydrophobic amino acid in this part of the VDR-LBP. This compensates for the loss of the 25-OH group (Figure 5A). Thus, the collection of small changes stabilizes helices H3, H11, and H12 and overall causes higher stability of VDR's LBD.

■ A-RING MODIFICATION

The biological profiles of the C-2-substituted vitamin D analogs 2-(3'-hydroxypropyl)-**1,25(OH)₂D₃** (**25**), 2-butyl-

1,25(OH)₂D₃ (**26**), 2-(4'-hydroxybutyl)-**1,25(OH)₂D₃** (**27**), 2-(2'-hydroxyethoxy)-**1,25(OH)₂D₃** (**28**), 2-(3'-hydroxypropoxy)-**1,25(OH)₂D₃** (**29**), and 2-(4'-hydroxybutoxy)-**1,25(OH)₂D₃** (**30**) (Figure 6) indicated that C-2β-substituted analogs have higher affinity for the serum vitamin D binding protein (DBP) and lower affinity for VDR but are superior to C-2α analogs in reporter gene assays.⁴⁵

The analog 2-(3'-hydroxypropyl)-1α-methyl-**1,25(OH)₂D₃** (**32**) was synthesized through a Pd-catalyzed ring-closure of enyne, and coupling with vinyl bromide and showed a 2-fold higher potency than **1,25(OH)₂D₃** in reporter gene assays.⁴⁶ Interestingly, when VDR's amino acid R274, which contacts the 1α-OH group of the ligand (Figure 2), is mutated to a hydrophobic residue (R274L), the compound is even 7 times more potent than the natural hormone, suggesting that the 1α-



Family	compound	VDR aff.	24OH trans	HL-60 diff
2 α -subst	25	++	-	=
	26	--	--	--
	27	=	--	=
	28	=	--	=
	29	++	--	=
	30	--	-	--
	31		++	
4-OH	32		--	
	33a	-		
2-methylen	33b	=		
	34a	--	--	-
	34b	=	-	+
	35	=	++	++

Ref: 1,25(OH)₂D₃: = = =

Figure 6. Vitamin D analogs with A-ring modifications. The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10 \times higher; (-) >10 \times lower; (--), >100 \times lower; VDR aff, VDR affinity; 24OH trans, CYP24A1 transactivation activity; HL60 diff, HL-60 cell differentiation induction.

methyl group is stabilized primarily by hydrophobic interactions.

The compounds 1 α ,4 α ,25(OH)₃D₃ (**33a**) and 1 α ,4 β ,25-(OH)₃D₃ (**33b**) were created through a Pd-catalyzed ring-closure and coupling and showed lower affinity for VDR than 1,25(OH)₂D₃.⁴⁷ However, the 4 β -analog (**33b**) displays higher VDR affinity and potency in reporter gene assays than the 4 α -compound (**33a**).

The analogs 2-methylene-25-(OH)D₃ (**34a**) and 2-methylen-20-epi-25-(OH)D₃ (**34b**) belong to the family of the potent lead compound 2-methylene-1,25(OH)₂D₃ (**35**, 2MD) and were synthesized through a Pd-catalyzed coupling between an enol triflate and an enyne.⁴⁸ These molecules are defined by the relocation of the exocyclic methylene group from C-10 to C-2 and the inversion of the C-20 configuration. Compound **34a** shows lower affinity for VDR than 1,25(OH)₂D₃, lower potency in inducing HL-60 cell differentiation and in reporter gene assays, while compound **34b** displays the same affinity for VDR as 1,25(OH)₂D₃, higher potency in HL-60 cell differentiation induction, and lower activity in reporter gene assays.

■ TRIENE SYSTEM MODIFICATIONS

The compounds PRI-1731 (**36**), PRI-1732 (**37**), PRI-1733 (**38**), and PRI-1734 (**39**) represent a series of vitamin D analogs with a branched side-chain (E)-stereochemistry at the C-5/C-6 double bond, both configurations at C-24 and a C-22/C-23 double bond or an OH group at C-22^{49,50} (Figure 7). They have moderate prodifferentiating activities on HL-60 cells and their maximal inhibition of proliferation ranged from




Figure 7. Triene system modified vitamin D analogs.




Figure 8. Vitamin D analogs with side-chain and A-ring modifications (I). The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10× higher; (++) >100× higher; (−) >10× lower; (--) >100× lower; VDR aff, VDR affinity; Antag act, antagonistic activity.

10% to 15% of that for 1,25(OH)₂D₃ and 20–30% of that for 1,25(OH)₂D₂.

■ SIDE-CHAIN AND A-RING MODIFICATIONS

25-Dehydro-1 α -hydroxy-vitamin D₃-26,23 lactones with double modifications of C-24 and C-2 α were synthesized via a convergent approach by Pd-catalyzed ring closure of a enyne and subsequent coupling with a functionalized vinyl bromide.^{51,52} Numerous analogs have been synthesized with variations in their C-23 configuration, C-24 mono- or disubstitution [H, Me, Et, *c*-Pr, *n*-Pr, *n*-Bu, *i*-Bu (40, 41, 48,

50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72)] and C-2 α substitution [H, Me, CH₂CH₂CH₂OH, OCH₂CH₂CH₂OH, (42–47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73)] (Figure 8). The principal characteristic of these analogs is their antagonist activity, probably due to locking the VDR-LBD in a conformation where it does not effectively interact with coactivator proteins. This can be seen also from the values of the β -factors found in the crystal structure (PDB code 3A2H) of the lactone analog TEI-9647, where the coactivator peptide shows very high values compared to 1,25(OH)₂D₃. In addition, helices H9 and 10 as well as H11 and to some




Figure 9. Vitamin D analogs with side-chain and A-ring modifications (II). The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10× higher; (++) >100× higher; (+++) >1000× higher; (–) >10× lower; (--) >100× lower; VDR aff, VDR affinity; MCF10 prol, MCF10 cell proliferation inhibition; NB4 diff, NB4 cell differentiation induction; MG63 trans, MG63 cell transactivation activity; U963 prol, U963 cell proliferation inhibition; HL60 diff, HL-60 cell differentiation induction.

extent H12 show higher fluctuation leading to overall destabilization of the LBD (Figure 5B). Structure–activity relationship studies demonstrated that the exomethylene group of the lactone is indispensable, the C-23S configuration provides higher activity, and an appropriate combination of C-24 and C-2 α substitution obtains the highest antagonist potency.

19-nor (**74**) and 1 α -F (**75**) Gemini analogs containing triple bonds and isohexafluoro-2-propanol or isohexadeutero-2-propanol side-chain end groups⁴² (Figure 9) were synthesized. Again (*R*)-analogues are more potent in inhibiting MCF10CA1 cell proliferation than their (*S*)-counterparts. Both isomers are active already at 100–1000 times lower concentrations than 1,25(OH)₂D₃. In both configurations the Gemini analogs are

equally potent to 1,25(OH)₂D₃ in inducing NB4 cell differentiation but are less calcemic than the natural hormone.

A series of 2-methylene-19-nor vitamin D analogs with or without C-22S alkyl substitution, such as 2-methylene-19,25,26,27-tetranor-vitamin D₃ (**76**), 2-methylene-19,25-dinor-vitamin D₃ (**77**), 2-methylene-19,26,27-trinor-vitamin D₃ (**78**), and 2-methylene-19-nor-vitamin D₃ (**79**) and their C-22S alkyl derivatives (a, R = H; b, R = Et; c, R = Bu) have been prepared and biologically tested.⁵³ The side-chain modifications in **76**, **77**, and **78** reduce the VDR binding affinity 10-fold compared to 1,25(OH)₂D₃. Interestingly, an increasing size of the C-22 substituent in 2-methylene-19-nor-vitamin D₃ (**79**) results in a decreased VDR binding affinity compared to 1,25(OH)₂D₃. Compounds with a normal side-chain (**79a**,





Figure 10. Vitamin D analogs with side-chain and A-ring modifications (III).

79b, and 79c) show strong activation in reporter gene assays and compounds without C-22 substitution (76a, 77a, 78a, and 79a) even full agonist activity. In contrast, C-22S butyl-substituted molecules (76c, 76c, 78c, and 79c) present little transactivation potency, while C-22S ethyl-substituted compounds (76b, 77b, 78b, and 79b) display intermediate activity. Moreover, the analogs 76a, 77a, 78a, and 79a induce the recruitment of the VDR partner receptor retinoid X receptor (RXR, Figure 3) and of a coactivator peptide in a concentration dependent manner, while C-22S-substituted compounds cause only moderate effects.

VDR-LBD crystal structures complexed with further 22S-alkyl-2-methylene-19-nor-1,25(OH)₂D₃ derivatives (80–83)⁵⁴ confirmed that the compounds trigger the creation of an extra cavity of the LBP by rotating L305 about 27° outward, in order to shelter the butyl group (Figure 5C). Ligands act as VDR antagonists when they do not interact with the C-terminal helix H12 of the receptor. Interestingly even though the position of helix H12 is maintained in the agonistic position, most likely forced by the presence of the coactivator peptide, there is a shift or rotation of multiple residues away from the 22-butyl analog. These changes weaken the interaction with H301 (loop H6/7), H393 (H11), or F418 (H12) and destabilize this



Family	compound	VDR aff	24OH trans	HL60 diff
2-methylen	116a	-	--	-
	116b	=	++	=
	117a	++	+++	+++
	117b	--	--	--
	118a	++	++	-
	119a	--	--	--
	119b	--	--	--
	120a	--	=	+
	120b	-	+	+
	121a	--	--	--
	121b	-	--	--
	122a	--	=	+
	122b	-	=	+

Ref: 1,25(OH)₂D₃: = = =

Figure 11. Vitamin D analogs with side-chain and A-ring modifications (IV). The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10× higher; (++) >100× higher; (+++) >1000× higher; (-) >10× lower; (--) >100× lower; VDR aff, VDR affinity; 24OH trans, CYP24A1 transactivation activity; HL60 diff, HL-60 cell differentiation induction.

region of the LBD (Figure 5C). Interestingly, in the presence of a coactivator peptide some of the antagonistic 22-butyl analogs take the agonistic conformation. Whether this is a technical artifact of the crystallization or has a physiological meaning, such as sensing of cofactor balance in the cellular context, needs to be clarified.

20-epi-Eldecalcitol (**84**), a 20-epi derivative of the antiosteoporotic drug eldecalcitol (**12**) (Table 1 and Figure 1), was synthesized through a convergent approach by Pd-catalyzed ring-closure of an enyne and coupling with vinyl bromide.⁵⁵ Since 20-epi-1,25(OH)₂D₃, a diastereomer of 1,25(OH)₂D₃ possessing an inverted C-21 methyl-substituent at C-20, shows enhanced biological activities compared to 1,25(OH)₂D₃, compound **84** displays a 50-fold increased inhibition of U937 human leukemia cell proliferation.⁵⁶

Since the presence of a 1 α -OH group in 1,25(OH)₂D₃ is crucial for VDR binding, its replacement with a 1 β -OH group [1 β ,25(OH)₂D₃, **86**] causes loss of physiological activity⁵⁷ (Figure 10). Replacing the 1 α -OH group with one fluorine atom [1 α -F,25(OH)₂D₃, **87**] also markedly diminishes biological activity,⁵⁸ while a compound with each a fluorine atom at C-1 and C-25 [1,25(F)₂(OH)₂D₃, **88**] is devoid of all activity.⁵⁹ Interestingly, the 3-OH group is not necessary if the 1 α -OH group is already in position, but the lack of the 3-OH group [1 α ,25(OH)₂-3-deoxy-D₃, **89**] reduces the biological activity.^{60,61} Additional methyl groups at C-1 β [1 β -methyl-1,25(OH)₂D₃, **90**] and C-3 α [3 α -methyl-1,25(OH)₂D₃, **91**] significantly reduce VDR binding affinity. Switching the 3-OH group from β to α position [3-epi-1,25(OH)₂D₃, **92**] causes a drastic reduction of physiological activity.⁶² In fact,




Figure 12. Vitamin D analogs with side-chain and A-ring modifications (V). The table summarizes the biological properties of the compounds: reference, $1,25(\text{OH})_2\text{D}_3$; (=) similar value; (+) $>10\times$ higher; (++) $>100\times$ higher; (-) $>10\times$ lower; (--) $>100\times$ lower; VDR aff, VDR affinity; HL60 diff, HL-60 cell differentiation induction.

compound **92** is an intermediate of $1,25(\text{OH})_2\text{D}_3$ degradation displaying lower VDR binding affinity and lower calcemic effects compared to $1,25(\text{OH})_2\text{D}_3$. Despite its decreased potency, the in vivo action of compound **92** is tissue-specific.

Taken together, most modifications of the A-ring result in decreased biological activity except for those modified at C-2. Therefore, a large number of C-2-substituted vitamin D analogs have been synthesized and were studied intensively for their biological activity. 2α -Substitutions [2α -methyl- $1,25(\text{OH})_2\text{D}_3$ (**93**)] are more potent than 2β -substitutions [2β -methyl- $1,25(\text{OH})_2\text{D}_3$ (**92**)].⁶³ Elongation of the C-2-alkyl group [2α -ethyl- $1,25(\text{OH})_2\text{D}_3$ (**94**) and 2α -propyl- $1,25(\text{OH})_2\text{D}_3$ (**95**)] reduces VDR binding affinity and biological potency, but ω -hydroxylation restores the activity.⁶⁴ 2α -Methyl- $1,25(\text{OH})_2\text{D}_3$ (**93**) is twice as calcemic as $1,25(\text{OH})_2\text{D}_3$. In combination with 20-epimerization [2α -methyl-20-epi- $1,25(\text{OH})_2\text{D}_3$] VDR binding affinity increases even 12-fold.⁶³ 2α -(3'-Hydroxypropyl)- $1,25(\text{OH})_2\text{D}_3$ (**97**) has a 3-fold increased VDR binding affinity than $1,25(\text{OH})_2\text{D}_3$, while 2β -(3'-hydroxypropyl)- $1,25(\text{OH})_2\text{D}_3$ (**98**) is 1.4 times more potent. Terminal hydroxylation of 2α - and 2β -prooxy groups at C-2, 2α -hydroxypropoxy- $1,25(\text{OH})_2\text{D}_3$ (**99**) and 2β -hydroxypropoxy- $1,25(\text{OH})_2\text{D}_3$ (**100**) also increases the VDR binding potential.

19-nor-Vitamin D analogs are known to be devoid of hypercalcemic and hyperphosphatemic effects.^{65,66} For example, 19-nor- $1,25(\text{OH})_2\text{D}_3$ (**101**) has a 5 times reduced the VDR binding affinity compared to $1,25(\text{OH})_2\text{D}_3$ paired with low or no bone calcification activity, while 19-nor- $1,25(\text{OH})_2\text{D}_2$ (paricalcitol, **9**, Figure 1) has similar affinity as the

natural hormone. Selected modifications at the C-2 position of 19-nor-vitamin D analogs are more potent inducers of gene activity. 2β -(3'-Hydroxypropoxy)-19-nor- $1,25(\text{OH})_2\text{D}_3$ (**102**) and 2α -(3'-hydroxypropoxy)-19-nor- $1,25(\text{OH})_2\text{D}_3$ (**103**) have reduced potency in bone and intestine.⁶⁷ 2-Methylene-19-nor-20-epi- $1,25(\text{OH})_2\text{D}_3$ (**2MD**, **35**) shows VDR binding affinity comparable to $1,25(\text{OH})_2\text{D}_3$ but a 100 times enhanced ability to mobilize calcium from bone.⁶⁸ Moreover, **2MD** is 10 times more potent than $1,25(\text{OH})_2\text{D}_3$ in activating *CYP24A1* gene activity and inducing HL-60 cell differentiation. Analogs of **2MD** with a shortened side-chain, such as **2MP** (**105**) and **2MbisP** (**106**) are able to reduce parathyroid hormone (PTH) production.⁶⁹

The compound 19-nor- $1\alpha,2\beta,25(\text{OH})_3\text{D}_3$ (**107**) is as potent as $1,25(\text{OH})_2\text{D}_3$ in intestinal calcium transport, while its epimer 19-nor- $1\alpha,2\alpha,25(\text{OH})_3\text{D}_3$ (**108**) possesses less activity.⁷⁰ Importantly, both compounds are not calcemic. Compound **107** has higher VDR binding affinity than **108** and is more potent in inhibiting MCF-7 cell proliferation. When the 2-ethylidene group is in *E*-configuration (**109**), VDR binding affinity is 2.4-fold increased. Moreover, a 3'-hydroxypropylidene group at C-2 in *E*-configuration resulted in the potent compounds **AGR** (**110**) and **IAGS** (**111**), which seem to be intestine-selective.⁶⁰

2α -(3'-Hydroxypropyl)-19-nor- $1,25(\text{OH})_2\text{D}_3$ (**112**), which is modified at both C-2 and C-10, shows a similar VDR binding affinity as $1,25(\text{OH})_2\text{D}_3$, displays a 36-fold higher potential in inducing HL-60 cell differentiation, and has a 500-fold higher antiproliferative potency in PZ-HPV-7 prostate cells. Its epimer 2β -(3'-hydroxypropyl)-19-nor- $1,25(\text{OH})_2\text{D}_3$





Figure 13. Structure–function relationship of various VDR ligands (II). The binding of 14-epi-2 α -methyl- and 14-epi-2 β -methyl-1,25(OH)₂-6,7-dehydro-19-norvitamin D₃. The overall spatial conservation of the OH groups is maintained, but the CD-ring structure for the epimers shows higher flexibility by changing the more planar conformation in 1,25(OH)₂D₃ to reverse V-letter shape. This is largely due to the bending of the C-ring downward. The other notable changes involve the proximal location of L227 in 2 β -methyl epimer and W286 for 1,25(OH)₂D₃ under 3.5 Å cutoff. Structural elements from 14-epi-2 α -methyl- (PDB code 3AUQ), 14-epi-2 β -methyl-1,25(OH)₂-6,7-dehydro-19-norvitamin D₃ (PDB code 3AUR), and 1,25(OH)₂D₃ (PDB code 1DB1) complexes are highlighted in green, blue, and white color, respectively (A). Possible mechanism governing the potency of fluorinated analogs. The effect of a fluorinated functional group is illustrated on CF₃ group in comparison with CH₃ group located at the terminal carbons C-26 and C-27 of the ligand's aliphatic chain. The high electronegativity of the fluorine atom has a pulling effect for the hydrophobic residues located in the proximity of the functional group (bottom), which cannot be seen for the 1,25(OH)₂D₃ under the same 3.5 Å cutoff (top). However, the hydrogen bonds between the conserved histidines and the 25-OH group are maintained in both scenarios. The positions of the CF₃ groups show a moderate opening about 5° with maintained distance between carbon C-26 and C-27. These changes have effect on the LBP, which is a bit smaller (middle panel) in the presence of CF₃ functional groups (B). Structural elements for the CF₃ and CH₃ groups are highlighted in orange and white color, respectively.

(113) has lower VDR binding and prodifferentiation activity, but it is as potent as 110 in inhibiting prostate cell proliferation.^{70,71}

Promising analogs with three different alterations in the vitamin D skeleton (the A-ring, the side-chain, and the CD-ring) are 1 β -hydroxymethyl-16-ene-24,24-F₂-26,27-bishomo-25(OH)₂D₃ (QW-1624-F2-2, 114)⁷² and 1 α -fluor-16-ene-20-epi-23-ene-26,27-bishomo-25(OH)₂D₃ (Ro-26.9228, 115).⁷³ In a skin cancer model compound 114 inhibits progression and molecule 115 restores bone loss, while both are not hypercalcemic.

Six new derivatives of compound 104 have been prepared by a convergent synthesis using the Wittig–Horner approach⁷⁴ in order to evaluate the influence of methyl groups at C-22 on biological activity (Figure 11). Single methylation of the (20*R*)-25-hydroxylated side-chain (117) did not change the VDR binding affinity in comparison to the parent compound 104. However, the addition of a 22-methyl group to the (20*S*)-25-hydroxylated side-chain (118) caused a much stronger effect. The 22*R*-compound 118a has a 2.5 times higher VDR binding affinity than 104 and is 250-fold more potent than its 22-epimer 118b. The prodifferentiation potential of an analog



Family	compound	MLR IFN- γ	LPS TNF- α
Gemini	138	=	+++
	139	=	+++
	140	+	--
	141	--	
	142	+	--
	143	=	+
	144	--	
	145	+	--
	146	=	--
	147	-	--
	148	=	=
	149	+++	+++

Ref: 1,25(OH) $_2$ D $_3$: = =

Figure 14. A-ring, D-ring, and side-chain modified vitamin D analogs. The table summarizes the biological properties of the compounds: reference, 1,25(OH) $_2$ D $_3$; (=) similar value; (+) >10 \times higher; (+++) >1000 \times higher; (–) >10 \times lower; (–) >100 \times lower; MLR IFN- γ , INFG inhibition; LPS TNF α , TNF inhibition.

with a 22S-methyl group in the “natural” side-chain (20R) (116a) is 10-fold higher than that of its 22-epimer (116b), whereas in the case of “unnatural” 20S-compound the 22R-epimer (117a) is 1000 times more potent than its 22-epimer (117b) and 4-fold more potent than the parent compound 104. When two methyl groups were introduced at C-22, such as in the 20R-compound (118a) and the 20S-compound (118b), VDR binding affinity is increased compared to their parent molecules.

C-20-isomers of 25(OH)-2-methylene-vitamin D $_3$ and 3-desoxy-1 α ,25(OH) $_2$ -2-methylene-vitamin D $_3$ (117–120) were synthesized through a convergent approach using a Sonogashira coupling⁷⁵ (Figure 11). The biological activities of compounds 119–122 are clearly lower than those of the parent compound 104. With the exception of the 1 α -hydroxylated compounds they were also less active than 1,25(OH) $_2$ D $_3$. Analogs without a 1 α -OH group show lower VDR binding affinity, HL-60 cell prodifferentiation activity, and CYP24A1 activation than those hydroxylated at C-1. The addition of the 10-exo-methylene group improved the in vitro activity of the (20S)-1-desoxy compounds. In contrast, in the (20S)-series only VDR binding affinity augmented. The

presence of the 2-exomethylene group resulted in enhanced intestinal calcium transport compared to 1,25(OH) $_2$ D $_3$, but bone calcium mobilization was 10-fold decreased in the (20R)-series.

A large structure–function analysis of 39 Gemini derivatives⁴³ showed five compounds (123–127) with enhanced antiproliferative activity (Figure 12). Compound 127 was stronger than 1,25(OH) $_2$ D $_3$ in inhibiting cancer cell growth, while both were equipotent in their calcemic effect.

The Sonogashira approach was used to synthesize novel 14-epi derivatives of 19-nor-1 α ,25(OH) $_2$ -previtamin D $_3$ (132) and 19-nor-1 α ,25(OH) $_2$ -tachysterol D $_3$ (135).⁷⁶ Dienynic compounds (129–131) showed moderate VDR binding affinity, where the 2-methylene compound (129) has higher activity than the 2-methyl-substituted diastereomers 130 and 131. Previtamin D $_3$ compounds (132–134) showed low VDR binding affinity. 14-epi-19-nor-tachysterol D $_3$ compounds displayed higher VDR binding affinity, of which 2-methylene-14-epi-19-nor-tachysterol D $_3$ (137) is most potent. Crystal structure analysis indicated unique binding conformations. The binding of both epimers, 14-epi-2 α -methyl- (PDB code 3AUQ) and 14-epi-2 β -methyl-1,25(OH) $_2$ -6,7-dehydro-19-nor-




Figure 15. Nonsteroidal VDR ligands. The table summarizes the biological properties of the compounds: reference, 1,25(OH)₂D₃; (=) similar value; (+) >10× higher; (++) >100× higher; (+++) >1000× higher; (–) >10× lower; (–) >100× lower; (–) >1000× lower; VDR aff, VDR affinity; calcemia, [Ca²⁺] level changes in serum; 24OH tr, CYP24A1 transactivation activity; HL60 diff, HL-60 cell differentiation induction; MCF7 pr, MCF7 proliferation inhibition.

vitamin D₃ (PDB code 3AUR) is very similar in maintaining the position of the anchoring OH groups seen from the

1,25(OH)₂D₃ complex (PDB code 1DB1). However, the modification between C-6 and C-7 provides rigidity for this

region introducing an unforeseen compensation in flexibility for the CD rings. Compared to their more planar conformation in $1,25(\text{OH})_2\text{D}_3$, here they take a reverse V-letter conformation such as by the bending of the C ring about 9° downward. For the 2β -methyl isomer the residue L227 is closer to the ligand under 3.5 \AA cutoff but under similar cutoff W286 seems to be closer to $1,25(\text{OH})_2\text{D}_3$, which is due to more planar conformation of the CD-rings (Figure 13A).

■ SIDE-CHAIN, D-RING, AND A-RING MODIFICATIONS

The synthesis of 12 analogs (138–149) of $1\alpha,25(\text{OH})_2$ -16-ene-20-cyclopropylvitamin D_3 relied on Wittig–Horner coupling⁷⁷ (Figure 14). These compounds have an unsaturated D-ring between C-16 and C-17 and a cyclopropyl group located at C-20. The structural diversity on the side-chain covered triple CC bonds (138–144), double CC bonds (145–147) and single CC bonds (148 and 149) together with the A-ring covered $1\alpha,3\beta$ -dihydroxy (138, 143, 146, and 149), 19-nor (139, 142, 145, and 148), 3-deoxy- 1α -hydroxy (141), and 1α -fluor- 3β -hydroxy (140 and 144). The anti-inflammatory properties of these compounds were studied via analyzing the inhibition of the secretion of the cytokines interferon- γ (IFNG) and tumor necrosis factor (TNF). Most of 16-ene-20-cyclopropyl analogs inhibited IFNG with similar potency to $1,25(\text{OH})_2\text{D}_3$, but compound 149 was more potent. The inhibition of TNF showed wide differences, some analogs (138, 140, 143–145) failed to induce TNF inhibition, whereas analog 149 inhibited TNF more efficiently than $1,25(\text{OH})_2\text{D}_3$. The metabolism of 149 was studied and the stable 24-oxo metabolite 150 accumulated during metabolism. Compound 150 mediates similar induction of primary vitamin D target genes as analog 149 but has a lower calcemic activity.

■ NONSTEROIDAL VDR LIGANDS

The synthesis of nonsteroidal VDR agonists containing a hydrophobic 1,12-dicarba-*closo*-dodecaborane (*p*-carborane) unit was achieved through bimolecular nucleophilic substitution⁷⁸ (Figure 15). The carborane cage replaced the CD-rings of the natural hormone exploiting the hydrophobicity of *p*-carborane. Despite their simple and flexible structure, the carborane-based VDR ligands show moderate binding affinity for VDR compared to $1,25(\text{OH})_2\text{D}_3$. The analogs are flexible acyclic triols; i.e., they lack an A-ring and conjugated triene structures. Their structures shared a branched side-chain on a carborane carbon, and in the other carbon three different chains are bound either to 3-oxaheptan-5,7-diol (151), 3-oxahexan-5,6-diol (152) or 4-oxaheptan-6,7-diol (153). The flexibility of the diol is favorable for VDR binding affinity, which, however, is more than 100 times lower than for $1,25(\text{OH})_2\text{D}_3$. Nevertheless, these mimics are rather active in inducing HL-60 cell differentiation [rac-151, 5%; (S)-151, 8%; (R)-151, 2%; 152, 0.05%; 153, 0.001%]. The *S*-isomers showed, compared to the *R*-enantiomer, higher prodifferentiation activity and VDR binding affinity.

LG190178 (154) is the first published nonsteroidal vitamin D analog.⁷⁹ In general, VDR ligands based on bisphenyl core compounds with γ -hydroxycarboxylic acid moiety (155) show agonist activity. From compounds with a fluorine-containing bisphenyl core⁸⁰ the hexafluoro analog (157) is 5 times more potent in reporter gene assays than the parent compound (155), shows 2 times higher prodifferentiation activity, and is 7

times more effective in inducing bone γ -carboxyglutamate protein (BGLAP) expression. Like in secosteroidal vitamin D analogs, fluorination is an effective modification as shown by crystal structure analysis of the VDR-LBD complexed with 155. Also in this case helix H12 is stabilized in the agonistic position allowing interaction with coactivator proteins. From physicochemical point of view the fluorine atom's ionic radius is 100% larger than that of the hydrogen, the van der Waals radius is only 27% larger. However, the high electronegativity of fluor has a possible "pulling" effect for residues residing in the nearest proximity of a fluorinated functional group. This can be illustrated on the comparison of the aliphatic chain of the ligands that have CH_3 or CF_3 functional groups at carbon C-26 and C-27. The latter shows additional five hydrophobic residues that cannot be seen for CH_3 at the cutoff 3.5 \AA (Figure 13B). The only maintained interactions are with conserved histidine residues. In addition, the possible reaction to the strong van der Waals forces from hydrophobic residues is a moderate opening of the functional groups by 5° with maintained distance between carbon C-26 and C-27 due to additional twist in the absolute position of the C-25 carbon. These net effects show also small variation the LBP size, which is slightly confined in the presence of CF_3 functional groups resulting in a tighter packing of the cavity (Figure 13B).

Nonsteroidal vitamin D mimics with phenylpyrrolyl pentane skeletons have been designed (159–165).^{81,82} Among them, 159 shows clear antiproliferative effects on MCF-7 cells. In order to improve the biological activity of compound 159, derivatives were designed comprising side-chains terminated in a diethylcarbinol, hydrophilic groups or hydrophobic groups (160–165). The antiproliferative activities of the compounds were tested in MCF-7 cells, PC3 human prostate cancer cells, Caco2 human colon cancer cells, and HepG2 human liver cancer cells. Compound 160b exhibits the best antiproliferative activity, being more potent than the prototype compound 159 and $1,25(\text{OH})_2\text{D}_3$. Also the compounds 160a, 160c, 160d, 160f, 160g, 164b, and 165b show in all four model systems better antiproliferative activities than 159 and $1,25(\text{OH})_2\text{D}_3$. The R^2 substitutions at the pyrrole-ring side-chains are crucial for the antiproliferative activity of the compounds. Molecules with hydrophilic groups at the end of the pyrrole-ring side-chain (160a, 160b, 160d, 160g, and 164b) are more potent than those bearing hydrophobic groups (161a, 161b, 162a–d, 163a, and 164b). Moreover, compounds 160a, 160c, 160d, 160g, and 164b were less cytotoxic than 159 and $1,25(\text{OH})_2\text{D}_3$. Compounds 160a–d, 164b, and 165b also display prodifferentiating activity. In reporter gene assays 164b is the most potent compound, whereas the transactivation potential of 160b and 160g is comparable to that of $1,25(\text{OH})_2\text{D}_3$.

A novel class of analogs,⁸³ where the C-ring and D-ring were replaced by an aromatic *m*-phenylene D-ring and an alkyl chain, were synthesized based on the formation of the triene system through a Pd-catalyzed ring-closure of an enol triflate and a subsequent Suzuki–Miyaura reaction with appropriate boronate in aqueous medium.⁸⁴ Compounds 166a–e efficiently induce the differentiation of human keratinocytes and show antiproliferative activity in MCF-7, PC-3, SKOV-3 (human ovary cancer), and HaCaT cells comparable to $1,25(\text{OH})_2\text{D}_3$. Compound 166a with the shortest chain at C-8 is most active not only in antiproliferative tests but also in reporter gene assays. Importantly, none of compounds 166a–e induce hypercalcemia. In a SCID mice xenograph model of aggressive MDA-MB-231 human breast cancer cells compound

166a shows high efficacy for tumor growth inhibition and overall survival.

CONCLUSIONS

This review demonstrated that clever and relevant chemistry significantly increased the number and variety of synthetic vitamin D analogs. Analog design had advanced and led to functional molecules, such as the *o*-carborane compounds, that are devoid of a 25-OH group. Moreover, there are now molecules that completely lack A- and/or CD-rings, such as *p*-carborane compounds, but still interact with VDR. Some of these nonsteroidal vitamin D analogs display high activity in vitro in combination with low calcemic effects in vivo. Thus, the area of nonsteroidal analogs and mimics is expected to further rise in future.

The assessment of the biological profile of VDR ligands is still primarily reduced to in vitro assays, such as VDR binding affinity, reporter gene assays, and antiproliferative and prodifferentiation measurements in different cancer cell lines. The variety in the assays makes a direct comparison of the different types of vitamin D analogs difficult. Moreover, a reliable extrapolation of the in vivo potential of the compounds is impossible without changing to a different set of assays, such as gene expression profiles in freshly isolated human peripheral blood mononuclear cells.⁸⁵

Nowadays research on vitamin D analogs is nearly exclusively performed in academia and many interesting approaches for optimizing the profile of VDR ligands have not been explored to their limits. Accordingly, a complete picture is still missing and there is potential for improvements. The number of nearly 150 solved crystal structures of the VDR-LBD complexed with synthetic ligands is impressive and demonstrates the active interest of academia in understanding the molecular actions of VDR agonists and antagonists.

Unfortunately, failures of clinical trials focused on cancer have majorly dampened the interest of pharma industry in further developing vitamin D compounds. Since the natural hormone $1,25(\text{OH})_2\text{D}_3$ primarily prevents bone- and immune-system-related diseases, the molecule and its synthetic derivatives may not be perfect drugs for the therapy of cancer. Nevertheless, calcipotriol-activated VDR in stroma of human pancreatic tumors had been shown to markedly reduce markers of inflammation and fibrosis in pancreatitis and human tumor stroma.⁸⁶ This suggests that vitamin D compounds rather affect immune cells of the microenvironment of tumors than directly inhibiting the proliferation of the cancer cells. In fact, to date, most genome-wide data on the action of VDR and its ligands are available from cells of the hematopoietic system.⁸⁷ This further emphasizes the impact of vitamin D and VDR for innate and adaptive immunity and suggests that these areas should be further explored for a commercial application.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.jmedchem.9b00208.

Table S1 describing all 143 publically available VDR ligand crystal structures with individual hyperlinks to the PDB and PubMed databases and citations of refs 90–144 (PDF)

AUTHOR INFORMATION

Corresponding Author

*Phone: +358-40-355-3062. E-mail: carsten.carlberg@uef.fi.

ORCID

Carsten Carlberg: 0000-0003-2633-0684

Notes

The authors declare no competing financial interest.

Biographies

Miguel A. Maestro received his Ph.D. in Chemistry from the University of Santiago de Compostela (Spain) in 1989. He did his postdoctoral stay in synthetic organic chemistry at the ETH-Zentrum (Zürich, Switzerland). In 1991 he joined the Faculty of Sciences at the University of A Coruña (Spain) studying synthetic methodologies towards vitamin D metabolites and analogs. Since 2018 he holds a Professor position at the Department of Chemistry. Prof. Maestro's interests are the synthesis of new vitamin D analogs with isotopic labeling and the atomic disposition of molecular structures through X-ray crystallography.

Ferdinand Molnár received his Ph.D. in Biochemistry from the University of Kuopio (Kuopio, Finland) in 2006. He did his postdoctoral training in Structural Biology at the IGBMC (Illkirch, France). In 2008 he joined the School of Pharmacy at the University of Eastern Finland (Kuopio, Finland) studying nuclear receptor–ligand, –protein, and –DNA interactions. In 2018 he moved to the Nazarbayev University (Astana, Kazakhstan) where he holds an Associate Professor position at the Department of Biology. Prof. Molnár's interests are integrative structural biology and bioinformatics, eukaryotic transcriptional regulation in health and disease, and recombinant protein production.

Carsten Carlberg graduated in 1989 with a Ph.D. in Biochemistry at the Free University Berlin (Germany). After positions as postdoc at Roche (Basel, Switzerland), group leader at the University of Geneva (Switzerland), and docent at the University of Düsseldorf (Germany) he is since 2000 Full Professor of Biochemistry at the University of Eastern Finland in Kuopio (Finland). His work focuses on mechanisms of gene regulation by nuclear hormones, in particular on vitamin D. At present Prof. Carlberg has projects on epigenome-wide effects of vitamin D on the human immune system.

ACKNOWLEDGMENTS

M.A.M. thanks Xunta de Galicia (Grant ED431B-2018/GI2105) for financial support. C.C. thanks the Academy of Finland for support.

ABBREVIATIONS USED

$1,25(\text{OH})_2\text{D}_3$, $1\alpha,25$ -dihydroxyvitamin D_3 ; $1,25(\text{OH})_2\text{D}_2$, $1\alpha,25$ -dihydroxyvitamin D_2 ; $25(\text{OH})\text{D}_3$, 25-hydroxyvitamin D_3 ; BGLAP, bone γ -carboxyglutamate protein (previously called osteocalcin); CAMP, cathelicidin antimicrobial peptide; CYP24A1, cytochrome P450, family 24, subfamily A, polypeptide 1; DBP, vitamin D binding protein; IFNG, interferon- γ ; LBD, ligand-binding domain; LBP, ligand-binding pocket; PDB, Protein Data Base; PTH, parathyroid hormone; RXR, retinoid X receptor; TNF, tumor necrosis factor; TSS, transcription start site; VDR, vitamin D receptor

REFERENCES

(1) Tremezaygues, L.; Sticherling, M.; Pfohler, C.; Friedrich, M.; Meineke, V.; Seifert, M.; Tilgen, W.; Reichrath, J. Cutaneous photosynthesis of vitamin D: an evolutionary highly-conserved endocrine system that protects against environmental hazards

including UV-radiation and microbial infections. *Anticancer Res.* **2006**, *26* (4A), 2743–2748.

(2) Jasinghe, V. J.; Perera, C. O.; Barlow, P. J. Bioavailability of vitamin D₂ from irradiated mushrooms: an in vivo study. *Br. J. Nutr.* **2005**, *93* (6), 951–955.

(3) Norman, A. W. From vitamin D to hormone D: fundamentals of the vitamin D endocrine system essential for good health. *Am. J. Clin. Nutr.* **2008**, *88* (2), 491S–499S.

(4) Hollis, B. W. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J. Nutr.* **2005**, *135* (2), 317–322.

(5) Carlberg, C.; Polly, P. Gene regulation by vitamin D₃. *Crit. Rev. Eukaryotic Gene Expression* **1998**, *8* (1), 19–42.

(6) Bouillon, R.; Suda, T. Vitamin D: calcium and bone homeostasis during evolution. *BoneKEY Rep.* **2014**, *3*, 480.

(7) Hewison, M. An update on vitamin D and human immunity. *Clin. Endocrinol. (Oxford, U. K.)* **2012**, *76* (3), 315–325.

(8) Rook, G. A. The role of vitamin D in tuberculosis. *Am. Rev. Respir. Dis.* **1988**, *138* (4), 768–770.

(9) Ramagopalan, S. V.; Maugeri, N. J.; Handunnetthi, L.; Lincoln, M. R.; Orton, S. M.; Dymont, D. A.; DeLuca, G. C.; Herrera, B. M.; Chao, M. J.; Sadovnick, A. D.; Ebers, G. C.; Knight, J. C. Expression of the multiple sclerosis-associated MHC class II Allele HLA-DRB1*1501 is regulated by vitamin D. *PLoS Genet.* **2009**, *5* (2), e1000369.

(10) Holick, M. F.; Binkley, N. C.; Bischoff-Ferrari, H. A.; Gordon, C. M.; Hanley, D. A.; Heaney, R. P.; Murad, M. H.; Weaver, C. M. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J. Clin. Endocrinol. Metab.* **2011**, *96* (7), 1911–1930.

(11) Carlberg, C. The physiology of vitamin D—far more than calcium and bone. *Front. Physiol.* **2014**, *5*, 335.

(12) Holick, M. F. Vitamin D deficiency. *N. Engl. J. Med.* **2007**, *357*, 266–281.

(13) Kupferschmidt, K. Uncertain verdict as vitamin D goes on trial. *Science* **2012**, *337* (6101), 1476–1478.

(14) Institute of Medicine. *Dietary Reference Intakes for Calcium and Vitamin D*; Ross, A. C., Taylor, C. L., Yaktine, A. L., Del Valle, H. B., Eds.; National Academies Press: Washington, DC, 2011.

(15) Cheskis, B. J.; Freedman, L. P.; Nagpal, S. Vitamin D receptor ligands for osteoporosis. *Curr. Opin. Invest. Drugs* **2006**, *7* (10), 906–911.

(16) Fogh, K.; Kragballe, K. New vitamin D analogs in psoriasis. *Curr. Drug Targets: Inflammation Allergy* **2004**, *3* (2), 199–204.

(17) Bouillon, R.; Okamura, W. H.; Norman, A. W. Structure-function relationships in the vitamin D endocrine system. *Endocr. Rev.* **1995**, *16*, 200–257.

(18) Carlberg, C.; Mouriño, A. New vitamin D receptor ligands. *Expert Opin. Ther. Pat.* **2003**, *13*, 761–772.

(19) Carlberg, C.; Molnár, F.; Mouriño, A. Vitamin D receptor ligands: the impact of crystal structures. *Expert Opin. Ther. Pat.* **2012**, *22* (4), 417–435.

(20) Evans, T. R.; Colston, K. W.; Lofts, F. J.; Cunningham, D.; Anthoney, D. A.; Gogas, H.; de Bono, J. S.; Hamberg, K. J.; Skov, T.; Mansi, J. L. A phase II trial of the vitamin D analog Seocalcitol (EB1089) in patients with inoperable pancreatic cancer. *Br. J. Cancer* **2002**, *86* (5), 680–685.

(21) Heikkinen, S.; Väisänen, S.; Pehkonen, P.; Seuter, S.; Benes, V.; Carlberg, C. Nuclear hormone 1 α ,25-dihydroxyvitamin D₃ elicits a genome-wide shift in the locations of VDR chromatin occupancy. *Nucleic Acids Res.* **2011**, *39* (21), 9181–9193.

(22) Gombart, A. F.; Borregaard, N.; Koeffler, H. P. Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D₃. *FASEB J.* **2005**, *19* (9), 1067–1077.

(23) Haussler, M. R.; Haussler, C. A.; Jurutka, P. W.; Thompson, P. D.; Hsieh, J. C.; Remus, L. S.; Selznick, S. H.; Whitfield, G. K. The

vitamin D hormone and its nuclear receptor: molecular actions and disease states. *J. Endocrinol.* **1997**, *154* (Suppl.), S57–S73.

(24) Carlberg, C. Vitamin D genomics: from in vitro to in vivo. *Front. Endocrinol.* **2018**, *9*, 250.

(25) Verstuyf, A.; Carmeliet, G.; Bouillon, R.; Mathieu, C. Vitamin D: a pleiotropic hormone. *Kidney Int.* **2010**, *78* (2), 140–145.

(26) Wang, Y.; Zhu, J.; DeLuca, H. F. Where is the vitamin D receptor? *Arch. Biochem. Biophys.* **2012**, *523* (1), 123–133.

(27) Carlberg, C.; Molnár, F. Vitamin D receptor signaling and its therapeutic implications: genome-wide and structural view. *Can. J. Physiol. Pharmacol.* **2015**, *93*, 311–318.

(28) Rochel, N.; Wurtz, J. M.; Mitschler, A.; Klaholz, B.; Moras, D. Crystal structure of the nuclear receptor for vitamin D bound to its natural ligand. *Mol. Cell* **2000**, *5*, 173–179.

(29) Molnár, F. Structural considerations of vitamin D signaling. *Front. Physiol.* **2014**, *5*, 191.

(30) Molnár, F.; Peräkylä, M.; Carlberg, C. Vitamin D receptor agonists specifically modulate the volume of the ligand-binding pocket. *J. Biol. Chem.* **2006**, *281* (15), 10516–10526.

(31) Carlberg, C. Molecular basis of the selective activity of vitamin D analogues. *J. Cell. Biochem.* **2003**, *88* (2), 274–281.

(32) Carlberg, C. Molecular endocrinology of vitamin D on the epigenome level. *Mol. Cell. Endocrinol.* **2017**, *453*, 14–21.

(33) Wei, Z.; Yoshihara, E.; He, N.; Hah, N.; Fan, W.; Pinto, A. F. M.; Huddy, T.; Wang, Y.; Ross, B.; Estepa, G.; Dai, Y.; Ding, N.; Sherman, M. H.; Fang, S.; Zhao, X.; Liddle, C.; Atkins, A. R.; Yu, R. T.; Downes, M.; Evans, R. M. Vitamin D switches BAF complexes to protect beta cells. *Cell* **2018**, *173* (5), 1135–1149.

(34) Carlberg, C.; Campbell, M. J. Vitamin D receptor signaling mechanisms: Integrated actions of a well-defined transcription factor. *Steroids* **2013**, *78* (2), 127–136.

(35) Carlberg, C.; Molnár, F. Current status of vitamin D signaling and its therapeutic applications. *Curr. Top. Med. Chem.* **2012**, *12* (6), 528–547.

(36) Pérez-García, X.; Rumbo, A.; Larriba, M. J.; Ordóñez, P.; Muñoz, A.; Mouriño, A. The first locked side-chain analogues of calcitriol (1 α ,25-dihydroxyvitamin D₃) induce vitamin D receptor transcriptional activity. *Org. Lett.* **2003**, *5*, 4033–4036.

(37) Siqueiro, R.; Maestro, M. A.; Mouriño, A. Synthesis of side-chain locked analogs of 1 α ,25-dihydroxyvitamin D₃ bearing a C17 methyl group. *Org. Lett.* **2018**, *20* (9), 2641–2644.

(38) Norman, A. W.; Manchand, P. S.; Uskokovic, M. R.; Okamura, W. H.; Takeuchi, J. A.; Bishop, J. E.; Hisatake, J.-I.; Koeffler, H. P.; Peleg, S. Characterization of a novel analog of 1 α ,25(OH)₂-vitamin D₃ with two side chains: interaction with its nuclear receptor and cellular actions. *J. Med. Chem.* **2000**, *43*, 2719–2730.

(39) Herdick, M.; Bury, Y.; Quack, M.; Uskokovic, M. R.; Polly, P.; Carlberg, C. Response element and coactivator-mediated conformational change of the vitamin D₃ receptor permits sensitive interaction with agonists. *Mol. Pharmacol.* **2000**, *57* (6), 1206–1217.

(40) Väisänen, S.; Peräkylä, M.; Kärkkäinen, J. I.; Uskokovic, M. R.; Carlberg, C. Structural evaluation of the agonistic activity of a vitamin D analog with two side chains binding to the nuclear vitamin D receptor. *Mol. Pharmacol.* **2003**, *63* (6), 1230–1237.

(41) Liu, Y. Y.; Collins, E. D.; Norman, A. W.; Peleg, S. Differential interaction of 1 α ,25-dihydroxyvitamin D₃ analogues and their 20-epi homologues with the vitamin D receptor. *J. Biol. Chem.* **1997**, *272* (6), 3336–3345.

(42) Maehr, H.; Lee, H. J.; Perry, B.; Suh, N.; Uskokovic, M. R. Calcitriol derivatives with two different side chains at C-20. V. Potent inhibitors of mammary carcinogenesis and inducers of leukemia differentiation. *J. Med. Chem.* **2009**, *52* (17), 5505–5519.

(43) Okamoto, R.; Gery, S.; Kuwayama, Y.; Borregaard, N.; Ho, Q.; Alvarez, R.; Akagi, T.; Liu, G. Y.; Uskokovic, M. R.; Koeffler, H. P. Novel Gemini vitamin D₃ analogs: large structure/function analysis and ability to induce antimicrobial peptide. *Int. J. Cancer* **2014**, *134* (1), 207–217.

(44) Otero, R.; Seoane, S.; Siqueiro, R.; Belorusova, A. Y.; Maestro, M. A.; Perez-Fernandez, R.; Rochel, N.; Mouriño, A. Carborane-based

design of a potent vitamin D receptor agonist. *Chem. Sci.* **2016**, *7*, 1033–1037.

(45) Takahashi, E.; Nakagawa, K.; Suhara, Y.; Kittaka, A.; Nihei, K.; Konno, K.; Takayama, H.; Ozono, K.; Okano, T. Biological activities of 2 α -substituted analogues of 1 α ,25-dihydroxyvitamin D₃ in transcriptional regulation and human promyelocytic leukemia (HL-60) cell proliferation and differentiation. *Biol. Pharm. Bull.* **2006**, *29* (11), 2246–2250.

(46) Honzawa, S.; Takahashi, N.; Yamashita, A.; Sugiura, T.; Kurihara, M.; Arai, M. A.; Kato, S.; Kittaka, A. Synthesis of a 1 α -C-methyl analogue of 25-hydroxyvitamin D₃: interaction with a mutant vitamin D receptor Arg274Leu. *Tetrahedron* **2009**, *65* (34), 7135–7145.

(47) Sawada, D.; Tsukuda, Y.; Yasuda, K.; Sakaki, T.; Saito, H.; Takagi, K.; Takenouchi, K.; Chen, T. C.; Reddy, G. S.; Kittaka, A. Synthesis and biological activities of 1 α ,4 α ,25- and 1 α ,4 β ,25-trihydroxyvitamin D₃ and their metabolism by human CYP24A1 and UDP-glucuronosyltransferase. *Chem. Pharm. Bull.* **2012**, *60* (10), 1343–1346.

(48) Sibilska, I. K.; Szybinski, M.; Sicinski, R. R.; Plum, L. A.; DeLuca, H. F. Highly potent 2-methylene analogs of 1 α ,25-dihydroxyvitamin D₃: synthesis and biological evaluation. *J. Steroid Biochem. Mol. Biol.* **2013**, *136*, 9–13.

(49) Piotrowska, A.; Wierzbicka, J.; Nadkarni, S.; Brown, G.; Kutner, A.; Zmijewski, M. A. Antiproliferative activity of double point modified analogs of 1,25-dihydroxyvitamin D₂ against human malignant melanoma cell lines. *Int. J. Mol. Sci.* **2016**, *17* (1), E76.

(50) Corcoran, A.; Nadkarni, S.; Yasuda, K.; Sakaki, T.; Brown, G.; Kutner, A.; Marcinkowska, E. Biological evaluation of double point modified analogues of 1,25-dihydroxyvitamin D₂ as potential anti-leukemic agents. *Int. J. Mol. Sci.* **2016**, *17* (2), E91.

(51) Saito, N.; Kittaka, A. Highly potent vitamin D receptor antagonists: design, synthesis, and biological evaluation. *ChemBioChem* **2006**, *7* (10), 1478–1490.

(52) Saito, N.; Matsunaga, T.; Saito, H.; Anzai, M.; Takenouchi, K.; Miura, D.; Namekawa, J.; Ishizuka, S.; Kittaka, A. Further synthetic and biological studies on vitamin D hormone antagonists based on C2 α -alkylation and C2 α -functionalization of 25-dehydro-1 α -hydroxyvitamin D₃-26,23-lactones. *J. Med. Chem.* **2006**, *49* (24), 7063–7075.

(53) Sakamaki, Y.; Inaba, Y.; Yoshimoto, N.; Yamamoto, K. Potent antagonist for the vitamin D receptor: vitamin D analogues with simple side chain structure. *J. Med. Chem.* **2010**, *53* (15), 5813–5826.

(54) Yoshimoto, N.; Sakamaki, Y.; Haeta, M.; Kato, A.; Inaba, Y.; Itoh, T.; Nakabayashi, M.; Ito, N.; Yamamoto, K. Butyl pocket formation in the vitamin D receptor strongly affects the agonistic or antagonistic behavior of ligands. *J. Med. Chem.* **2012**, *55* (9), 4373–4381.

(55) Yoshino, M.; Eto, K.; Takahashi, K.; Ishihara, J.; Hatakeyama, S.; Ono, Y.; Saito, H.; Kubodera, N. Synthesis of 20-eldecalcitol [20-epi-1 α ,25-dihydroxy-2 β -(3-hydroxypropoxy)vitamin D₃: 20-epi-ED-71]. *Heterocycles* **2010**, *81*, 381–394.

(56) Binderup, L.; Latini, S.; Binderup, E.; Bretting, C.; Calverley, M.; Hansen, K. 20-epi-vitamin D₃ analogues: a novel class of potent regulators of cell growth and immune responses. *Biochem. Pharmacol.* **1991**, *42*, 1569–1575.

(57) Paaren, H. F.; Schones, H. K.; DeLuca, H. F. Synthesis of 1 β -hydroxyvitamin D₃ and 1 β ,25-dihydroxyvitamin D₃. *J. Chem. Soc., Chem. Commun.* **1977**, 890–892.

(58) Napoli, J. L.; Fivizzani, M. A.; Schones, H. K.; DeLuca, H. F. 1-Fluorovitamin D₃, a vitamin D₃ analogue more active on bone-calcium mobilization than on intestinal-calcium transport. *Biochemistry* **1979**, *18* (9), 1641–1646.

(59) Paaren, H. E.; Fivizzani, M. A.; Schones, H. K.; De Luca, H. F. 1 α ,25-difluorovitamin D₃: an inert vitamin D analog. *Arch. Biochem. Biophys.* **1981**, *209* (2), 579–583.

(60) Glebocka, A.; Chiellini, G. A-ring analogs of 1,25-dihydroxyvitamin D₃. *Arch. Biochem. Biophys.* **2012**, *523* (1), 48–57.

(61) Ishida, H.; Shimizu, M.; Yamamoto, K.; Iwasaki, Y.; Yamada, S.; Yamaguchi, K. Synthesis of 1-alkyl-1,25-dihydroxyvitamin D₃. *J. Org. Chem.* **1995**, *60*, 1828–1833.

(62) Sekimoto, H.; Siu-Caldera, M. L.; Weiskopf, A.; Vouros, P.; Muralidharan, K. R.; Okamura, W. H.; Uskokovic, M. R.; Reddy, G. S. 1 α ,25-dihydroxy-3-epi-vitamin D₃: *in vivo* metabolite of 1 α ,25-dihydroxyvitamin D₃ in rats. *FEBS Lett.* **1999**, *448* (2–3), 278–282.

(63) Nakagawa, K.; Kurobe, M.; Ozono, K.; Konno, K.; Fujishima, T.; Takayama, H.; Okano, T. Novel ring A stereoisomers of 2-methyl-1 α ,25-dihydroxyvitamin D₃ and 2-methyl-20-epi-1 α ,25-dihydroxyvitamin D₃: transactivation of target genes and modulation of differentiation in human promyelocytic leukemia (HL-60) cells. *Biochem. Pharmacol.* **2000**, *59* (6), 691–702.

(64) Suhara, Y.; Nihei, K. I.; Tanigawa, H.; Fujishima, T.; Konno, K.; Nakagawa, K.; Okano, T.; Takayama, H. Syntheses and biological evaluation of novel 2 α -substituted 1 α ,25-dihydroxyvitamin D₃ analogues. *Bioorg. Med. Chem. Lett.* **2000**, *10* (10), 1129–1132.

(65) Perlman, K. L.; Swenson, R. E.; Paaren, H. E.; Schones, H. K.; DeLuca, H. F. Novel synthesis of 19-nor-vitamin D compounds. *Tetrahedron Lett.* **1991**, *32*, 7663–7666.

(66) Bouillon, R.; Sarandeses, L. A.; Allewaert, K.; Zhao, J.; Mascarenas, J. L.; Mouriño, A.; Vrielynck, S.; de Clercq, P.; Vandewalle, M. Biologic activity of dihydroxylated 19-nor-(pre)-vitamin D₃. *J. Bone Miner. Res.* **1993**, *8* (8), 1009–1015.

(67) Sicinski, R. R.; Perlman, K. L.; DeLuca, H. F. Synthesis and biological activity of 2-hydroxy and 2-alkoxy analogs of 1 α ,25-dihydroxy-19-norvitamin D₃. *J. Med. Chem.* **1994**, *37* (22), 3730–3738.

(68) Yamamoto, H.; Shevde, N. K.; Warriar, A.; Plum, L. A.; DeLuca, H. F.; Pike, J. W. 2-Methylene-19-nor-(20S)-1,25-dihydroxyvitamin D₃ potentially stimulates gene-specific DNA binding of the vitamin D receptor in osteoblasts. *J. Biol. Chem.* **2003**, *278* (34), 31756–31765.

(69) Williams, S.; Bledsoe, R. K.; Collins, J. L.; Boggs, S.; Lambert, M. H.; Miller, A. B.; Moore, J.; McKee, D. D.; Moore, L.; Nichols, J.; Parks, D.; Watson, M.; Wisely, B.; Willson, T. M. X-ray crystal structure of the liver X receptor beta ligand binding domain: regulation by a histidine-tryptophan switch. *J. Biol. Chem.* **2003**, *278* (29), 27138–27143.

(70) Tsugawa, N.; Nakagawa, K.; Kurobe, M.; Ono, Y.; Kubodera, N.; Ozono, K.; Okano, T. *In vitro* biological activities of a series of 2 beta-substituted analogues of 1 α ,25-dihydroxyvitamin D₃. *Biol. Pharm. Bull.* **2000**, *23* (1), 66–71.

(71) Shimizu, M.; Miyamoto, Y.; Kobayashi, E.; Shimazaki, M.; Yamamoto, K.; Reischl, W.; Yamada, S. Synthesis and biological activities of new 1 α ,25-dihydroxy-19norvitamin D₃ analogs with modifications in both A-ring and the side chain. *Bioorg. Med. Chem.* **2006**, *14*, 4277–4294.

(72) Posner, G. H.; Lee, J. K.; Wang, Q.; Peleg, S.; Burke, M.; Brem, H.; Dolan, P.; Kensler, T. W. Noncalcemic, antiproliferative, transcriptionally active, 24-fluorinated hybrid analogues of the hormone 1 α ,25-dihydroxyvitamin D₃: synthesis and preliminary biological evaluation. *J. Med. Chem.* **1998**, *41*, 3008–3014.

(73) Peleg, S.; Ismail, A.; Uskokovic, M. R.; Avnur, Z. Evidence for tissue- and cell-type selective activation of the vitamin D receptor by Ro-26-9228, a noncalcemic analog of vitamin D₃. *J. Cell. Biochem.* **2003**, *88* (2), 267–273.

(74) Flores, A.; Sicinski, R. R.; Grzywacz, P.; Thoden, J. B.; Plum, L. A.; Clagett-Dame, M.; DeLuca, H. F. A 20S combined with a 22R configuration markedly increases both *in vivo* and *in vitro* biological activity of 1 α ,25-dihydroxy-22-methyl-2-methylene-19-norvitamin D₃. *J. Med. Chem.* **2012**, *55* (9), 4352–4366.

(75) Sibilska, I. K.; Szybinski, M.; Sicinski, R. R.; Plum, L. A.; DeLuca, H. F. Synthesis and biological activity of 2-methylene analogues of calcitriol and related compounds. *J. Med. Chem.* **2015**, *58* (24), 9653–9662.

(76) Sawada, D.; Tsukuda, Y.; Saito, H.; Kakuda, S.; Takimoto-Kamimura, M.; Ochiai, E.; Takenouchi, K.; Kittaka, A. Development of 14-epi-19-nortachysterol and its unprecedented binding config-

uration for the human vitamin D receptor. *J. Am. Chem. Soc.* **2011**, *133*, 7215–7221.

(77) Laverny, G.; Penna, G.; Uskokovic, M.; Marczak, S.; Maehr, H.; Jankowski, P.; Ceailles, C.; Vouros, P.; Smith, B.; Robinson, M.; Reddy, G. S.; Adorini, L. Synthesis and anti-inflammatory properties of $1\alpha,25$ -dihydroxy-16-ene-20-cyclopropyl-24-oxo-vitamin D_3 , a hypocalcemic, stable metabolite of $1\alpha,25$ -dihydroxy-16-ene-20-cyclopropyl-vitamin D_3 . *J. Med. Chem.* **2009**, *52* (8), 2204–2213.

(78) Fujii, S.; Masuno, H.; Taoda, Y.; Kano, A.; Wongmayura, A.; Nakabayashi, M.; Ito, N.; Shimizu, M.; Kawachi, E.; Hirano, T.; Endo, Y.; Tanatani, A.; Kagechika, H. Boron cluster-based development of potent nonsecosteroidal vitamin D receptor ligands: direct observation of hydrophobic interaction between protein surface and carborane. *J. Am. Chem. Soc.* **2011**, *133* (51), 20933–20941.

(79) Boehm, M. F.; Fitzgerald, P.; Zou, A.; Elgort, M. G.; Bischoff, E. D.; Mere, L.; Mais, D. E.; Bissonnette, R. P.; Heyman, R. A.; Nadzan, A. M.; Reichman, M.; Allegretto, E. A. Novel nonsecosteroidal vitamin D mimics exert VDR-modulating activities with less calcium mobilization than $1,25$ -dihydroxyvitamin D_3 . *Chem. Biol.* **1999**, *6* (5), 265–275.

(80) Kashiwagi, H.; Ohta, M.; Ono, Y.; Morikami, K.; Itoh, S.; Sato, H.; Takahashi, T. Effects of fluorines on nonsecosteroidal vitamin D receptor agonists. *Bioorg. Med. Chem.* **2013**, *21* (3), 712–721.

(81) Hao, M.; Hou, S.; Xue, L.; Yuan, H.; Zhu, L.; Wang, C.; Wang, B.; Tang, C.; Zhang, C. Further developments of the phenyl-pyrrolyl pentane series of nonsteroidal vitamin D receptor modulators as anticancer agents. *J. Med. Chem.* **2018**, *61* (7), 3059–3075.

(82) Shen, W.; Xue, J.; Zhao, Z.; Zhang, C. Novel nonsecosteroidal VDR agonists with phenyl-pyrrolyl pentane skeleton. *Eur. J. Med. Chem.* **2013**, *69*, 768–778.

(83) Gogoi, P.; Seoane, S.; Sigüeiro, R.; Guiberteau, T.; Maestro, M. A.; Perez-Fernandez, R.; Rochel, N.; Mouriño, A. Aromatic-based design of highly active and noncalcemic vitamin D receptor agonists. *J. Med. Chem.* **2018**, *61* (11), 4928–4937.

(84) Gogoi, P.; Sigüeiro, R.; Eduardo, S.; Mouriño, A. An expeditious route to $1\alpha,25$ -dihydroxyvitamin D_3 and its analogues by an aqueous tandem palladium-catalyzed A-ring closure and Suzuki coupling to the C/D unit. *Chem. - Eur. J.* **2010**, *16* (5), 1432–1435.

(85) Stio, M.; Martinesi, M.; Bruni, S.; Treves, C.; Mathieu, C.; Verstuyf, A.; d'Albasio, G.; Bagnoli, S.; Bonanomi, A. G. The vitamin D analogue TX 527 blocks NF- κ B activation in peripheral blood mononuclear cells of patients with Crohn's disease. *J. Steroid Biochem. Mol. Biol.* **2007**, *103* (1), 51–60.

(86) Sherman, M. H.; Yu, R. T.; Engle, D. D.; Ding, N.; Atkins, A. R.; Tiriach, H.; Collisson, E. A.; Connor, F.; Van Dyke, T.; Kozlov, S.; Martin, P.; Tseng, T. W.; Dawson, D. W.; Donahue, T. R.; Masamune, A.; Shimosegawa, T.; Apte, M. V.; Wilson, J. S.; Ng, B.; Lau, S. L.; Gunton, J. E.; Wahl, G. M.; Hunter, T.; Drebin, J. A.; O'Dwyer, P. J.; Liddle, C.; Tuveson, D. A.; Downes, M.; Evans, R. M. Vitamin D receptor-mediated stromal reprogramming suppresses pancreatitis and enhances pancreatic cancer therapy. *Cell* **2014**, *159* (1), 80–93.

(87) Carlberg, C. Genome-wide (over)view on the actions of vitamin D. *Front. Physiol.* **2014**, *5*, 167.

(88) Plum, L. A.; DeLuca, H. F. Vitamin D, disease and therapeutic opportunities. *Nat. Rev. Drug Discovery* **2010**, *9* (12), 941–955.

(89) Leyssens, C.; Verlinden, L.; Verstuyf, A. The future of vitamin D analogs. *Front. Physiol.* **2014**, *5*, 122.

(90) Tocchini-Valentini, G.; Rochel, N.; Wurtz, J. M.; Mitschler, A.; Moras, D. Crystal structures of the vitamin D receptor complexed to superagonist 20-epi ligands. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98* (10), 5491–5496.

(91) Tocchini-Valentini, G.; Rochel, N.; Wurtz, J. M.; Moras, D. Crystal structures of the vitamin D nuclear receptor liganded with the vitamin D side chain analogues calcipotriol and seocalcitol, receptor agonists of clinical importance: insights into a structural basis for the switching of calcipotriol to a receptor antagonist by further side chain modification. *J. Med. Chem.* **2004**, *47* (8), 1956–1961.

(92) Eelen, G.; Verlinden, L.; Rochel, N.; Claessens, F.; De Clercq, P.; Vandewalle, M.; Tocchini-Valentini, G.; Moras, D.; Bouillon, R.; Verstuyf, A. Superagonistic action of 14-epi-analogs of $1,25$ -dihydroxyvitamin D explained by vitamin D receptor-coactivator interaction. *Mol. Pharmacol.* **2005**, *67* (5), 1566–1573.

(93) Hourai, S.; Fujishima, T.; Kittaka, A.; Suhara, Y.; Takayama, H.; Rochel, N.; Moras, D. Probing a water channel near the A-ring of receptor-bound $1\alpha,25$ -dihydroxyvitamin D_3 with selected 2α -substituted analogues. *J. Med. Chem.* **2006**, *49*, 5199–5205.

(94) Kakuda, S.; Ishizuka, S.; Eguchi, H.; Mizwicki, M. T.; Norman, A. W.; Takimoto-Kamimura, M. Structural basis of the histidine-mediated vitamin D receptor agonistic and antagonistic mechanisms of (23S)- 25 -dehydro- 1α -hydroxyvitamin D_3 -26,23-lactone. *Acta Crystallogr., Sect. D: Biol. Crystallogr.* **2010**, *66* (Part 8), 918–926.

(95) Antony, P.; Sigüeiro, R.; Huet, T.; Sato, Y.; Ramalanjaona, N.; Rodrigues, L. C.; Mouriño, A.; Moras, D.; Rochel, N. Structure-function relationships and crystal structures of the vitamin D receptor bound 2α -methyl-(20S,23S)- and 2α -methyl-(20S,23R)-epoxymethano- $1\alpha,25$ -dihydroxyvitamin D_3 . *J. Med. Chem.* **2010**, *53* (3), 1159–1171.

(96) Molnár, F.; Sigüeiro, R.; Sato, Y.; Araujo, C.; Schuster, I.; Antony, P.; Peluso, J.; Muller, C.; Mouriño, A.; Moras, D.; Rochel, N. $1\alpha,25(\text{OH})_2$ -3-epi-vitamin D_3 , a natural physiological metabolite of vitamin D_3 : its synthesis, biological activity and crystal structure with its receptor. *PLoS One* **2011**, *6*, e18124.

(97) Shindo, K.; Kumagai, G.; Takano, M.; Sawada, D.; Saito, N.; Saito, H.; Kakuda, S.; Takagi, K.; Ochiai, E.; Horie, K.; Takimoto-Kamimura, M.; Ishizuka, S.; Takenouchi, K.; Kittaka, A. New C15-substituted active vitamin D_3 . *Org. Lett.* **2011**, *13*, 2852–2855.

(98) Kashiwagi, H.; Ono, Y.; Shimizu, K.; Haneishi, T.; Ito, S.; Iijima, S.; Kobayashi, T.; Ichikawa, F.; Harada, S.; Sato, H.; Sekiguchi, N.; Ishigai, M.; Takahashi, T. Novel nonsecosteroidal vitamin D_3 carboxylic acid analogs for osteoporosis, and SAR analysis. *Bioorg. Med. Chem.* **2011**, *19* (16), 4721–4729.

(99) Hourai, S.; Rodrigues, L.; Antony, P.; Reina-San-Martin, B.; Ciesielski, F.; Magnier, B.; Schoonjans, K.; Mouriño, A.; Rochel, N.; Moras, D. Structure-based design of a superagonist ligand for the vitamin D nuclear receptor. *Chem. Biol.* **2008**, *15*, 383–392.

(100) Rochel, N.; Moras, D. Crystal structure of a vitamin D_3 analog, ZK203278, showing dissociated profile. *Anticancer Res.* **2012**, *32* (1), 335–339.

(101) Rochel, N.; Hourai, S.; Moras, D. Crystal structure of hereditary vitamin D-resistant rickets-associated mutant H305Q of vitamin D nuclear receptor bound to its natural ligand. *J. Steroid Biochem. Mol. Biol.* **2010**, *121*, 84–87.

(102) Verlinden, L.; Verstuyf, A.; Eelen, G.; Bouillon, R.; Ordóñez-Morán, P.; Larriba, M. J.; Muñoz, A.; Rochel, N.; Sato, Y.; Moras, D.; Maestro, M.; Seoane, S.; Dominguez, F.; Eduardo-Canosa, S.; Nicoletti, D.; Moman, E.; Mouriño, A. Synthesis, structure, and biological activity of des-side chain analogues of $1\alpha,25$ -dihydroxyvitamin D_3 with substituents at C18. *ChemMedChem* **2011**, *6* (5), 788–793.

(103) Fraga, R.; Zacconi, F.; Sussman, F.; Ordóñez-Moran, P.; Muñoz, A.; Huet, T.; Molnár, F.; Moras, D.; Rochel, N.; Maestro, M.; Mouriño, A. Design, synthesis, evaluation, and structure of vitamin D analogues with furan side chains. *Chem. - Eur. J.* **2012**, *18* (2), 603–612.

(104) Saitoh, H.; Chida, T.; Takagi, K.; Horie, K.; Sawai, Y.; Nakamura, Y.; Harada, Y.; Takenouchi, K.; Kittaka, A. Synthesis of C-2 substituted vitamin D derivatives having ringed side chains and their biological evaluation, especially biological effect on bone by modification at the C-2 position. *Org. Biomol. Chem.* **2011**, *9* (10), 3954–3964.

(105) Kashiwagi, H.; Ono, Y.; Ohta, M.; Itoh, S.; Ichikawa, F.; Harada, S.; Takeda, S.; Sekiguchi, N.; Ishigai, M.; Takahashi, T. A series of nonsecosteroidal vitamin D receptor agonists for osteoporosis therapy. *Bioorg. Med. Chem.* **2013**, *21* (7), 1823–1833.

(106) Saitoh, H.; Watanabe, H.; Kakuda, S.; Takimoto-Kamimura, M.; Takagi, K.; Takeuchi, A.; Takenouchi, K. Synthesis and biological

activities of vitamin D₃ derivatives with cyanoalkyl side chain at C-2 position. *J. Steroid Biochem. Mol. Biol.* **2015**, *148*, 27–30.

(107) Sawada, D.; Kakuda, S.; Kamimura-Takimoto, M.; Takeuchi, A.; Matsumoto, Y.; Kittaka, A. Revisiting the 7,8-cis-vitamin D₃ derivatives: Synthesis, evaluating the biological activity, and study of the binding configuration. *Tetrahedron* **2016**, *72*, 2838–2848.

(108) Ciesielski, F.; Sato, Y.; Chebaro, Y.; Moras, D.; Dejaegere, A.; Rochel, N. Structural basis for the accommodation of bis- and tris-aromatic derivatives in vitamin D nuclear receptor. *J. Med. Chem.* **2012**, *55* (19), 8440–8449.

(109) Matsuo, M.; Hasegawa, A.; Takano, M.; Saito, H.; Kakuda, S.; Chida, T.; Takagi, K.; Ochiai, E.; Horie, K.; Harada, Y.; Takimoto-Kamimura, M.; Takenouchi, K.; Sawada, D.; Kittaka, A. Synthesis of 2 α -heteroarylalkyl active vitamin D₃ with therapeutic effect on enhancing bone mineral density in vivo. *ACS Med. Chem. Lett.* **2013**, *4* (7), 671–674.

(110) Belorusova, A. Y.; Eberhardt, J.; Potier, N.; Stote, R. H.; Dejaegere, A.; Rochel, N. Structural insights into the molecular mechanism of vitamin D receptor activation by lithocholic acid involving a new mode of ligand recognition. *J. Med. Chem.* **2014**, *57* (11), 4710–4719.

(111) Zheng, J.; Chang, M. R.; Stites, R. E.; Wang, Y.; Bruning, J. B.; Pascal, B. D.; Novick, S. J.; Garcia-Ordenez, R. D.; Stayrook, K. R.; Chalmers, M. J.; Dodge, J. A.; Griffin, P. R. HDX reveals the conformational dynamics of DNA sequence specific VDR co-activator interactions. *Nat. Commun.* **2017**, *8* (1), 923.

(112) Sawada, D.; Kakuda, S.; Takeuchi, A.; Kawagoe, F.; Takimoto-Kamimura, M.; Kittaka, A. Effects of 2-substitution on 14-epi-19-nortachysterol-mediated biological events: based on synthesis and X-ray co-crystallographic analysis with the human vitamin D receptor. *Org. Biomol. Chem.* **2018**, *16* (14), 2448–2455.

(113) Vanhooke, J. L.; Benning, M. M.; Bauer, C. B.; Pike, J. W.; DeLuca, H. F. Molecular structure of the rat vitamin D receptor ligand binding domain complexed with 2-carbon-substituted vitamin D₃ hormone analogues and a LXXLL-containing coactivator peptide. *Biochemistry* **2004**, *43* (14), 4101–4110.

(114) Vanhooke, J.; Tadi, B.; Benning, M.; Plum, L.; DeLuca, H. New analogs of 2-methylene-19-nor-(20S)-1,25-dihydroxyvitamin D₃ with conformationally restricted side chains: evaluation of biological activity and structural determination of VDR-bound conformations. *Arch. Biochem. Biophys.* **2007**, *460*, 161–165.

(115) Kakuda, S.; Okada, K.; Eguchi, H.; Takenouchi, K.; Hakamata, W.; Kurihara, M.; Takimoto-Kamimura, M. Structure of the ligand-binding domain of rat VDR in complex with the nonsecosteroidal vitamin D₃ analogue YR301. *Acta Crystallogr., Sect. F: Struct. Biol. Cryst. Commun.* **2008**, *64*, 970–973.

(116) Shimizu, M.; Miyamoto, Y.; Takaku, H.; Matsuo, M.; Nakabayashi, M.; Masuno, H.; Udagawa, N.; DeLuca, H.; Ikura, T.; Ito, N. 2-Substituted-16-ene-22-thia-1 α ,25-dihydroxy-26,27-dimethyl-19-norvitamin D₃ analogs: Synthesis, biological evaluation, and crystal structure. *Bioorg. Med. Chem.* **2008**, *16*, 6949–6964.

(117) Nakabayashi, M.; Yamada, S.; Yoshimoto, N.; Tanaka, T.; Igarashi, M.; Ikura, T.; Ito, N.; Makishima, M.; Tokiwa, H.; DeLuca, H.; Shimizu, M. Crystal structures of rat vitamin D receptor bound to adamantyl vitamin D analogs: structural basis for vitamin D receptor antagonism and partial agonism. *J. Med. Chem.* **2008**, *51*, 5320–5329.

(118) Inaba, Y.; Yoshimoto, N.; Sakamaki, Y.; Nakabayashi, M.; Ikura, T.; Tamamura, H.; Ito, N.; Shimizu, M.; Yamamoto, K. A new class of vitamin D analogues that induce structural rearrangement of the ligand-binding pocket of the receptor. *J. Med. Chem.* **2009**, *52*, 1438–1449.

(119) Inaba, Y.; Nakabayashi, M.; Itoh, T.; Yoshimoto, N.; Ikura, T.; Ito, N.; Shimizu, M.; Yamamoto, K. 22S-butyl-1 α ,24R-dihydroxyvitamin D₃: Recovery of vitamin D receptor agonistic activity. *J. Steroid Biochem. Mol. Biol.* **2010**, *121* (1–2), 146–150.

(120) Demizu, Y.; Takahashi, T.; Kaneko, F.; Sato, Y.; Okuda, H.; Ochiai, E.; Horie, K.; Takagi, K.; Kakuda, S.; Takimoto-Kamimura, M.; Kurihara, M. Design, synthesis and X-ray crystallographic study of

new nonsecosteroidal vitamin D receptor ligands. *Bioorg. Med. Chem. Lett.* **2011**, *21* (20), 6104–6107.

(121) Nakabayashi, M.; Tsukahara, Y.; Iwasaki-Miyamoto, Y.; Mihori-Shimazaki, M.; Yamada, S.; Inaba, S.; Oda, M.; Shimizu, M.; Makishima, M.; Tokiwa, H.; Ikura, T.; Ito, N. Crystal structures of hereditary vitamin D-resistant rickets-associated vitamin D receptor mutants R270L and W282R bound to 1,25-dihydroxyvitamin D₃ and synthetic ligands. *J. Med. Chem.* **2013**, *56* (17), 6745–6760.

(122) Kudo, T.; Ishizawa, M.; Maekawa, K.; Nakabayashi, M.; Watarai, Y.; Uchida, H.; Tokiwa, H.; Ikura, T.; Ito, N.; Makishima, M.; Yamada, S. Combination of triple bond and adamantane ring on the vitamin D side chain produced partial agonists for vitamin D receptor. *J. Med. Chem.* **2014**, *57* (10), 4073–4087.

(123) Asano, L.; Ito, I.; Kuwabara, N.; Waku, T.; Yanagisawa, J.; Miyachi, H.; Shimizu, T. Structural basis for vitamin D receptor agonism by novel non-secosteroidal ligands. *FEBS Lett.* **2013**, *587*, 957–963.

(124) Masuno, H.; Ikura, T.; Morizono, D.; Orita, I.; Yamada, S.; Shimizu, M.; Ito, N. Crystal structures of complexes of vitamin D receptor ligand-binding domain with lithocholic acid derivatives. *J. Lipid Res.* **2013**, *54* (8), 2206–2213.

(125) Anami, Y.; Itoh, T.; Egawa, D.; Yoshimoto, N.; Yamamoto, K. A mixed population of antagonist and agonist binding conformers in a single crystal explains partial agonism against vitamin D receptor: active vitamin D analogues with 22R-alkyl group. *J. Med. Chem.* **2014**, *57* (10), 4351–4367.

(126) Watarai, Y.; Ishizawa, M.; Ikura, T.; Zacconi, F. C.; Uno, S.; Ito, N.; Mouriño, A.; Tokiwa, H.; Makishima, M.; Yamada, S. Synthesis, biological activities, and X-ray crystal structural analysis of 25-hydroxy-25(or 26)-adamantyl-17-[20(22),23-diyanyl]-21-norvitamin D compounds. *J. Med. Chem.* **2015**, *58* (24), 9510–9521.

(127) Anami, Y.; Sakamaki, Y.; Itoh, T.; Inaba, Y.; Nakabayashi, M.; Ikura, T.; Ito, N.; Yamamoto, K. Fine tuning of agonistic/antagonistic activity for vitamin D receptor by 22-alkyl chain length of ligands: 22S-Hexyl compound unexpectedly restored agonistic activity. *Bioorg. Med. Chem.* **2015**, *23* (22), 7274–7281.

(128) Anami, Y.; Shimizu, N.; Ekimoto, T.; Egawa, D.; Itoh, T.; Ikeguchi, M.; Yamamoto, K. Apo- and antagonist-binding structures of vitamin D receptor ligand-binding domain revealed by hybrid approach combining small-angle X-ray scattering and molecular dynamics. *J. Med. Chem.* **2016**, *59* (17), 7888–7900.

(129) Kato, A.; Itoh, T.; Anami, Y.; Egawa, D.; Yamamoto, K. Helix12-stabilization antagonist of vitamin D receptor. *Bioconjugate Chem.* **2016**, *27* (7), 1750–1761.

(130) Asano, L.; Waku, T.; Abe, R.; Kuwabara, N.; Ito, I.; Yanagisawa, J.; Nagasawa, K.; Shimizu, T. Regulation of the vitamin D receptor by vitamin D lactam derivatives. *FEBS Lett.* **2016**, *590* (18), 3270–3279.

(131) Egawa, D.; Itoh, T.; Kato, A.; Kataoka, S.; Anami, Y.; Yamamoto, K. SRC2–3 binds to vitamin D receptor with high sensitivity and strong affinity. *Bioorg. Med. Chem.* **2017**, *25* (2), 568–574.

(132) Kato, A.; Yamao, M.; Hashihara, Y.; Ishida, H.; Itoh, T.; Yamamoto, K. Vitamin D analogues with a p-hydroxyphenyl group at the C25 position: Crystal structure of vitamin D receptor ligand-binding domain complexed with the ligand explains the mechanism underlying full antagonistic action. *J. Med. Chem.* **2017**, *60* (20), 8394–8406.

(133) Otero, R.; Ishizawa, M.; Numoto, N.; Ikura, T.; Ito, N.; Tokiwa, H.; Mouriño, A.; Makishima, M.; Yamada, S. 25 S-Adamantyl-23-yne-26,27-dinor-1 α ,25-dihydroxyvitamin D₃: synthesis, tissue selective biological activities, and X-ray crystal structural analysis of its vitamin D receptor complex. *J. Med. Chem.* **2018**, *61* (15), 6658–6673.

(134) Yoshizawa, M.; Itoh, T.; Hori, T.; Kato, A.; Anami, Y.; Yoshimoto, N.; Yamamoto, K. Identification of the histidine residue in vitamin D receptor that covalently binds to electrophilic ligands. *J. Med. Chem.* **2018**, *61* (14), 6339–6349.

(135) Rochel, N.; Hourai, S.; Perez-Garcia, X.; Rumbo, A.; Mouriño, A.; Moras, D. Crystal structure of the vitamin D nuclear receptor ligand binding domain in complex with a locked side chain analog of calcitriol. *Arch. Biochem. Biophys.* **2007**, *460* (2), 172–176.

(136) Ciesielski, F.; Rochel, N.; Moras, D. Adaptability of the vitamin D nuclear receptor to the synthetic ligand Gemini: remodelling the LBP with one side chain rotation. *J. Steroid Biochem. Mol. Biol.* **2007**, *103*, 235–242.

(137) Eelen, G.; Valle, N.; Sato, Y.; Rochel, N.; Verlinden, L.; De Clercq, P.; Moras, D.; Bouillon, R.; Muñoz, A.; Verstuyf, A. Superagonistic fluorinated vitamin D₃ analogs stabilize helix 12 of the vitamin D receptor. *Chem. Biol.* **2008**, *15* (10), 1029–1034.

(138) Huet, T.; Maehr, H.; Lee, H.; Uskokovic, M.; Suh, N.; Moras, D.; Rochel, N. Structure function study of Gemini derivatives with two different side chains at C-20, Gemini-0072 and Gemini-0097. *MedChemComm* **2011**, *2*, 424–429.

(139) Fischer, J.; Wang, T. T.; Kaldre, D.; Rochel, N.; Moras, D.; White, J. H.; Gleason, J. L. Synthetically accessible non-secosteroidal hybrid molecules combining vitamin D receptor agonism and histone deacetylase inhibition. *Chem. Biol.* **2012**, *19* (8), 963–971.

(140) Huet, T.; Laverny, G.; Ciesielski, F.; Molnar, F.; Ramamoorthy, T. G.; Belorusova, A. Y.; Antony, P.; Potier, N.; Metzger, D.; Moras, D.; Rochel, N. A vitamin D receptor selectively activated by Gemini analogs reveals ligand dependent and independent effects. *Cell Rep.* **2015**, *10* (4), 516–526.

(141) Belorusova, A. Y.; Suh, N.; Lee, H. J.; So, J. Y.; Maehr, H.; Rochel, N. Structural analysis and biological activities of BXL0124, a Gemini analog of vitamin D. *J. Steroid Biochem. Mol. Biol.* **2017**, *173*, 69–74.

(142) Lin, Z.; Chen, H.; Belorusova, A. Y.; Bollinger, J. C.; Tang, E. K. Y.; Janjetovic, Z.; Kim, T. K.; Wu, Z.; Miller, D. D.; Slominski, A. T.; Postlethwaite, A. E.; Tuckey, R. C.; Rochel, N.; Li, W. 1 α ,20S-Dihydroxyvitamin D₃ interacts with vitamin D receptor: Crystal structure and route of chemical synthesis. *Sci. Rep.* **2017**, *7* (1), 10193.

(143) Belorusova, A. Y.; Martinez, A.; Gandara, Z.; Gomez, G.; Fall, Y.; Rochel, N. Structure-activity relationship study of vitamin D analogs with oxolane group in their side chain. *Eur. J. Med. Chem.* **2017**, *134*, 86–96.

(144) Lin, Z.; Marepally, S. R.; Goh, E. S. Y.; Cheng, C. Y. S.; Janjetovic, Z.; Kim, T. K.; Miller, D. D.; Postlethwaite, A. E.; Slominski, A. T.; Tuckey, R. C.; Peluso-Iltis, C.; Rochel, N.; Li, W. Investigation of 20S-hydroxyvitamin D₃ analogs and their 1 α -OH derivatives as potent vitamin D receptor agonists with anti-inflammatory activities. *Sci. Rep.* **2018**, *8* (1), 1478.